PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: A61K 38/00, 31/54, 31/535, 31/495, 31/445

(11) International Publication Number:

WO 99/08697

(43) International Publication Date:

25 February 1999 (25.02.99)

(21) International Application Number:

PCT/US98/17201

A1

(22) International Filing Date:

19 August 1998 (19.08.98)

(30) Priority Data:

60/056,135

19 August 1997 (19.08.97)

US

(71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): KAUFFMAN, Raymond, Francis [US/US]; 11420 Saint Andrews Lane, Carmel, IN 46032 (US). PALKOWITZ, Alan, David [US/US]; 10737 Kingsmill Drive, Carmel, IN 46032 (US).

(74) Agents: STRODE, Janelle, D. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: TREATMENT OF CONGESTIVE HEART FAILURE WITH GROWTH HORMONE SECRETAGOGUES

(57) Abstract

The invention provides methods for the modulation of cardiac function by the administration of a growth hormone secretagogue, wichch results in an increase in the levels of endogenous growth hormone. Also provided are methods for the treatment of congestive heart failure by the administration of a growth hormone secretagogue. Further provided are methods for the treatment of congestive heart failure by the administration of a growth hormone secretagogue in combination with a growth hormone releasing hormone, or in combination with an antihypertensive agent, diuretic, or other suitable agents.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE.	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

1

TITLE

TREATMENT OF CONGESTIVE HEART FAILURE WITH GROWTH HORMONE SECRETAGOGUES

5

10

15

20

25

30

Congestive heart failure (CHF; cardiac failure) is a condition in which weakened heart function exists together with a build-up of body fluid. Cardiac failure often occurs when cardiac output is insufficient to meet metabolic demands of the body, or when the heart cannot meet the demands of operating at increased levels of filling/diastolic pressure.

Congestive heart failure may be caused by many forms of heart disease. Common causes of congestive heart failure include: narrowing of the arteries supplying blood to the heart muscle (coronary heart disease); prior heart attack (myocardial infarction) resulting in scar tissue large enough to interfere with normal function of the heart; high blood pressure; heart valve disease due to past rheumatic fever or an abnormality present at birth; primary disease of the heart muscle itself (cardiomyopathy); defects in the heart present at birth (congenital heart disease) and infection of the heart valves and/ or muscle itself (endocarditis and/or myocarditis). Each of these disease processes can lead to congestive heart failure by reducing the strength of the heart muscle contraction, by limiting the ability of the heart's pumping chambers to fill with blood due to mechanical problems or impaired diastolic relaxation, or by filling the heart's chambers with too much blood.

Numerous compounds are known in the art to be useful for the prevention and treatment of congestive heart failure, including α -adrenergic antagonists, angiotensin II antagonists, angiotensin-converting enzyme (ACE) inhibitors,

2

 β -adrenergic antagonists, antihypertensives, calcium channel blockers, diuretics, potassium channel opening vasodilators, renin inhibitors, and serotonin antagonists.

Growth hormone, which is secreted from the pituitary gland, stimulates growth of all tissues of the body that are capable of growing. In addition, growth hormone is known to have the following basic effects on the metabolic processes of the body: (1) increased rate of protein synthesis in all cells of the body; (2) decreased rate of carbohydrate 10 utilization in cells of the body; and (3) increased mobilization of free fatty acids and use of fatty acids for energy. The cardiac effects of growth hormone have been reviewed, and it is suggested that growth hormone plays a role in the modulation of cardiac performance and the maintenance of normal cardiac structure and performance. 15 The effects of growth hormone on heart function is discussed in Sacca, et al., Endocrine Rev. 15: 555-573, 1994. Further, dramatic effects of growth hormone on heart failure have been observed clinically (Fazio et al., New England 20 Journal of Medicine, 334: 809-14, 1996).

A deficiency in growth hormone secretion can result in various medical disorders, depending on the age of onset. In children, the syndrome is characterized by short stature, with normal body proportions and reduced growth rate, also known as dwarfism. A deficiency in growth hormone secretion later in life may be characterized by excessive adiposity, reduced muscle mass, impaired exercise capacity, reduced body water, decreased bone mineral density, and psychological disorders. For example, a deficiency in growth hormone may result in myocardial dysfunction, in particular left ventricular diastolic dysfunction. Shahj, et al., Br. Heart J., 67, 92-96 (1992). A deficiency in growth hormone also has been suggested to be a factor in increased mortality from cardiovascular disease of adults

25

3

with growth-hormone deficiency. Rosen, Lancet, 336, 285-288 (1990).

Supplemental growth hormone administered for a four month period to growth hormone-deficient adults was reported to have no significant effect on left ventricular mass. Jorgenson, et al., Lancet, 1221-1225 (1989); Shajh, et al., Br. Heart J., 67, 92-96 (1992). However, when administered to growth-hormone deficient adults over a six month period, supplemental growth hormone increased left ventricular mass, 10 and provided favorable cardiovascular effects including increased cardiac output and glomerular filtration rate and reduced peripheral vascular resistance. Caldahl, et al., Clinical Endocrinology, 40, 393-400 (1994). In a patient suffering from growth hormone deficiency and poor cardiac function, supplemental growth hormone resulted in 15 improvement in myocardial contractility and cardiac output. Cuneo et al., Lancet, 1, 838-839 (1989). Supplemental growth hormone therapy in growth hormone deficient adults was found to improve resting and exercise cardiac function, but at the expense of ventricular hypertrophy. Fort, et 20 al., Circulation, 90, (4, Part 2) pg. I-610, abs. 3290 (1994). However, the effects of growth hormone in human heart failure patients without growth hormone deficiency have not been reported.

Certain compounds have been developed which stimulate the release of endogenous growth hormone. These compounds are also called growth hormone secretagogues. Growth hormone releasing peptides GHRP-6 and GHRP-1 (GRP-1 and GRP-6) are described in U.S. Patent No. 4,411,890 and PCT Patent Publications WO 89/07110 and WO 89/07111. Growth hormone releasing peptide GHRP-2 (GRP-2) is described in PCT Patent Publication WO 93/04081, and hexarelin is described in J. Endocrinol. Invest., 15 (Suppl 4), 45 (1992). Growth hormone releasing peptides (GHRPs) are also described in

25

4

Bowers et al., Endocrinology 114: 1537-45 1984. In addition, small organic molecules having growth hormone secretagogue activity have also been described in Jacks et al., Endocrinology 137: 5284-9, 1996.

Other compounds with growth hormone secretagogue 5 activity are discussed in United States Patent 3,239,345; United States Patent 4,036,979; United States Patent 4,411,890; United States Patent 5,206,235; United States Patent 5,248,841; United States Patent 5,310,737; United States Patent 5,310,017; European Patent Publication 10 144,230; European Patent Publication 513,974; Patent Cooperation Treaty Patent Publication WO 94/07486; Patent Cooperation Treaty Patent Publication WO 94/08583; Patent Cooperation Treaty Patent Publication WO 94/13696; Science, 15 260:1640-1643 (1993), Ann. Rep. Med. Chem., 28, 177-186 (1993); Bioorg. Med. Chem. Ltrs., 4 (22), 2709-2714 (1994); PNAS 92, 7001-7005 (July 1995); British Patent Application GB 2308064A, United States Serial Number 08/704,494, filed August 20, 1996, and United Sates Serial Number 08/700,206, filed August 20, 1996. Additional compounds with growth 20 hormone secretagogue activity are described herein.

Recent advances in heart failure therapy have demonstrated it is possible not only to alter the progression of heart failure, but also, in the case of antiadrenergic therapy with β -blockers, to partially reverse systolic dysfunction and ventricular remodeling of the failed myocardium (Eichhorn and Bristow, Circulation 94: 2285-96, 1996. In the case of Carvedilol®, these effects upon heart failure appear to result in dramatic reductions in mortality in patients (Packer et al., New England Journal of Medicine 334: 1349-55, 1996). Improved biological function of the cardiomyocyte appears to play an important role in the beneficial effects of β -blockade in experimental

25

5

heart failure (Tsutsui *et al.*, *J. Clin. Invest.* 93: 2639-2648, 1994). These observations indicate the reversible nature of congestive heart failure.

Although β -blockers and ACE inhibitors represent significant cornerstones for the treatment of heart failure, important medical issues remain concerning quality of life and survival. Thus, novel therapeutic approaches to congestive heart failure that improve survival rate as well as the indices of quality of life, and that are also convenient for both patient and physician remain an important unmet medical need.

6

Thus, it would be a significant contribution to the art to provide compounds and methods of using same that promote the secretion of growth hormone for use in the prevention, inhibition, or treatment of congestive heart failure as indicated herein.

The instant invention relates to methods for the modulation of cardiac function which comprise the administration of a compound having growth hormone secretagogue activity.

The present invention yet further relates to methods for the treatment or prevention of congestive heart failure, which method comprises administering to an animal in need of said treatment an effective amount of a compound having growth hormone secretagogue activity.

The present invention still further relates to methods for the treatment or prevention of congestive heart failure, which method comprises administering to an animal in need of said treatment an effective amount of GRP-2.

The present invention still further relates to methods for the treatment or prevention of congestive heart failure, which method comprises administering to an animal in need of said treatment an effective amount of a compound of formula I having growth hormone secretagogue activity, wherein a compound of formula I is represented by

25

5

10

15

```
wherein:
```

A is C_1 - C_6 alkyl, aryl, C_1 - C_6 alkylaryl, C_1-C_6 alkyl(0) C_1-C_6 alkylaryl, C_1-C_6 alkyl(S) C_1-C_6 alkylaryl, indoly1, indoliny1, thieny1, (C_1-C_6alky1) thieny1,

benzothienyl, benzofuranyl, naphthanyl, cyclohexyl, $(C_1 C_6$ alkyl)indolyl, $(C_1-C_6$ alkyl)benzothienyl,

 (C_1-C_6alkyl) naphthanyl, (C_1-C_6alkyl) benzofuranyl, and (C_1-C_6alkyl) cyclohexyl;

B is NH_2 , NHR_1 , C_1 - C_6 alky $1NH_2$, C_1 - C_6 alky $1NHR_1$, 10

 C_1 - C_6 alkylaryl NH_2 , C_1 - C_6 alkylaryl NHR_1 ,

 $C_1\text{-}C_6$ alkylcyclohexylNH2, $C_1\text{-}C_6$ alkylcyclohexylNHR1,

 R_1 -piperidin-3-yl(C_1 - C_6 alkyl),

 R_1 -piperidin-2-yl(C_1 - C_6 alkyl), R_1 -piperidin-4-yl(C_1 - C_6 alkyl),

 R_1 -quinolin-2-yl(C_1 - C_6 alkyl),

 R_1 -(2,4-dihydroquinolin-2-yl(C_1 - C_6 alkyl),

 R_1 -isoquinolin-2-yl(C_1 - C_6 alkyl), and

 R_1 -(2,4-dihydroisoquinolin-2-yl(C_1 - C_6 alkyl);

 R_1 is hydrogen, $C_1\text{-}C_6alkyl$, $C_1\text{-}C_6alkyl$ (OH), or


 C_1 - C_6 alkylidenyl(OH) R_2 ; 20

 R_2 is C_1 - C_6 alkyl, C_1 - C_6 alkenyl,

 C_1-C_6 alkyl(0) C_1-C_6 alkyl, $C(0)0-C_1-C_6$ alkyl, aryl, or C₁-C₆alkylaryl;

X is C_1 - C_6 alkylidenyl, O, S, NH, or $N(C_1$ - C_6 alkyl);

V is selected from the group consisting of 25



9

and

W is S, O, NH, or CH_2 ; 5 Y is N or CH; Z is N or CH; Y' is N or CH; Z' is N or CH; R₄ and R₅ are independently hydrogen, C₁-C6alkyl, 10 aryl, C_1 - C_6 alkylaryl, $C(0)O(C_1$ - C_6 alkyl), $C(0)N(C_1-C_6alkyl)_2$, or $C_1-C_6alkylCOR_7$; R₇ is hydrogen, C₁-C₆alkyl, pyrrolidinyl, piperidinyl, homoproline, or proline; D is hydrogen, C₁-C₆alkyl, 15 C_1-C_6 alkyl(0)(CO) C_1-C_6 alkyl, C_1-C_6 alkyl(0)(CO) $N(C_1-C_6$ alkyl)₂, C_1-C_6 alkylaryl, $C(0)R_6$, C_1-C_6 alkyl $(0)R_6$, C_1-C_6 alkyl(0H), C_1-C_6 $alkylC(0)R_6$, $C_1-C_6alkylR_6$, aryl, $(C_1-C_6alkyl)NHSO_2(C_1-C_6alkyl)$, (C_1-C_6alkyl) NHSO₂ (aryl); R_6 is H, C_1 - C_6 alkyl, aryl, naphthyl, 20 C_1 - C_6 alkylaryl, acetyl, NH_2 , $NH(C_1$ - C_6 alkyl), $NH(C_1-C_6alky1)O(C_1-C_6alky1)$, $NH(C_1-C_6alky1)S(C_1-C_6alky1)$, $NH(C_1-C_6alkylidenyl)OCH_3$, $NH(C_1-C_6alkyl)aryl$, $NH(C_3-C_6 \text{ cycloalkyl})$, $NH(C_1-C_6\text{alkyl})C(O)(C_1-C_6\text{alkyl})$, $NH(C_1-C_6alkyl)NH(C_1-C_6alkyl)$, $NH(C_1-C_6alkyl)NH(C_1-C_6alkylaryl)$, 25 $NHSO_2(C_1-C_6alkylaryl)$, $NH(C_1-C_6alky1)C(0)O(C_1-C_6alky1)$, NH (naphthyl), $N(C_1-C_6alkyl)_2$, $N(C_1-C_6alkyl)$ (aryl), $N(C_1-C_6alkyl)(C_1-C_6alkylaryl), O(C_1-C_6alkyl), O(aryl),$ $O(C_1-C_6alkylaryl)$, piperidinyl, piperidinyl- $C(0)NH(C_1-C_6alkyl)$, piperidinyl- $C(0)NH(C_1-C_6alkylaryl)$,

piperidinyl- $C(0)N(C_1-C_6alkyl)_2$,

10


piperidinyl-C(O)N(C₁-C₆alkyl)(aryl),
pyrrolidinyl, pyrrolidinyl C(O)NH(aryl)-,
pyrrolidinyl C(O)NH(C₁-C₆alkyl)-,
pyrrolidinyl C(O)NH(C₁-C₆alkylaryl)-,
pyrrolidinyl C(O)N(C₁-C₆alkyl)₂-,
pyrrolidinyl C(O)N(C₁-C₆alkyl)(aryl)-,
pyrrolinyl, morpholino, hexamethyleneimino,

pyrroliny1, morpholino, hexamethyleneimino, heptamethyleneimino, quinoliny1, 2,4-dihydroquinoliny1, 1,2,3,4-tetrahydroquinoliny1,

10 2,4-dihydroisoquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl,

- indolinyl, an amino acid selected from the group consisting of proline, homoproline, glycine, alanine, valine, leucine, isoleucine, tyrosine, tryptophan, phenylalanine, serine, threonine, asparagine, glutamic acid, aspartic acid,
- lysine, arginine, glutamine, histidine, cysteine, and methionine, or a nitrogen-containing heterocycle selected from the group consisting of

11



 $\label{eq:condition} E \mbox{ is hydrogen, C_1-C_6alkyl, $C(O)C_1-C_6alkyl$, aryl, $(aryl)C(O)NR_6$, $(aryl)(C_1-C_6alkyl)C(O)R_6$,}$

C₁-C₆alkylaryl, C(0)aryl, C₁-C₆ alkylC(0)aryl, naphthyl, C₁-C₆alkylnaphthyl, C(0)naphthyl, C₁-C₆alkylC(0)naphthyl, heteroaryl, C₁-C₆alkylheteroaryl, C(0)heteroaryl, C₁-C₆ alkylC(0)heteroaryl, indanyl,

12

 C_1 - C_6 alkylindanyl, C(0)indanyl, C_1 - C_6 alkylC(0)indanyl, cycloalkyl;

or D and E combine to form indanyl, fluorenyl, or cycloalkyl;

G is hydrogen, C_1 - C_6 alkyl, aryl, C_1 - C_6 alkylaryl, and C_1 - C_6 alkenyl;

J is hydrogen, $C_1\text{-}C_6alkyl,\ aryl,\ and <math display="inline">C_1\text{-}C_6alkylaryl;$

15

20

25

 $L is hydrogen, C_1-C_6alkyl, C(0)OC_1-C_6alkyl, aryl, \\ C_1-C_6alkylaryl, C(0)OC_1-C_6alkylaryl, C_1-C_6alkenyl, -F, and \\ -CN, C_1-C_6alkyl-OH, C_1-C_6alkyl-O-C_1-C_6alkyl, \\ C_1-C_6alkyl-C(0)R_6;$

or a pharmaceutically acceptable salt or solvate thereof.

The present invention also relates to pharmaceutical formulations containing a growth hormone secretagogue alone or in combination with additional therapeutic agents useful for the treatment or prevention of congestive heart failure.

Also provided are pharmaceutical formulations of GRP-2 alone or in combination with additional therapeutic agents useful for the treatment or prevention of congestive heart failure.

Further provided are pharmaceutical formulations of a compound of formula I alone or in combination with additional therapeutic agents useful for the treatment or prevention of congestive heart failure.

Figure 1 illustrates the differences in heart and lung weight observed between the administration of growth hormone and GRP-2 in the SHHF rat model of CHF.

Figure 2 illustrates pre-dosing levels of corticosterone in rats.

Figure 3 illustrates levels of corticosterone in rats treated with growth hormone or GRP-2 after one month.

Figure 4 illustrates levels of corticosterone in rats treated with growth hormone or GRP-2 at the end of the study.

5

20

30

13

PCT/US98/17201

Figure 5 illustrates pre-dosing levels of atrial natriuretic peptide (ANP) in rats.

Figure 6 illustrates levels of atrial natriuretic peptide (ANP) in rats treated with growth hormone or GRP-2 after one month.

Figure 7 illustrates levels of atrial natriuretic peptide (ANP) in rats treated with growth hormone or GRP-2 at the end of the study.

Figure 8 illustrates end systolic volume in rats 10 treated with growth hormone or GRP-2.

(Note: Data presented in Figures 1-7 is +/- standard error of means. Data presented in Figure 8 is +/- standard deviation.)

The term "growth hormone secretagogue" means any exogenously administered compound that directly or indirectly stimulates or increases the endogenous release of growth hormone in an animal. The growth hormone secretagogue may be peptidyl or nonpeptidyl in nature.

The growth hormone secretagogue may be used alone or in combination with other growth hormone secretagogues, or with other therapeutic agents known to be beneficial in the treatment of hypertension or the treatment or prevention of congestive heart failure, at least as discussed *supra*.

25 Administration of same may be concomitant or sequential.

Improving overall pulmonary function in a patient suffering from congestive heart failure is also included in the scope of the present invention. In addition, the administration of a growth hormone secretagogue is also useful for diminishing or preventing loss of body weight and enhancing recovery following congestive heart failure.

Administration of a growth hormone secretagogue more closely mimics biological functionality than the administration of growth hormone subcutaneously. For

10

15

20

25

30

35

14

PCT/US98/17201

example, bolus or acute administration of GHRPs leads to an acute, transient release of growth hormone from pituitary stores, but the chronic administration of same results in an enhanced, pulsatile release of growth hormone. The latter therapeutic approach more closely mimics the natural, physiological release of endogenous growth hormone. The terms and abbreviations used in the instant examples have their normal meanings unless otherwise designated. For example "°C" refers to degrees Celsius; "N" refers to normal or normality; "mmol" refers to millimole or millimoles; "g" refers to gram or grams; "ml" means milliliter or milliliters; "M" refers to molar or molarity; "MS" refers to mass spectrometry; "FDMS" refers to field desorption mass spectrometry; "UV" refers to ultraviolet spectroscopy; "IR" refers to infrared spectroscopy; and "NMR" refers to nuclear magnetic resonance spectroscopy.

As used herein, the term " C_1 - C_6 alkyl" refers to straight or branched, monovalent, saturated aliphatic chains of 1 to 6 carbon atoms and includes, but is not limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, pentyl, isopentyl, and hexyl. The term " C_1 - C_6 alkyl" includes within its definition the term " C_1 - C_4 alkyl".

As used herein, the term "cycloalkyl" refers to cyclized chains of 1 to 6 carbon atoms and includes, but is not limited to, cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

"Halo" represents chloro, fluoro, bromo or iodo. "C1-C6 alkoxy" represents a straight or branched alkyl

chain having from one to six carbon atoms attached to an oxygen atom. Typical C_1 - C_6 alkoxy groups include methoxy, ethoxy, propoxy, isopropoxy, butoxy, t-butoxy, pentoxy and the like. The term " C_1 - C_6 alkoxy" includes within its definition the term " C_1 - C_4 alkoxy".

"C2-C6 alkanoyl" represents a straight or branched alkyl chain having from one to five carbon atoms attached through a carbonyl moiety. Typical C2-C6 alkanoyl groups

25

30

35

15

PCT/US98/17201

include ethanoyl (also referred to as acetyl), propanoyl, isopropanoyl, butanoyl, t-butanoyl, pentanoyl, hexanoyl, and the like.

"C1-C6 alkylidenyl" refers to a straight or branched, divalent, saturated aliphatic chain of one to six carbon atoms and includes, but is not limited to, methylenyl, ethylenyl, propylenyl, isopropylenyl, butylenyl, isobutylenyl, t-butylenyl, pentylenyl, isopentylenyl, hexylenyl, and the like.

The term "aryl" represents an aromatic ring or rings including phenyl, napthyl, biphenyl, and aromatic residues of 5 to 7-membered rings with 1 to 4 heteroatoms (a "heteroaryl"), all of which may be optionally substituted with one or more substituents, including

15 C1-C6 alkyl, -OC1-C6 alkyl, -OCF3, amide, NHamide, carboxamide, sulfonamide, NHsulfonamide, imide, hydroxy, carboxy, nitro, chloro, fluoro, tri(chloro or fluoro)methyl, cyano, and the like. The aromatic ring may be attached at any carbon atom or heteroatom which affords a stable structure. 3,4-methylenedioxyphenyl is included here.

The term "heterocycle" represents a stable 5- to 7-membered monocyclic or 7- to 10-membered bicyclic heterocyclic ring which is saturated or unsaturated and which consists of carbon atoms and from 1 to 4 heteroatoms selected from the group consisting of nitrogen, oxygen or sulfur, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized and including a bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which affords a stable structure, and may be optionally substituted with one or more substituents selected from the group consisting of C1-C6 alkyl, -OC1-C6 alkyl, hydroxy, nitro, chloro, fluoro, or tri(chloro or fluoro)methyl, and the like.

25

30

PCT/US98/17201

The term "carboxy-protecting group" as used herein refers to substituents of the carboxy group commonly employed to block or protect the carboxy functionality while reacting other functional groups on the compound. Examples of such protecting groups include methyl, ethyl, pnitrobenzyl, p-methylbenzyl, p-methoxybenzyl, 3,4dimethoxybenzyl, 2,4-dimethoxybenzyl, 2,4,6trimethoxybenzyl, 2,4,6-trimethylbenzyl, pentamethylbenzyl, 3,4-methylene-dioxybenzyl, benzhydryl, 4,4'-dimethoxy-10 benzhydryl, 2,2',4,4'-tetramethoxybenzhydryl, t-butyl, tamyl, trityl, 4-methoxytrityl, 4,4'-dimethoxytrityl, 4, 4', 4"-trimethoxytrity1, 2-phenylprop-2-yl, trimethylsily1, tbutyldimethylsilyl, phenacyl, 2,2,2-trichloroethyl, 2-(di(nbuty1) methylsily1) ethyl, p-toluenesulfonylethyl, 4nitrobenzylsulfonylethyl, allyl, cinnamyl, 1-15 (trimethylsilylmethyl)prop-1-en-3-yl, and the like. A preferred carboxy-protecting group for the practice of the present invention is methyl or ethyl. Further examples of these groups may be found in E. Haslam, supra, at Chapter 5, 20 and T.W. Greene, et al., supra, at Chapter 5.

16

The term "amino-protecting group" as used herein refers to substituents of the amino group commonly employed to block or protect the amino functionality while reacting other functional groups on the compound. Examples of such amino-protecting groups include formyl, trityl, phthalimido, trichloroacetyl, chloroacetyl, bromoacetyl, iodoacetyl, and urethane-type blocking groups such as benzyloxycarbonyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl, 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, n-butoxycarbonyl, (NBoc) t-butoxycarbonyl,

35 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl,

17

2-phenylprop-2-yloxycarbonyl,

2-(p-toluy1)-prop-2-yloxycarbonyl, cyclopentanyloxycarbonyl,

1-methylcyclopentanyloxycarbonyl, cyclohexanyloxycarbonyl,

1-methylcyclohexanyloxycarbonyl,

5 2-methylcyclohexanyloxycarbonyl,

2-(4-toluylsulfonyl)-ethoxycarbonyl,

2-(methylsulfonyl)ethoxycarbonyl,

2-(triphenylphosphino)-ethoxycarbonyl,

fluorenylmethoxy-carbonyl (FMOC),

10 2-(trimethylsilyl)ethoxycarbonyl, allyloxycarbonyl,

1-(trimethylsilylmethyl)prop-1-enyloxycarbonyl,

5-benzisoxalylmethoxycarbonyl, 4-acetoxybenzyloxycarbonyl,

2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl,

cyclopropylmethoxycarbonyl, 4-(decyloxy)benzyloxycarbonyl,

isobornyloxycarbonyl, 1-piperidyloxycarbonyl, and the like; benzoylmethylsulfonyl group, 2-nitrophenylsulfenyl,

diphenylphosphine oxide and like amino-protecting groups.

The amino-protecting group employed is usually not critical so long as the derivatized amino group is stable to the condition of subsequent reactions on other positions of the intermediate molecule, and may be selectively removed at the appropriate point without disrupting the remainder of the molecule including any other amino-protecting groups. A preferred amino-protecting group for the practice of the present invention is t-butoxycarbonyl (NBoc). Further examples of groups referred to by the above terms are

described by E. Haslam, Protective Groups in Organic Chemistry, (J.G.W. McOmie, ed., 1973), at Chapter 2; and T.W. Greene and P.G.M. Wuts, Protective Groups in Organic

30 Synthesis (1991), at Chapter 7.

20

25

The term "leaving group" (Q) refers to a group of atoms that is displaced from a carbon atom by the attack of a nucleophile in a nucleophilic substitution reaction. Suitable leaving groups include bromo, chloro, and iodo,

35 benzenesulfonyloxy, methanesulfonyloxy, and

18

toluenesulfonyloxy. The term "leaving group" (Q) includes activating groups.

The term "activating group" as used herein refers a leaving group which, when taken with the carbonyl (-C=O) group to which it is attached, is more likely to take part in an acylation reaction than would be the case if the group were not present, as in the free acid. Such activating groups are well-known to those skilled in the art and may be, for example, succinimidoxy, phthalimidoxy,

10 benzotriazolyloxy, azido, or -O-CO-(C4-C7 alkyl).

15

The compounds used in the method of the present invention may have one or more asymmetric centers. As a consequence of these chiral centers, the compounds of the present invention occur as racemates, mixtures of enantiomers and as individual enantiomers, as well as diastereomers and mixtures of diastereomers. All asymmetric forms, individual isomers and combinations thereof, are within the scope of the present invention.

The terms "R" and "S" are used herein as commonly used
in organic chemistry to denote specific configuration of a
chiral center. The term "R" (rectus) refers to that
configuration of a chiral center with a clockwise
relationship of group priorities (highest to second lowest)
when viewed along the bond toward the lowest priority group.

The term "S" (sinister) refers to that configuration of a

The term "S" (sinister) refers to that configuration of a chiral center with a counterclockwise relationship of group priorities (highest to second lowest) when viewed along the bond toward the lowest priority group. The priority of groups is based upon their atomic number (in order of

decreasing atomic number). A partial list of priorities and a discussion of stereochemistry is contained in *Nomenclature* of Organic Compounds: Principles and Practice, (J.H. Fletcher, et al., eds., 1974) at pages 103-120.

In addition to the (R)-(S) system, the older D-L system is also used in this document to denote absolute configuration, especially with reference to amino acids. In

19

this system, a Fischer projection formula is oriented so that the number 1 carbon of the main chain is at the top. The prefix "D" is used to represent the absolute configuration of the isomer in which the functional (determining) group is on the right side of the carbon atom at the chiral center and "L", that of the isomer in which it is on the left.

In order to preferentially prepare one optical isomer over its enantiomer, a number of routes are available. As an example, a mixture of enantiomers may be prepared, and then the two enantiomers may be separated. A commonly employed method for the resolution of the racemic mixture (or mixture of enantiomers) into the individual enantiomers is to first convert the enantiomers to diastereomers by way of forming a salt with an optically active acid or base. These diastereomers may then be separated using differential solubility, fractional crystallization, chromatography, or the like. Further details regarding resolution of enantiomeric mixtures may be found in J. Jacques, et al., Enantiomers, Racemates, and Resolutions, (1991).

Preferred compounds of the present invention are compounds of formula I wherein:


A is

25

5

10

15



20

B is



J is H;

G is H;

5 X is NH;

10

V is



 \sum_{N}



21

E is

5

D is $-C(0)R_6$, where R_6 is 1-pyrrolidinyl, 1-piperidinyl, 4-methyl-1-piperidinyl, N,N-dimethyl,

or
$$N \longrightarrow F$$

10

L is H or CH_3 ; or a pharmaceutically acceptable salt or solvate thereof.

A preferred compound includes a compound of formula Id provided below:

Also preferred are compounds of formula IA and IB provided hereinabove.

10

15

20

25

During any of the following synthetic sequences it may be necessary or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by employing conventional protecting groups as described, supra.

The compounds of the present invention may be prepared by a number of routes, many of which are known to those of skill in the art. The particular order of steps to be employed in the synthesis of compounds of formula I is dependent upon the compound to be synthesized, the starting material employed, and the relative lability of the various substituted moieties. Examples of such synthetic routes may be found in Schemes I through IV provided below, as well as in the Examples.

One synthetic route to compounds of the present invention is provided in Scheme I below. The compounds of formula IV' and IV are commercially available, or may be prepared using techniques known in the art. A compound of formula IV may be prepared from a compound of formula IV' through an intermediate acid chloride prepared by standard methods using thionyl chloride or oxalyl chloride.

Treatment of the resulting acid chloride with a bromine source, such as N-bromosuccinimide, followed by quenching of

23

the acid chloride with ethanol, results in compounds of formula IV. It is to be understood that the bromine group on the compound of formula IV may in fact be any suitable leaving group (Q), as defined herein. This preparation is provided below in Scheme IA.

Scheme IA

15

20

25

30

wherein R is representative of E as defined in a compound of formula I above.

The starting material further includes compounds of formula V, which are commercially available, or may be routinely synthesized using techniques readily known in the art. Compounds of formula IV may be coupled with a compound of formula V (4-nitroimidazole) by methods known in the art to generate a compound of formula IIb'. Suitable agents to be employed in the coupling of these compounds include the treatment of a compound of formula IV with an organic or inorganic base, followed by reaction with the bromo compound of formula IV. Standard organic bases include trialkylamines, potassium hexamethyldisilazide, lithium hexamethyldisilazide, lithium diisopropylamide, potassium carbonate, and the like. Preferred for the practice of the present invention is sodium hydride or potassium carbonate in dimethylformamide. A compound of formula IIb' is then deprotected to provide a compound of formula IIb, using lithium hydroxide, although other deprotecting reagents may be employed in this reaction. Such deprotecting agents include standard saponification reagents such as sodium hydroxide, potassium hydroxide, and lithium hydroxide.

24

Substantially pure (R) enantiomers of compounds of formula IIb may also be synthesized by methods provided in U.S. 5,344,937 and 5,380,866, the disclosures of which are herein incorporated by reference.

A compound of formula IIb is then converted to the 5 corresponding amide under appropriate conditions with a compound of formula VI to generate a compound of formula In general, amidation of primary or secondary amines of formula VI may be accomplished by a number of methods known in the art in which activation of the acid to form a 10 better leaving group is the initial step. Suitable activating agents for this are also known in the art and include dicyclohexycarbodiimide (DCC), 1-(3dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) with hydroxybenzotriazole (HOBT), oxalyl chloride, thionyl 15 chloride, PyBOP® (benzotriazol-1-yloxytripyrrolidinephosphonium hexafluorophosphate), and the Preferred for the practice of the present invention is hydroxybenzotriazole (HOBT). The nitro group on the resulting compound of formula IIa may then be reduced to an 20 amino group using any suitable means, employing a suitable reducing agent. Preferred for the practice of the present invention is a catalytic reduction employing hydrogen and 5% palladium on carbon. A compound of formula II is produced by this reduction reaction. 25

The preferred reaction temperature range employed in these reactions is between -40 and 150 °C, and the most preferred range is between 10 and 40 °C. These reactions may be conveniently carried out *in situ*, without isolation of the particular compound after its preparation.

Examples of these reactions are provided below in Scheme I.

PCT/US98/17201

Scheme I

5 wherein R is representative of E as previously defined, and R_2R_1N is R_6 as previously defined.

26

A second portion of the overall synthesis of compounds of formula I is provided in Scheme II below. Representative starting material for this synthesis is a compound of formula IIIb', which is a chemically-protected form of the amino acid O-serine. By chemically-protected it is meant that both the amino- and carboxy- functional groups have been suitably protected in order to facilitate further reactions with this molecule. Such protection reactions are known to those of skill in the art, and may be applied to other suitable starting materials. Intermediates of formula IIIb' are commercially available, or may be prepared by standard syntheses of amino acids. Such syntheses are well known to persons of ordinary skill in the art and are described, for example, in Chemistry and Biochemistry of Amino Acids, (G.C. Chapman ed., 1985). The protected amino group may be specifically deprotected using trifluoroacetic acid and methylene chloride to allow for further reactions with this amino functional group. This deprotection reaction results in a compound of formula IIIb.

10

15

20

25

30

A compound of formula IIIb may then be N-acylated with an amino-protected compound of formula X to produce a compound of formula IIIa'. Suitable activating agents for this N-acylation reaction are known in the art and include DCC, HOBT, EDC, and oxalyl chloride. Preferred for the practice of the present invention is HOBT. Compounds of formula X are commercially available, or are readily prepared from suitable available starting materials. The protected carboxy group on the compound of formula IIIa' is then selectively deprotected, typically using lithium hydroxide, to generate a compound of formula III. Compounds of formula III in which the starting material IIIb' is 2-Nboc-amino-pentanoic acid methyl ester may also be prepared by the route described in Scheme II.

A compound of formula III is then coupled with a 35 compound prepared from the reduction of IIb' with hydrogen and a palladium catalyst employing a coupling reaction to

27

generate a compound of formula Ia. Again, typical reagents for this N-acylation are known in the art, and include DCC and HOBT, which is the preferred method of coupling employed in the practice of the present invention. A compound of formula Ia is then selectively deprotected at the carboxy group, coupled at this site with a compound of formula VI, and then further deprotected at the amino group to generate a compound of formula Ia. Suitable agents for these deprotection and coupling reactions are discussed, infra, and are known in the art. Compounds of formula Ia are encompassed by formula I, and are pharmaceutically active.

The preferred reaction temperature range employed in these reactions is between -40 and 150 °C, and the most preferred range is between 10 and 40 °C. These reactions may be conveniently carried out *in situ*, without isolation of the particular compound after its preparation.

10

15

20


Alternatively, compounds of formula IIa can be coupled with compounds of formula III to provide intermediates which can be deprotected to give compounds of formula Ia.

Representative reactions are provided below in Scheme II.

28

Sch me II

III



30

wherein R is E as previously defined, and R_2R_1N is R_6 as previously defined.

An alternative synthetic scheme is provided in Scheme III below. A compound of formula VII

- 5 (5-nitrobenzimidazole) is commercially available, or may be conveniently prepared using reactions known in the art. A compound of formula VII is coupled with a compound of formula IV in an alkylation reaction, using coupling agents as discussed, infra. A compound of formula VIII' is
- produced in which the carboxy functional group is protected. This protecting group is then removed as previously discussed, typically using lithium hydroxide, followed by coupling with a compound of formula XII. The nitro group on the resulting compound of formula VIII is then reduced,
- followed by coupling with a compound of formula III. The resulting compound of formula Ib' is then deprotected to provide a compound of formula Ib. Compounds of formula Ib are encompassed by formula I, and are pharmaceutically active. These reactions are provided below in Scheme III.

PCT/US98/17201 WO 99/08697

31

Scheme III

Ph

CO₂Me

Ib'

ö

Ιb

ĆO₂Me

32

A still further representative synthesis of compounds of formula I is provided below in Scheme IV. Starting materials of formula IX (3-amino-nitrobenzene) are commercially available. Initially, a compound of formula IX is coupled with a compound of formula IV by means discussed previously. The resulting compound of formula XI' is then deprotected, followed by coupling with a compound of formula XII to provide a compound of formula XI. A compound of formula XI is then reduced and further coupled in an N-acylation reaction with a compound of formula III. The resulting compound of formula Ic' is then deprotected to result in a compound of formula Ic. Conditions for these reactions have been discussed previously. Compounds of formula Ic are encompassed by formula I, and are pharmaceutically active.

10

Ic

Scheme IV

Ic'

34

In addition to the Schemes described hereinabove, an enantiospecific protocol for the preparation of the compounds of formula I may be employed. Typically, a synthetic reaction design which maintains the chiral center present in the starting material in a desired orientation is chosen. The preferred reaction schemes are those that generally produce compounds in which greater than 95 percent of the product is the desired enantiomer. In Scheme V below, R-substituted phenyl is representative of the E substituents as provided in compounds of formula I above. Further, the -methylene-X moiety is representative of the A substituents in compounds of formula I above.

Scheme V

36

The following discussion is directed to the reactions provided in Scheme V. Specifically, the reactions of compounds of formula I, II, and III are as provided in the discussion of Scheme I hereinabove.

5

10

30

A compound of formula IV may be prepared by the alkylation of a compound of formula III by standard methods using a base, such as sodium hydride, followed by treatment with an electrophile, such as methyl iodide. Preferred bases for this reaction include sodium-, lithium-, or potassium hexamethyldisilazide, lithium diisopropylamide, and sodium hydride. Preferred methylating agents include methyl halides or any methyl group with a suitable substituted leaving group such as toslyate, mesylate, and the like.

A compound of formula V may be prepared by hydrolysis of a compound of formula IV using standard saponfication conditions known in the art. Suitable reagents for this transformation include sodium hydroxide or lithium hydroxide. The resulting carboxylic acid may be converted into the acid chloride by standard methods using thionyl chloride or, preferably, oxalyl chloride. The acid chloride may then be reacted with the lithium salt of a chiral auxiliary, such as (4R, 5S)-(+)-4-methyl-5-phenyl-2-oxazolidinone, to provide compounds of formula V and VI, which are readily separable by silica gel chromatography.

A compound of formula VII may be prepared by the removal of the chiral auxiliary under basic conditions, such as lithium hydroxide. Other reagents known in the art for removing oxazolidinone-type chiral auxiliaries may be used for this transformation. These include lithium hydroxide/hydroperoxide conditions, reduction/oxidation protocols, alkyl sulfur displacements, and transaminations.

A compound of formula VIII may be prepared from a compound of formula VII by standard methods known in the

37

art. Formation of the acid chloride using oxalyl or thionyl chloride followed by reaction with a suitable substituted amine (NR $_2$) provide compounds of formula VIII.

A compound of formula IX may be prepared by the reduction of a compound of formula VIII using hydrogen with 5 palladium on carbon. Other methods known in the art which may be employed for the reduction of the nitro group include the use of tin(II)chloride, iron in an acidic solution, ferrous sulfate and aqueous alkali, activated alumina, and sodium sulfite. The resulting 4-amino imidazole compound of 10 formula VIIa is then reacted directly with the appropriate dipeptide acid (a compound of formula IIX) under standard peptide coupling conditions involving formation of the active ester of the dipeptide followed by reaction with amine VIIa. Conditions suitable for amide formation include 15 DCC, EDC, with HOBT. A compound of formula IIX may be prepared from the methyl ester of unnatural D-amino acids such as D-benzyloxyserine, D-tryptophan, and D-2-amino-5phenyl-pentanoic acid and the like which are known in the art. Standard coupling protocols involving formation of the 20 active ester of the amino acid using DCC/HOBt followed by reaction with N-Boc-aminoisobutyric acid provide dipeptide acids of formula IIX.

The Boc protecting group of a compound of formula IX may be removed under standard acidic conditions such as hydrochloric acid in acetic acid or ethyl acetate, trifluoroacetic acid, tetramethyliodosilane, aluminum chloride, sulfuric acid in dioxane, and methanesulfonic acid.

25

30

An additional method of preparing diastereomeric compounds of formula I involves the use of a chromatographic column which employs a chiral phase. An example of such a preparation may be found in Examples Part 6 as provided hereinbelow.

38

Preferred for the practice of the present invention are those compounds of formula I wherein the indicated stereochemistry is (R,R) at the two chiral centers. An example of this preferred stereochemistry is provided by compounds of formula IA and IB as provided hereinabove.

5

Two additional Schemes for providing chiral intermediates are provided hereinbelow as Schemes VA and VB. As described in Scheme VA, optically pure aryl glycine amino acids may be protected at the amino position by reaction 10 with a suitable protecting group, such as Boc. Reaction of the Boc protected intermediate with a standard methylating agent, such as methyl iodide, may provide the corresponding phenolic methyl ether. The carboxamide may be prepared by coupling with an amine, such as dimethylamine, pyrrolidine, or 4-methyl piperidine, using standard coupling techniques. 15 Preferred coupling agents for the invention are diethy cyanophosphorane (DECP), triethylamine and the amine at O°C. The Boc protecting group may be removed under standard acidic conditions, with trifluoroacetic acid being preferred. The desired 4-nitroimidazole compounds can be 20 prepared by reaction of the free amine with 1, 4dinitroimidazole to give optically pure compounds, as determined by chiral HPLC. Such chiral intermediates can be processed as described in Schemes I and II to provide diastereomerically pure products. For example, the chiral 25 nitroimidazoles described in Scheme VA or VB may be reduced under standard conditions, such as hydrogenation with a palladium catalyst, to provide the corresponding chiral amino intermediate II. Such intermediates may be subsequently coupled with compounds of type III as 30 previously described to provide a chiral intermediate which can be deprotected to give diastereomerically pure compounds of formula Ia.

39

Scheme VA

CHIRAL SYNTHESIS of D-Phenylglycine Imidazole Subunit

Scheme VB

Chiral Synthesis of L-Phenylglycine Imidazole

Chiral HPLC>97%ee

An additional approach and corresponding synthetic scheme for the preparation of compounds of the instant invention is provided below in Scheme VI:

41

Scheme VI

42

Pharmaceutically active compounds of formula I include at least compounds of formula IA, IB, Id, and Ia' as described herein.

Compounds of formula I may be conveniently screened for growth hormone secretagogue activity. A typical assay may employ pituitary cells established in culture, followed by a challenge with the various compounds of formula I, and the levels of growth hormone determined accordingly. Growth hormone levels may be calculated using various radioimmunoassay techniques known to those of skill in the art. Screening of compounds for growth hormone secretagogue activity may conveniently be scaled up for high throughput screening.

The invention further encompasses methods employing the pharmaceutically acceptable salts of the compounds defined by formula I. Although generally neutral, a compound of this invention can possess a sufficiently acidic, a sufficiently basic, or both functional groups, and accordingly react with any of a number of inorganic bases, and inorganic and organic acids, to form a pharmaceutically acceptable salt.

15

20

25

30

. 35

The term "pharmaceutically acceptable salt" as used herein refers to salts of the compounds of formula I which are substantially non-toxic to living organisms. Typical pharmaceutically acceptable salts include those salts prepared by reaction of the compounds of the present invention with a pharmaceutically acceptable mineral or organic acid or an inorganic base. Such salts are known as acid addition and base addition salts.

Acids commonly employed to form acid addition salts are inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as p-toluenesulfonic, methanesulfonic acid, oxalic acid,

43

p-bromophenylsulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like. Examples of such pharmaceutically acceptable salts are the sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caproate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, 10 chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, γ-hydroxybutyrate, glycollate, tartrate, methanesulfonate, propanesulfonate, 15 naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate, mesylate, and the like. Preferred pharmaceutically acceptable acid addition salts are those formed with mineral acids such as hydrochloric acid and hydrobromic acid, and those formed with organic acids such as maleic acid and 20 methanesulfonic acid.

Salts of amine groups may also comprise quaternary ammonium salts in which the amino nitrogen carries a suitable organic group such as an alkyl, alkenyl, alkynyl, or aralkyl moiety.

25

30

35

Base addition salts include those derived from inorganic bases, such as ammonium or alkali or alkaline earth metal hydroxides, carbonates, bicarbonates, and the like. Such bases useful in preparing the salts of this invention thus include sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, potassium bicarbonate, calcium hydroxide, calcium carbonate, and the like. The potassium and sodium salt forms are particularly preferred.

It should be recognized that the particular counterion forming a part of any salt of this invention is not of a

44

critical nature, so long as the salt as a whole is pharmacologically acceptable and as long as the counterion does not contribute undesired qualities to the salt as a whole.

This invention further encompasses methods employing pharmaceutically acceptable solvates of the compounds of Formula I. Many of the formula I compounds can combine with solvents such as water, methanol, ethanol and acetonitrile to form pharmaceutically acceptable solvates such as the corresponding hydrate, methanolate, ethanolate and acetonitrilate.

10

15

20

30

This invention also encompasses methods employing the pharmaceutically acceptable prodrugs of the compounds of formula I. A prodrug is a drug which has been chemically modified and may be biologically inactive at its site of action, but which may be degraded or modified by one or more enzymatic or other *in vivo* processes to the parent bioactive form. This prodrug should have a different pharmacokinetic profile than the parent, enabling easier absorption across the mucosal epithelium, better salt formation or solubility, or improved systemic stability (an increase in plasma halflife, for example).

Typically, such chemical modifications include:

- 1) ester or amide derivatives which may be cleaved by 25 esterases or lipases;
 - 2) peptides which may be recognized by specific or nonspecific proteases; or
 - 3) derivatives that accumulate at a site of action through membrane selection of a prodrug form or a modified prodrug form; or any combination of 1 to 3, supra. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in H, Bundgaard, Design of Prodrugs, (1985).

Suitable dosing ranges of compounds of formula I include 0.01 $\mu g/kg/day$ to 60 mg/kg/day. Representative

45

pharmaceutical formulations containing compounds of formula I are provided below.

Pharmaceutical formulations containing a growth hormone secretagogue as described herein may also include an effective amount of agents having therapeutic activity as α -adrenergic antagonists, angiotensin II antagonists, angiotensin-converting enzyme (ACE) inhibitors, β -adrenergic antagonists, antihypertensives, calcium channel blockers, diuretics, potassium channel opening vasodilators, renin inhibitors, and serotonin antagonists.

5

10

15

20

25

30

In addition, growth hormone secretagogues may be administered along with cardiac stimulants such as dobutamine and xamoterol and phosphodiesterase inhibitors including amrinone and milrinone.

Optionally, the active ingredient of the pharmaceutical formulations can include an anabolic agent in addition to at least one growth hormone secretagogue or another composition which exhibits another activity, for example an antibiotic growth promoting agent, or a corticosteroid employed to minimize the catabolic side effects, or another compound which enhances efficacy and minimizes side effects. Growth promoting and anabolic agents include TRH, diethylstilbesterol, estrogens, β -antagonists, theophilline, anabolic steroids, enkephalins, E series prostaglandins, retinoic acid, zeranol, and sulbenox.

The different forms of these additional therapeutic agents available as well as the various utilities associated with same and the applicable dosing regimens are well known to those of skill in the art. The method of administration of each additional therapeutic agent is consistent with that known in the art.

Also included are compounds of formula I employed in the manufacture of a medicament for use in treating or preventing congestive heart failure.

As used herein, the term "effective amount" means an amount of compound of the instant invention which is capable

46

of inhibiting, alleviating, ameliorating, treating, or preventing further symptoms in mammals, including humans, suffering from congestive heart failure.

By "pharmaceutically acceptable formulation" it is meant that the carrier, diluent, excipients and salt must be 5 compatible with the active ingredient (a compound of formula I) of the formulation, and not be deleterious to the recipient thereof. Pharmaceutical formulations can be prepared by procedures known in the art. For example, the 10 compounds of this invention can be formulated with common excipients, diluents, or carriers, and formed into tablets, capsules, and the like. Examples of excipients, diluents, and carriers that are suitable for such formulations include the following: fillers and extenders such as starch, 15 sugars, mannitol, and silicic derivatives; binding agents such as carboxymethyl cellulose and other cellulose derivatives, alginates, gelatin, and polyvinyl pyrrolidone; moisturizing agents such as glycerol; disintegrating agents such as agar agar, calcium carbonate, and sodium 20 bicarbonate; agents for retarding dissolution such as paraffin; resorption accelerators such as quaternary ammonium compounds; surface active agents such as cetyl alcohol, glycerol monostearate; adsorptive carriers such as kaolin and bentonite; and lubricants such as talc, calcium 25 and magnesium stearate and solid polyethylene glycols. Final pharmaceutical forms may be: pills, tablets, powders, lozenges, syrups, aerosols, saches, cachets, elixirs, suspensions, emulsions, ointments, suppositories, sterile injectable solutions, or sterile packaged powders, and the

Additionally, the compounds of this invention are well suited to formulation as sustained release dosage forms. The formulations can also be so constituted that they release the active ingredient only or preferably in a particular part of the intestinal tract, possibly over a period of time. Such formulations would involve coatings,

like, depending on the type of excipient used.

30

47

envelopes, or protective matrices which may be made from polymeric substances or waxes.

The particular dosage of a compound required to treat, inhibit, or prevent the symptoms and/or disease of

5 congestive heart failure in a mammal, including humans, according to this invention will depend upon the particular disease, symptoms, and severity. Dosage, routes of administration, and frequency of dosing is best decided by the attending physician. Generally, accepted and effective doses will be from 15mg to 1000mg, and more typically from 15mg to 80mg. Such dosages will be administered to a patient in need of treatment from one to three times each day or as often as needed for efficacy.

48

The formulations which follow are given for purposes of illustration and are not intended to be limiting in any way. The total active ingredients in such formulations comprises from 0.1% to 99.9% by weight of the formulation. The term "active ingredient" means a growth hormone secretagogue, including GRP-2 or a compound of formula I.

49

Formulation 1

Hard gelatin capsules containing the following ingredients are prepared:

5

		Quantity
	Ingredient	(mg/capsule)
	Active Ingredient	30.0
10	Starch	305.0
	Magnesium stearate	5.0

The above ingredients are mixed and filled into hard gelatin capsules in 340 mg quantities.

50

Formulation 2

A tablet formula is prepared using the ingredients below:

5

		Quantity
	Ingredient	(mg/tablet)
	Active Ingredient	25.0
10	Cellulose, microcrystalline	200.0
	Colloidal silicon dioxide	10.0
15	Stearic acid	5.0

The components are blended and compressed to form tablets, each weighing 240 mg.

51

Formulation 3

A dry powder inhaler formulation is prepared containing the following components:

	<u>Ingredient</u>	Weight %
	Active Ingredient	5
10	Lactose	95

The active mixture is mixed with the lactose and the mixture is added to a dry powder inhaling appliance.

Formulation 4

Tablets, each containing 30 mg of active ingredient, are prepared as follows:

		Quantity
	Ingredient	(mg/tablet)
10	Active Ingredient	30.0 mg
	Starch	45.0 mg
1.5	Microcrystalline cellulose	35.0 mg
15	Polyvinylpyrrolidone (as 10% solution in water)	4.0 mg
	Sodium carboxymethyl starch	4.5 mg
20	Magnesium stearate	0.5 mg
	Talc	1.0 mg
25	Total	120 mg

The active ingredient, starch and cellulose are passed through a No. 20 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders, which are then passed through a 16 mesh U.S. sieve. The granules so produced are dried at 50-60°C and passed through a 16 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate, and talc, previously passed through a No. 30 mesh U.S. sieve, are then added to the

53

granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 120 mg.

Formulation 5

5

Capsules, each containing 40 mg of medicament are made as follows:

		Quantity
10	Ingredient	(mg/capsule)
	Active Ingredient	40.0 mg
	Starch	109.0 mg
15	Magnesium stearate	1.0 mg
	Total	150.0 mg

The active ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 20 mesh U.S. sieve, and filled into hard gelatin capsules in 150 mg quantities.

54

Formulation 6

5 Suppositories, each containing 25 mg of active ingredient are made as follows:

	Ingredient	Amount
	Active Ingredient	25 mg
10		
	Saturated fatty acid glycerides to	2,000 mg

The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid
15 glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2.0 g capacity and allowed to cool.

55

Formulation 7

Suspensions, each containing 50 mg of medicament per 5.0 ml dose are made as follows:

5

25

	Ingredient Active Ingredient	Amount 50.0 mg
10	Xanthan gum	4.0 mg
10	Sodium carboxymethyl cellulose (11%) Microcrystalline cellulose (89%)	50.0 mg
15	Sucrose	1.75 g
13	Sodium benzoate	10.0 mg
	Flavor and Color	q.v.
20	Purified water to	5.0 ml

The medicament, sucrose and xanthan gum are blended, passed through a No. 10 mesh U.S. sieve, and then mixed with a previously made solution of the microcrystalline cellulose and sodium carboxymethyl cellulose in water. The sodium benzoate, flavor, and color are diluted with some of the water and added with stirring. Sufficient water is then added to produce the required volume.

56

Formulation 8

Capsules, each containing 15 mg of medicament, are made as follows:

5

		Quantity
	Ingredient	(mg/capsule)
	Active Ingredient	15.0 mg
10	Starch	407.0 mg
	Magnesium stearate	3.0 mg
	Total	425.0 mg

15

The active ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 20 mesh U.S. sieve, and filled into hard gelatin capsules in 425 mg quantities.

20

Formulation 9

An intravenous formulation may be prepared as follows:

<u>Ingredient</u>	Quantity
Active Ingredient	250.0 mg
Isotonic saline	1000 ml

57

Formulation 10

A topical formulation may be prepared as follows:

5

	Ingredient Active Ingredient	Quantity 1-10 g
10	Emulsifying Wax	30 g
	Liquid Paraffin	20 g
	White Soft Paraffin	to 100 g

The white soft paraffin is heated until molten. The liquid paraffin and emulsifying wax are incorporated and stirred until dissolved. The active ingredient is added and stirring is continued until dispersed. The mixture is then cooled until solid.

58

Formulation 11

Sublingual or buccal tablets, each containing 10 mg of active ingredient, may be prepared as follows:

5

25

30

_		
		Quantity
	Ingredient	Per Tablet
	Active Ingredient	10.0 mg
10	Glycerol	210.5 mg
	Water	143.0 mg
15	Sodium Citrate	4.5 mg
	Polyvinyl Alcohol	26.5 mg
	Polyvinylpyrrolidone	15.5 mg
20	Total	410.0 mg

The glycerol, water, sodium citrate, polyvinyl alcohol, and polyvinylpyrrolidone are admixed together by continuous stirring and maintaining the temperature at about 90°C.

When the polymers have gone into solution, the solution is cooled to about 50-55°C and the medicament is slowly admixed. The homogenous mixture is poured into forms made of an inert material to produce a drug-containing diffusion matrix having a thickness of about 2-4 mm. This diffusion matrix is then cut to form individual tablets having the appropriate size.

Another formulation employed in the methods of the present invention employs transdermal delivery devices or patches. Such transdermal patches may be used to provide

59

continuous or discontinuous infusion of the compounds of the present invention in controlled amounts. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art. See, for example, U.S. Patent 5,023,252, the disclosure of which is herein incorporated by reference. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

Frequently, it will be desirable or necessary to introduce the pharmaceutical composition to the brain, either directly or indirectly. Direct techniques usually involve placement of a drug delivery catheter into the host's ventricular system to bypass the blood-brain barrier. One such implantable delivery system, used for the transport of biological factors to specific anatomical regions of the body, is described in U.S. Patent 5,011,472, the disclosure of which is herein incorporated by reference.

10

15

20

25

Indirect techniques, which are generally preferred, usually involve formulating the compositions to provide for drug latentiation by the conversion of hydrophilic drugs into lipid-soluble drugs or prodrugs. Latentiation is generally achieved through blocking of the hydroxy, carbonyl, sulfate, and primary amine groups present on the drug to render the drug more lipid soluble and amenable to transportation across the blood-brain barrier.

Alternatively, the delivery of hydrophilic drugs may be enhanced by intra-arterial infusion of hypertonic solutions which can transiently open the blood-brain barrier.

The following Examples are provided to better elucidate the practice of the present invention and should not be interpreted in any way as to limit the scope of same. Those skilled in the art will recognize that various modifications may be made while not departing from the spirit and scope of

60

the invention. All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains.

61

EXAMPLES PART 1

Preparation 1

5

10

15

20

To a solution of boc-(OBz)-D-Ser-OH (25.0 g, 84.7 mmol) stirring in anhydrous N,N-dimethylformamide (500 mL) at room temperature was added sodium bicarbonate (14.2 g, 169 mmol) followed by methyl iodide (26.4 mL, 424 mmol). After 18 h, the reaction mixture was concentrated to approximately 100 mL. Ethyl acetate was added and the mixture washed with aqueous sodium bicarbonate and brine. The organic extract was dried and concentrated to give the desired compound (25 g, 96%) as a light yellow oil: ¹H NMR (300 MHz, CDCl3) d 1.45 (s, 9H), 3.70 (m, 1H), 3.75 (s, 3H), 3.85 (m, 1H), 4.50 (m, 3H), 7.30 (m, 5H); MS (FD) m/e 310; Anal. calc'd for C16H23NO5: C, 62.12; H, 7.49; N, 4.53. Found: C, 62.31; H, 7.49; N, 4.43.

Preparation 2

25

To a solution of a compound of Preparation 1 (5.0 g, 16 mmol) stirring in dichloromethane (25 mL) and anisole (1 mL) at 0 $^{\circ}$ C was added trifluoroacetic acid. After 4 h at room

62

temperature, saturated sodium bicarbonate solution was added and the mixture extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sufate, and concentrated. The crude product was used in the next step without further purification.

Preparation 3

$$\bigcap_{\text{MeO }_2\text{C}} \bigcap_{\text{O}} \bigcap_{\text{NHBoc}}$$

10

5

To a solution of a compound of Preparation 2 (65.4 mmol), boc- α -aminoisobutyric acid (13.2 g, 65.4 mmol), 1hydroxybenzotriazole (8.8 g, 65.4 mmol), and N,Ndiisopropylethylamine (22.8 mL, 130.7 mmol) stirring in dichloromethane (500 mL) at 0 °C was added 1-(3-15 dimethylaminopropyl)-3-ethylcarbodiimide (12.3 g, 71.9 mmol). After 18 h, ethyl acetate and saturated ammonium chloride were added and the mixture extracted with ammonium chloride, sodium bicarbonate, and brine. The organic 20 extracts were dried over sodium sulfate and concentated. Purification by silica gel chromatography (25% ethyl acetate/hexanes) yielded the desired compound (21.6 g, 83%) as a white solid: ${}^{1}H$ NMR (300 MHz, CDCl₃) d 1.39 (s, 9H), 1.48 (s, 6H), 3.62 (dd, J = 3.4, 9.1 Hz, 1H), 3.70 (s, 3H), 3.85 (dd, J = 3.4, 9.1 Hz, 1H), 4.48 (dd, J = 12.5, 22.7 Hz,25 2H), 4.75 (m, 1H), 4.92 (s, 1H), 7.11 (d, J = 8.6 Hz, 1H), 7.35 (m, 5H); MS (FD) m/e 395; Anal. calc'd for C20H30N2O6: C, 60.90; H, 7.67; N, 7.10. Found: C, 61.02; H, 7.78; N, 7.10.

63

Preparation 4

5 To a solution of a compound of Preparation 3 (5.30 g, 13.4) stirring in dioxane (100 mL)/water (50 mL) at room temperature was added lithium hydroxide (2.80 g, 67.3 mmol). After 18 h, water was added and the solution concentrated. The resulting mixture was extracted with diethyl ether. Sodium chloride was added to the aqueous layer and the pH 10 adjusted to 3.5 with 1 N HCl. The resulting mixture was extracted with ethyl acetate and the combined organic extracts dried over sodium sulfate then concentrated to yield the title compound (4.40 g, 86%) as a white foam: 1H NMR (300 MHz, CDCl3) d 1.39 (s, 9H), 1.45 (s, 3H), 1.47 (s, 15 3H), 3.68 (m, 1H), 3.95 (m, 1H), 4.54 (s, 2H), 4.70 (m, 1H), 5.51 (bs, 1H), 7.18 (d, J = 9.1 Hz, 1H), 7.25 (m, 5H), 9.90(bs, 1H); MS (FD) m/e 381; Anal. calc'd for C19H28N2O6: C, 59.99; H, 7.42; N, 7.36. Found: C, 59.74; H, 7.26; N, 7.30.

20

Preparation 5

To a solution of α -bromophenylacetic acid (100 g, 466 mmol) stirring in absolute ethanol (500 mL) at room temperature was added p-toluenesulfonic acid monohydrate (10 g, 53 mmol). This solution was heated to reflux and, after

64

8 h, concentrated to dryness. The resulting residue was dissolved in ethyl acetate, washed with saturated aqueous sodium bicarbonate, brine, dried over sodium sulfate, filtered, and concentrated to yield 77 g (68 %) of the desired product as an orange oil: ¹H-NMR is consistent with structure; MS (FD) 241.9, 243.9.

Preparation 6

10

15

20

25

To a slurry of sodium hydride (13.6 g of a 60% dispersion in mineral oil, 341 mmol) stirring in N,Ndimethylformamide (240 mL) was carefully added 4nitroimidazole (38.6 g, 341 mmol) such that the temperature during the addition was maintained below 40°C. This resulting slurry was stirred for 1 h and then cooled to 5°C. To this mixture was slowly added BX8-MEZ-148 (76 g, 310 mmol) at a rate such that the reaction temperature was maintained below 20°C. After 4 h, the reaction was concentrated and subsequently extracted with ethyl acetate. The combined organic extracts were filtered, washed with water, brine, dried over sodium sulfate, filtered and concentrated. The resulting residue was purified by silica gel chromatography (methanol/chloroform gradient) to yield the 60.1 g (70%) of the desired product as a white solid: H-NMR is consistent with structure; MS (FD) 275 (M+); Anal. Calc'd. for: C, 56.73; H, 4.73; N, 15.27. Found: C, 56.48; H, 4.78; N, 15.08.

65

Preparation 7

5 To a suspension of 5% Pd/C (0.85 g) and a compound of Preparation 6 (2.13 g, 7.21 mmol) stirring in dioxane (50 mL) at room temperature was added hydrogen (g) (35 psi) on a Parr apparatus. After 4 h, the mixture was purged with nitrogen, celite added, and the solution filtered through a 10 pad of celite. To the resulting filtrate, under nitrogen atmosphere, was added a compound of Preparation 4 (2.74 g, 7.21 mmol), 1-hydroxybenzotriazole (0.97 g, 7.21 mmol), N,Ndiisopropylethylamine (2.5 mL, 14.4 mmol), and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (1.36 g, 7.93 mmol). After 18 hours, ethyl acetate was added and the 15 mixture washed with saturated aqueous ammonium chloride, saturated aqueous sodium bicarbonate, and brine. organic extract was dried over sodium sulfate and concentrated. Purification by silica gel chromatography (5% methanol/dichloromethane) yielded the title compound (1.25 20 g, 29 %) as a yellow foam: 1 H NMR (300 MHz, CDCl₃) d 1.30 (t, J = 6.9 Hz, 3H), 1.40 (s, 9H), 1.42 (s, 3H), 1.51 (s, 33H), 3.60 (dd, J = 5.1, 9.7 Hz, 1H), 4.05 (m, 1H), 4.28 (m, 2H), 4.54 (dd, J = 14.08, 26.3 Hz, 2H), 4.62 (m, 1H), 5.08(bs, 1H), 5.82 (s, 1H), $7.12 \cdot (d, J = 11.5 Hz, 1H), <math>7.35$ (m, 25 12H), 9.75 (bs, 1H); MS (FD) m/e 607; Anal. calc'd for C32H41N5O7: C, 63.29; H, 6.80; N, 11.52. Found: C, 63.07; H, 6.81; N, 11.74.

66

Preparation 8

To a solution of a compound of Preparation 7 (5.3 g, 8.75) stirring in dioxane (50 mL)/water (25 mL) at room temperature was added lithium hydroxide (0.73 g, 17.50 mmol). After 20 min, water was added and the reaction concentrated to approximately 30 mL. The resulting mixture was extracted with diethyl ether and the aqueous layer saturated with sodium chloride then adjusted to pH 3.5 with 1 N HCl. The mixture was extracted with ethyl acetate and the combined organic extracts dried over sodium sulfate and concentrated to yield the title compound (4.90 g, 97%) as a light orange foam: ¹H NMR (300 MHz, CDCl3) d; MS (FD) m/e; Anal. calc'd for: C,; H,; N, Found: C,; H,; N,

Preparation 9

67

To a solution of a compound of Preparation 8 (2.09 g, 3.61 mmol), pyrrolidine (0.30 mL, 3.61 mmol), and 1hydroxybenzotriazole (0.54 g, 3.97 mmol) stirring in anhydrous DMF (50 mL) at 0 °C was added 1,3-dicyclohexyl carbodiimide (0.82 g, 3.97 mmol). After 18 hours at room temperature, the reaction was concentrated, dissolved in. dichloromethane, filtered, and concentrated. Purification by silica gel chromatography (5% methanol/dichloromethane) yielded the title compound (1.74 g, 76%) as a light orange solid: ¹H NMR (300 MHz, CDCl₃) d 1.41 (s, 9H), 1.43 (s, 3H), 1.52 (s, 3H), 2.88 (m, 4H), 3.42 (m, 1H), 3.50 (m, 4H), 4.08 (m, 1H), 4.55 (dd, J = 14.9, 27.4 Hz, 2H), 4.70 (m,1H), 4.96 (d, J = 4.0 Hz, 1H), 5.86 (s, 1H), 7.15 (d, J =6.9 Hz, 1H), 7.35 (m, 12H), 9.28 (bs, 1H); MS (FD) m/e 632; Anal. calc'd for C34H44N6O6: C, 64.54; H, 7.01; N, 13.28. Found: C, 63.48; H, 6.95; N, 12.19.

Example 1

20

25

10

15

To a solution of a compound of Preparation 9 (1.00 g, 1.58 mmol) and anisole (0.3 mL) stirring in anhydrous dichloromethane (12 mL) at 0 $^{\circ}$ C was added trifluoroacetic acid (3 mL) and the reaction mixture warmed to room temperature. After 4 h, the dichloromethane was removed in

68

vacuo and excess diethyl ether added. After 20 min, the reaction mixture was filtered to yield the title compound (1.02 g, 85%) as a white solid: ¹H NMR (300 MHz, CDCl₃) d 1.60 (s, 6H), 1.90 (m, 4H), 3.08 (m, 1H), 3.58 (m, 3H), 3.88 (m, 2H), 4.52 (m, 2H), 4.72 (m, 1H), 6.10 (m, 2H), 7.25 (m, 6H), 7.46 (m, 5H), 7.70 (m 1H), 8.00 (m, 1H), 8.40 (m, 1H), 11.15 (m, 1H); MS (FD) m/e 532 (M-2TFA); Anal. calc'd for C₃₃H₃₈F₆N₆O₈: C, 52.10; H, 5.03; N, 11.05. Found: C, 51.54; H, 5.25; N, 11.21.

10

5

Preparation 10

To a slurry of $d, 1-\alpha$ -amino-4-phenylbutyric acid (20.0) 15 q, 111 mmol) stirring in 3N sulfuric acid (200 mL) at 0°C was added finely ground potassium bromide (48 g, 403 mmol). This slurry was cooled to -10°C, then a solution of sodium nitrite (11.0 g, 160 mmol in water (75 mL)) was added 20 dropwise. The resulting solution was stirred for 4 h while slowly warming to ambient temperature. The resulting precipitate was filtered to give 20.0 g of a yellow solid. To a solution of the yellow solid (18.8 g, 80 mmol) in absolute ethanol (400 mL) was added p-toluenesulfonic acid monohydrate (4.6 g, 24 mmol). This solution was refluxed 25 for 4 h, filtered and concentrated. The resulting residue was purified by silica gel chromatography (ethyl acetate/hexanes gradient) to give 7.2 g (24%) of the desired product as a clear oil. H-NMR is consistent with structure; 30 MS (FD) 269, 27

69

Preparation 11

$$EtO \bigvee_{O}^{NO_2} \bigvee_{O}^{N}$$

5 To a slurry of sodium hydride (1.0 g of a 60% dispersion in mineral oil, 24 mmol) stirring in N, Ndimethylformamide (200 mL) at ambient temperature was carefully added a solution of 4-nitroimidazole (5.7 g, 20 mmol). This mixture was cooled to 0 $^{\circ}$ C and a solution of a compound of Preparation 10 (15.2 g, 60 mmol) in N, N-10 dimethylformamide (10 mL) was added. After 16 h, the mixture was slowly warmed to ambient temperature, concentrated, and the resulting residue extracted with chloroform. The combined organic extracts were washed with water, brine, dried over sodium sulfate, filtered and 15 concentrated. The resulting residue was purified by silica gel chromatography (chloroform) to give 5.0 g (82%) of the desired product as a clear oil. H-NMR is consistent with structure; MS (FD) 303 (M+); Anal. Calc'd for: C, 59.40; H, 5.65; N, 13.85. Found: C, 59.73; H, 5.71; N, 13.40. 20

70

Preparation 12

5

To a solution of a compound of Preparation 11 (4.24 g, 14 mmol) stirring in tetrahydrofuran (30 mL) and ethanol (30 mL) at room temperature was added 2N NaOH (35 mL, 70 mmol). After 1 h, this mixture was treated with 5N HCl until pH = 2.5. Ethyl acetate (30 mL) and water (30 mL) were added and the resulting solution was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated to give 3.8 g (98%) of the desired product as a yellow oil: ¹H-NMR is consistent with structure; MS (FD) 276 (M+).

71

Preparation 13

To a solution of a compound of Preparation 12 (3.8 g, 14 mmol), 1-proline methylester (1.8 g, 14 mmol) and 1-hydroxybenzotriazole hydrate (2.1 g, 15 mmol) stirring in N,N-dimethylformamide (150 mL) at room temperature was added 1,3-dicyclohexylcarbodiimide (3.2 g, 15.4 mmol). After 16 h, the mixture was concentrated and the resulting residue partitioned between ethyl acetate and water. The combined organic extracts were washed with water, brine, dried over sodium sulfate, and concentrated. The resulting orange oil was purified by silica gel chromatography

15 (methanol/chloroform gradients) to give 3.8 g (70%) of the desired product as a yellow oil: 1H-NMR is consistent with

structure; MS (FD) 386.2 (M+).

72

Preparation 14

5 To a slurry of 10% Pd/C in dioxane (10 mL) was added a solution of a compound of Preparation 13 (2.4 g, 6.2 mmol) in dioxane (100 mL). The mixture was then treated with hydrogen gas (40 psi) on a Parr apparatus. After 5 h, an amount of 10% Pd/C (0.5 g) in dioxane (10 mL) was added. The mixture was hydrogenated for 4 h then carefully filtered 10 through celite. To the resulting filtrate was added a compound of Preparation 4 (2.4 g, 6.2 mmol), 1hydroxybenzotriazole hydrate (0.92 g, 6.8 mmol), followed by 1,3-dicyclohexylcarbodiimide (1.4 g, 6.8 mmol). After 16 h, 15 the reaction was concentrated and the resulting residue extracted with ethyl acetate. The combined organic extracts were washed with saturated aqueous sodium bicarbonate, brine, dried over sodium sulfate, filtered, and concentrated. Purification silica gel chromatography 20 (methanol/chloroform gradient) gave 2.2 g (50%) of the desired product as a tan foam: 1H-NMR is consistent with structure; MS (FD) 718.7 (M+); Anal Calc'd for: C, 63.49; H, 7.01; N, 11.69. Found: C, 63.30; H, 6.91; N, 11.84.

73

Example 2

$$\begin{array}{c|c}
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & & \\
& & &$$

5

15

20

To a solution of a compound of Preparation 14 (2.1 g, 3.0 mmol) stirring in dichloromethane (25 mL) was added trifluoroacetic acid (8 mL, 104 mmol). After one h, water (25 mL) was added and the solution was quenched carefully with sodium carbonate, then extracted with chloroform. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. To a solution of the resulting residue in diethyl ether (40 mL) was added a saturated solution of HCl in diethyl ether (40 mL). The resulting slurry was concentrated to dryness to yield 1.6 g (80%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 618.3 (M+); Anal. Calc'd. for: C, 57.31; H, 6.41; N, 12.15. Found: C, 57.52; H, 6.19; N, 12.04. IR (KBr) 2954, 1743, 1656, 1559, 1496, 1453 cm⁻¹.

74

Preparation 15

Reaction of d,1-α-phenylalanine (20.0 g, 120 mmol), potassium bromide (48 g, 400 mmol), sodium nitrite (11.0g, 160 mmol), water (75 mL), 3N sulfuric acid (200 mL), ptoluenesulfonic acid monohydrate (5.7 g, 30 mmol) and absolute ethanol (500 mL) according to Preparation 10 gave 18.0 g (70%) of the desired product as a colorless oil: ¹H-NMR is consistent with structure; MS (FD) 256, 258.

Preparation 16

15

20

Reaction of a compound of Preparation 15 (15.22 g, 60 mmol), sodium hydride (2.84 g of a 60% dispersion in mineral oil, 72 mmol), 4-nitroimidazole (8.1 g, 72 mmol) in N,N-dimethylformamide (400 mL) according to Preparation 11 gave 9.5 g (55%) of the desired product as a yellow foam: ¹H-NMR is consistent with structure; MS (FD) 289.1 (M+); Anal. Calc'd. for: C, 58.13; H, 5.23; N, 14.53. Found: C, 58.40; H, 5.17; N 14.24.

25

75

Preparation 17

5

Reaction of a compound of Preparation 16 (3.3 g, 12.0 mmmol), 2N NaOH (30 mL, 60 mmol) in ethyl acetate (30 mL)/ethanol (30 mL) according Preparation 12 gave 2.85 g (90%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 262 (M+); Anal. Calc'd. for: C, 55.17; H, 4.24; N, 16.09. Found: C, 55.14; H, 4.24; N, 15.94.

Preparation 18

15

Reaction of a compound of Preparation 17 (2.8 g, 11.0 mmol), 1-proline methylester (1.4 g, 11.0 mmol), 1
20 hydroxybenzotriazole hydrate (1.63 g, 12.1 mmol), and 1,3-dicyclohexylcarbodiimide (2.5 g, 12.1 mmol) in N,N-dimethylformamide (150 mL according to Preparation 13 gave

PCT/US98/17201

3.2 g (70.4%) of the desired product as a white solid: $^{1}\text{H-}$ NMR is consistent with structure; MS (FD) 372 (M+).

Preparation 19

5

Reaction of a compound of Preparation 18 (0.6 g, 1.6 mmol), 5% Pd/C (0.66 g) in ethyl acetate (50 mL), ethanol (50 mL) and dichloromethane (4 mL), a compound of Preparation 4 (0.46 g, 1.2 mmol), 1-hydroxybenzotriazole hydrate (0.18 g, 1.3 mmol) and 1,3-dicyclohexylcarbodiimide (0.27 g, 1.3 mmol) in N,N-dimethylformamide (100 mL)

15 according to Preparation 14 gave 0.29 g (34%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 704.5 (M+).

77

Exampl 3

Reaction of a compound of Preparation 19 (0.23 g, 0.33 mmol), trifluoroacetic acid (4.0 mL, 24 mmol) in dichloromethane (12 mL), followed by treatment with HCl/ethyl acetate solution (40 mL), according to Example 2 gave 0.17 g (77%) of the desired product as a white foam:

10

1 H-NMR is consistent with structure; MS (FD) 604 (M+); Anal. Calc'd for: C, 56.72; H, 6.25; N, 12.40. Found: C, 56.53; H, 6.31; N, 12.19. IR (KBr) 2931.09, 1743.64, 1653.48, 1533.67, 1453.73 (cm⁻¹).

15


Preparation 20

Reaction of ethylbromoacetate (4.9 mL, 44 mmol), 4-nitroimidazole (5.00 g, 44 mmol) and potassium carbonate (12.2 g, 88 mmol) at ambient temperature in N,N-

78

dimethylformamide (50 mL) according to Preparation 3 from Examples Part 2A gave 7.77 g (88%) of the desired product as an orange solid: ¹H-NMR was consistent with structure; MS (FD) 199 (M+); Anal. Calc'd for: C, 42.21; H, 4.55; N, 21.10. Found: C, 42.51; H, 4.66; N, 21.24.

Preparation 21




10

15

Reaction of a compound of Preparation 20 (2.00 g, 10.0 mmol) and 2N NaOH (30 mL, 60 mmol) in tetrahydrofuran (5 mL) and ethanol (5 mL) according Preparation 12 gave 1.3 g (76%) of the desired product as a tan solid which is carried on without further purification.

Preparation 22



20

25

Reaction of a compound of Preparation 21 (1.20 g, 7.0 mmol), 1-proline methylester hydrochloride (1.27 g, 8.4 mmol), 1-hydroxybenzotriazole hydrate (1.04 g, 8.4 mmol), triethylamine (1.95 mL, 14.0 mmol) and 1,3-dicyclohexylcarbodiimide (1.6 g, 8.4 mmol) in N,N-

dimethylformamide according to Preparation 13 gave 0.6 g

79

(30%) of the desired compound as a tan semi-solid: ¹H-NMR is consistent with structure; MS (FD) 282 (M+).

Preparation 23

5

Hydrogenation of a compound of Preparation 22 (0.47 mg, 1.7 mmol) and 5 % Pd-C (0.15 g) in ethyl acetate (20 mL)/ethanol (20 mL) followed by treatment with 1-hydroxybenzotriazole hydrate (225 mg, 1.7 mmol), 1,3-dicyclohexylcarbodiimide (340 mg, 1.7 mmol) and 368979 (633 mg, 1.7 mmol) according to Preparation 14 gave 0.45 g (39%) of the desired product: ¹H-NMR is consistent with structure; MS (FD) 614 (M+).

80

Example 4

5

Reaction of a compound of Preparation 23 (0.40 g, 0.65 mmol) and trifluoroacetic acid (5 mL, 64 mmol) in dichloromethane (20 mL) according to Example 2 gave 0.22 g (67%) of the desired product as an off-white solid: ¹H-NMR is consistent with structure; MS (FD) 514 (M+); Anal. Calc'd for: C, 58.35; H, 6.66; N, 16.33. Found: C, 58.25; H, 6.40; N, 16.16.

Preparation 24

15

Preparation 1 gave 3.9 g (65%) of the desired product: 'H-

Reaction of 5-nitroindole (3.0 g, 18.5 mmol), α -bromophenylacetic acid ethylester (4.5 g, 18.5 mmol), and sodium hydride (0.8 g, 20 mmol, 60% dispersion in mineral oil) in N,N-dimethylformamide (75 mL) according to

81

NMR is consistent with structure; MS (FD) 324 M+; Anal. Calc'd for: C, 66.66; H, 4.97; N, 8.64. Found: C, 66.80; H, 5.11; N, 8.81.

Preparation 25

5

NO 2 HO NO

Reaction of a compound of Preparation 24 (2.0 g, 6.2 mmol) and 2N NaOH (50 mL, 100 mmol) in tetrahydrofuran (10 mL)/ethanol (8 mL) according to Preparation 12 gave 1.4 g (76%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 296 (M+); Anal. Calc'd for: C, 64.86; H, 4.08; N, 9.45. Found: C, 64.60; H, 4.14; N, 9.29.

82

Pr paration 26

Reaction of a compound of Preparation 25 (1.0 g, 5.7 mmol), 1-hydroxybenzotriazole hydrate (0.85 g, 6.3 mmol), 1-proline methylester hydrochloride (1.03 g, 6.3 mmol), triethylamine (1.6 mL, 11.4 mmol) and 1,3-dicyclohexylcarbodiimide (1.3 g, 6.3 mmol) in N,N-dimethylformamide (25 mL) according to Preparation 13 gave 1.35 g (58%) of the desired product as yellow solid: ¹H-NMR is consistent with structure; MS (FD) 407 (M+); Anal. Calc'd for: C, 64.86; H, 5.20; N, 10.31. Found: C, 65.20; H, 5.50; N, 10.10.

15

83

Pr paration 27

Hydrogenation of a compound of Preparation 26 (0.41 g, 1.0 mmol) with 5 % Pd-C (0.08 g) in ethanol (25 mL)/ethyl acetate (25 mL) followed by treatment with 1-hydroxybenzotriazole hydrate (0.15 g, 1.1 mmol), 1,3-dicyclohexylcarbodiimide (0.23 g, 1.1 mmol) and 368979 (0.42 g, 1.1 mmol) according to Preparation 14 gave 0.38 g (51%) of the desired product: ¹H-NMR is consistent with structure; MS (FD) 739.7 (M+)

84

Example 5

Reaction of a compound of Preparation 27 (0.38 g, 0.51 mmol) and trifluoroacetic acid (2 mL, 26 mmol) in dichloromethane (10 mL) according to Example 2 gave 0.125 g (38%) of the desired product: ¹H-NMR is consistent with structure; MS (FD) 639 (M+); Anal. Calc'd for 1 H₂O: C, 65.74; H, 6.59; N, 10.65. Found: C, 65.75; H, 6.42; N, 10.98.

Preparation 28

15

20

To a solution of a compound of Preparation 6 (27 g, 98 mmol) stirring in tetrahydrofuran (60 mL) and absolute ethanol (60 mL) at ambient temperature was added 2N NaOH (250 mL, 500 mmol). After 3.5 h, the mixture was washed

85

with diethyl ether and the organic extract subsequently washed with water. The combined aqueous extracts were acidified and the resulting mixture extracted with ethyl acetate. The combined organic extracts were washed once with brine, dried over sodium sulfate, filtered, and concentrated to give 24.2 g (75%) of the desired product as a tan solid: ¹H-NMR was consistent with structure; MS (FD) 246.9 (M+); Anal. Calc'd for: C, 53.44; H, 3.67; N, 17.00. Found: C, 53.71; H, 3.67; N, 16.83. mp = 218-221°C.

10

Preparation 29

15 To a slurry of a compound of Preparation 28 (8.15 g, 33 mmol) stirring in dichloromethane (100 mL) was added oxalyl chloride (11.5 mL, 130 mmol) and N, N-dimethylformamide (2 drops). After 90 min at ambient temperature, the mixture was concentrated and the residue was dissolved in 20 dichloromethane (40 mL). The resulting solution was added a N, N-diisopropylethylamine (6.5 mL, 360 mmol) and 1prolinemethylester (3.9 g, 20 mmol) in dichloromethane (4 mL). After 2 h at ambient temperature, the mixture was extracted with ethyl acetate and the combined organic 25 extracts washed with water, brine, dried over sodium sulfate, filtered, and concentrated. The resulting residue was purified by silica gel chromatography (ethyl acetate/hexanes) to give 10.7 g (71%) of the desired product

86

as tan foam: ¹H-NMR is consistent with structure; Anal. Calc'd for: C, 56.98; H, 5.06; N, 15.63. Found: C, 56.75; H, 5.14; N, 15.44. Mp, 103-111°C.

Preparation 30

5

To a slurry of 5% Pd/C (0.28 g) in ethyl acetate (30 10 mL) was added a solution of a compound of Preparation 29 (1.0 g, 2.8 mmol) in ethanol (100 mL). The mixture was hydrogenated at 40 psi on a Parr apparatus. After 25 min, additional 5% Pd/C (0.5 g) was added and the mixture susbsequently hydrogenated for 45 min, then filtered through 15 celite and concentrated. To a slurry of the resulting residue in N,N-dimethylformamide (100 mL) was added boc-dbenzyloxyserine (0.62 g, 2.1 mmol), 1-hydroxybenzotriazole hydrate(0.31 g, 2.3 mmol) followed by 1,3dicyclohexylcarbodiimide (0.48 g, 2.3 mmol). After 48 h, the 20 mixture was filtered and concentrated and the residue purified by radial chromatography (silica gel, methanol/chloroform gradient). The resulting product was dissolved in ethyl acetate and washed with water, dried over sodium sulfate, filtered, and concentrated to give 0.5 g (30%) of the desired product as a tan foam: 'H-NMR is 25 consistent with structure; MS (FD) 605 (M+).

87

Preparation 31

5

10

To a solution of a compound of Preparation 30 (3.1 g, 5.1 mmol) stirring in methanol (200 mL) at room temperature was added 5N HCl (51.0 mmol). After 16 h, the residue was partitioned between ethyl acetate and water and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated to give 2.1 g (81%) of the desired compound as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 506 (M+); Anal. Calc'd. for: C, 64.14; H, 6.18; N, 13.85.

15 Found: C, 63.92, H, 6.18; N, 13.56.

88

Preparation 32

To a solution of a compound of Preparation 31 (2.1 g, 4.2 mmol) stirring in N,N-dimethylformamide (200 mL) was added Boc-α-aminoisobutyric acid (0.85 g, 4.2 mmol), 1-hydroxybenzotriazole hydrate(0.62 g, 4.6 mmol). After 16 h, mixture was concentrated to dryness and the resulting residue extracted with ethyl acetate. The combined organic extracts were washed with water, brine, dried over sodium sulfate, filtered, and concentrated. Purification by silica gel chromatography (methanol/chloroform) gave 2.3 g (80%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 690 (M+).

89

Example 6

2.5 HCl

To a solution of the compound of Preparation 32 (1.75 g, 2.5 mmol) stirring in dichloromethane (190 mL) was added trifluoroacetic acid (63 mL, 780 mmol). After 1 h, the mixture was poured carefully into saturated aqueous sodium bicarbonate and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated. The residue was dissolved in ethyl acetate (250 mL) and subsequently treated with a saturated solution of HCl in ethyl acetate (100 mL). The resulting mixture was concentrated to dryness,

15 triturated with diethyl ether, and filtered to give 0.6 g (38%) of the desired product as a tan solid: ¹H-NMR is consistent with structure; MS (FD) 590 (M+); Anal. Calc'd for: C, 54.60; H, 5.92; N, 12.33. Found: C, 54.47; H, 5.72; N, 12.16. IR (KBr) 3164, 3030, 2978, 2952, 2878, 1743,

20 1664, 1531, 1456, 1436, 1498, 1197, 1179 cm⁻¹.

90

Preparation 33

The optically enriched S-isomer was isolated by selective crystallization (ethyl acetate/hexanes) of a compound of Preparation 29 to give 1.3 g of the desired isomer: ¹H-NMR is consistent with structure; MS (FD) 358 (M+); Anal. Calc'd for: C, 56.98; H, 5.06; N, 15.63. Found: C, 57.22; H, 4.87; N, 15.34. mp = 114-118°C.

Preparation 34

15

Hydrogenation of a compound of Preparation 29 (1.0 g, 2.8 mmol) and 5% Pd/C (0.756 g) in absolute ethanol (20 mL)/ethyl acetate (20 mL), followed by treatment of the resulting mixture with boc-d-benzyloxyserine (0.83 g, 2.8 mmol), 1-hydroxybenzotriazole hydrate (0.42 g, 3.4 mmol) and 1,3-dicyclohexylcarbodiimide (0.64 g, 3.1 mmol according to Preparation 1 gave 0.69 g (41%) of the desired product as a

91

crystalline solid. Purification by silica gel chromatography (methanol/chloroform) followed by recrystalization from ethyl acetate: ¹H-NMR is consistent with structure; MS (FD) 605 (M+); Anal. Calc'd. for: C, 63.46; H, 6.49; N, 11.56. Found: C, 63.61; H, 6.31; N, 11.38; mp = 184-186°C.

Preparation 35

10

Reaction of a compound of Preparation 34 (0.61 g, 1.0 mmol) and trifluoroacetic acid (1.7 mL, 22 mmol) in dichloromethane (40 mL) according to Preparation 1 gave 0.5 g (100%) of the desired product as a foam: ¹H-NMR is consistent with structure; MS (FD) 506 (M+); mp = 55-60°C.

92

Preparation 36

Reaction of a compound of Preparation 35 (0.5 g, 1 mmol), 1-hydroxybenzotriazole hydrate (0.15 g, 1.1 mmol) and 1,3-dicyclohexylcarbodiimide (0.23 g, 1.1 mmol) in N,N-dimethylformamide (15 mL) according to Preparation 32 gave 0.69 g (100%) of the desired product as a foam: ¹H-NMR is consistent with structure; MS (FD) 690.2 (M+); mp = 81-84°C.

10

Example 7

Reaction of a compound of Preparation 36 (0.595 g, 0.95 mmol) and trifluoroacetic acid (0.7 mL, 9.0 mmol) in dichloromethane (25 mL) according to Preparation 1 gave 0.37 g (75%) of the desired product as a solid: ¹H-NMR was consistent with structure; MS (FD) 590 (M+); Anal. Calc'd for: C, 63.04; H, 6.48; N, 14.23. Found: C, 62.98; H, 6.59; N, 14.01. Mp, 156-159°C.

93

Preparation 37

5

Reaction of a compound of Preparation 29 (2.63 g, 8.0 mmol), boc-1-benzyloxyserine (2.4 g, 8.0 mmol), 1-hydroxybenzotriazole hydrate (1.2 g, 8.8 mmol), 1,3-dicyclohexylcarbodiimide (1.8 g, 8.8 mmol) in N,N-dimethylformamide (250 mL) gave 2.4 g (50%) of the desired product as tan foam: ¹H-NMR is consistent with structure; MS (FD) 605 (M+).

94

Preparation 38

Reaction of a compound of Preparation 37 (2.3 g, 3.8 mmol), trifluoroacetic acid (35 mL, 45 mmol) in dichloromethane (90 mL) gave 1.4 g (74%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 506 (M+).

10 Preparation 39

Reaction of a compound of Preparation 38 (1.1 g, 2.2 mmol), boc-α-aminoisobutyric acid (0.45 g, 2.2 mmol), 1-hydroxybenzotriazole hydrate (0.33 g, 2.4 mmol) and 1,3-dicyclohexylcarbodiimide (0.5 g, 2.4 mmol) in N,N-dimethylformamide (100 mL) gave 0.84 g (55%) of the desired

95

product as a tan foam: $^{1}H-NMR$ is consistent with structure; MS (FD) 690 (M+).

Example 8

5

Reaction of a compound of Preparation 39 (0.7 g, 1.0 mmol), trifluoroacetic acid (25 mL, 320 mmol) in

10 dichloromethane, and followed by dissolution in ethyl acetate (100 mL) and treatment with ethyl acetate saturated with HCl (100 mL) yielded 0.29 g (44%) of the desired compound as a white solid: ¹H-NMR is consistent with structure; MS (FD) 590 (M+); Anal. Calc'd for: C, 56.11; H, 6.08; N, 12.66. Found: C, 56.16; H, 5.92; N, 12.56. IR (KBr) 3163.75, 3031.15, 2952.46, 2876.38, 1745.07, 1664.94, 1530.69, 1497.79, 1453.37, 1435.81, 1197.21, 1177.62, 1094.93, 747.95, 701.04 cm⁻¹.

96

Preparation 40

5 To a solution of a compound of Preparation 28 (1.0 g, 4.0 mmol), morpholine (0.35 mL, 4.0 mmol), 1hydroxybenzotriazole hydrate (0.6 g, 4.4 mmol) stirring in N,N-dimethylformamide (50 mL) at room temperature was added 1,3-dicyclohexylcarbodiimide (0.9 g, 4.4 mmol). After 16 h, 10 the mixture was concentrated, and the residue extracted with ethyl acetate. The combined organic extracts were filtered, washed with saturated aqueous sodium bicarbonate, water, brine, dried over sodium sulfate, filtered, and concentrated. The resulting residue was purified by silica gel chromatography (methanol/chloroform) to give 0.75 g 15 (60%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (FD) 316 (M+).

97

Preparation 41

5 To a slurry of 5% Pd/C (0.18 g) in ethyl acetate (5 mL) was carefully added a solution of a compound of Preparation 40 (0.67 g, 2.0 mmol) in ethyl acetate (25 mL)/ethanol (25 mL). The resulting slurry was treated with hydrogen gas at 40 psi on a Parr apparatus. After 1 h, a slurry of 5% Pd/C 10 (0.18 g) in ethyl acetate (10 mL) was added to this mixture, followed by hydrogenation at 40 psi. After 1 h, the mixture was filtered through celite and concentrated. To the residue stirring in N,N-dimethylformamide (100 mL) was added a compound of Preparation 4 (0.53 g, 1.4) and 1-15 hydroxybenzotriazole hydrate (0.21 g, 1.54 mmol) followed by 1,3-dicyclohexylcarbodiimide (0.32 g, 1.54 mmol). After 16 h at room temperature, the solution was concentrated and extracted with ethyl acetate. The combined organic extracts were washed with water, brine, dried over sodium sulfate, 20 filtered, and concentrated. The resulting residue was purified by silica gel chromatography (methanol/chloroform) to yield 0.27 g (30%) of the desired product as a tan foam: H-NMR is consistent with structure; MS (FD) 448 (M+).

98

Example 9

To a solution of a compound of Preparation 41 (0.27 g, 0.42 mmol) stirring in dichloromethane (12 mL) at room temperature was added trifluoroacetic acid (4 mL, 51 mmol).

After 1.5 h, water (40 mL) was added and the reaction mixture quenched carefully with solid sodium bicarbonate.

- The resulting mixture was extracted with ethyl acetate and the combined organic extracts washed with brine, dried over sodium sulfate, filtered and concentrated. The concentrate was dissolved in ethyl acetate (40 mL) and subsequently treated with a saturated solution of HCl in ethyl acetate
- 15 (40 mL). After 15 min, the mixture was concentrated to give 0.14 g (54%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 548 (M+); Anal.

Calc'd. for: C, 56.04; H, 6.16; N, 13.52. Found: C, 55.78; H, 6.11; N, 13.27; IR (KBr) 2927, 2858.9, 1659.3, 1542.2,

20 1114.4 cm⁻¹.

99

Preparation 42

Reaction of a compound of Preparation 28 (1.0 g, 4.0 mmol), piperidine (0.4 mL, 4.0 mmol), 1-hydroxybenzotriazole hydrate (0.6 g, 4.4 mmol) and 1,3-dicyclohexylcarbodiimide (0.9 g, 4.4 mmol) in N,N-dimethylformamide (50 mL) gave 0.95 g (75%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 314 (M+).

Preparation 43

15

Hydrogenation of a compound of Preparation 42 (0.91 g, 2.9 mmol) in ethyl acetate (50 mL)/ethanol (50 mL), 5% Pd/C (0.36 g) in ethyl acetate (5 mL) followed by reaction with a compound of Preparation 4 (0.95 g, 2.5 mmol), 1-

20 hydroxybenzotriazole hydrate (0.37 g, 2.75 mmol), and 1,3-

100

dicyclohexylcarbodiimide (0.57 g, 2.75 mmol) gave 0.43 g (25%) of the desired product as a tan foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 646 (M+).

5 Example 10

2HC1

Reaction of a compound of Preparation 43 (0.38 g, 0.59 mmol) and trifluoroacetic acid (4 mL, 51 mmol) in dichloromethane (12 mL) followed by acidification with HCl gave 0.03 g (8.3%) of the desired product as a tan solid:

1H-NMR is consistent with structure; MS (FD) 546 (M+); IR (KBr) 3141, 2937, 2859, 1642, 1534, 1453, 1444 cm⁻¹.

15

101

Pr paration 44

5 To a slurry of 5% Pd/C (1.0 g) in ethyl acetate (25 mL) was added a solution of a compound of Preparation 6 (8.25 g, 30 mmol) in ethyl acetate (25 mL)/absolute ethanol (25 mL). The slurry was hydrogenated at 40 psi on a Parr apparatus. After 75 min, a slurry of 5% Pd/C (0.7 g) in ethyl acetate 10 (25 mL) was added to the reaction mixture. After hydrogenation at 40 psi for 1.5 h, the mixture was filtered through celite and concentrated. The concentrate was dissolved in N, N-dimethylformamide (500 mL) and boc-dbenzyloxyserine (9.0 g, 30.8 mmol), 1-hydroxybenzotriazole 15 hydrate (4.5 g, 33 mmol) and 1,3-dicyclohexylcarbodiimide (6.8 g, 33 mmol) added. After 16 h at ambient temperature, the mixture was concentrated and the residue extracted with ethyl acetate. The combined organic extracts were washed with water, brine, dried over sodium sulfate, filtered, and 20 concentrated. Purification by silica gel chromatography (methanol /chloroform) gave 8.33 g (53%) of the desired product as a tan solid: 1H-NMR is consistent with structure; MS (FD) 522 (M+); Anal. Calc'd. for: C, 64.35; H, 6.56; N, 10.72. Found: C, 64.59; H, 6.83; N, 10.77.

102

Preparation 45

To a solution of a compound of Preparation 44 (8.1 g, 15.5 mmol) stirring at room temperature in dichloromethane (75 mL) was added trifluoroacetic acid (25 mL, 320 mmol). After 50 min, the mixture was carefully poured into a saturated aqueous solution of sodium bicarbonate and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated to give 6.5 g (99%) of the desired product as a tan solid. ¹H-NMR is consistent with structure; MS (FD) 422 (M+).

15

Preparation 46

To a solution of a compound of Preparation 45 (6.5 g, 15.0 mmol), boc- α -aminoisobutyric acid (3.05 g, 15.0 mmol),

103

1-hydroxybenzotriazole hydrate (2.23 g, 16.5 mmol) stirring in N,N-dimethylformamide (400 mL) at room temperature was added 1,3-dicyclohexylcarbodiimide (3.4 g, 16.5 mmol).

After 16 h, the mixture was concentrated and the resulting residue extracted with ethyl acetate. The combined organic extracts were washed with water, brine, dried over sodium sulfate, filtered, and concentrated. Purification by silica gel chromatography (methanol/chloroform) gave 6.39 g (70%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 607 (M+). Anal. Calc'd. for: C, 63.25; H, 6.80; N, 11.52. Found: C, 63.36; H, 6.92; N, 11.59.

Preparation 47

15

To a solution of a compound of Preparation 46 (6.04 g, 9.9 mmol) stirring in absolute ethanol

20 (50 mL)/tetrahydrofuran (50 mL) at room temperature was added 1N NaOH (50 mL, 49.5 mmol). After 30 min, the mixture was acidified with 1N HCl and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated to give 5.4

g (94%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 580 (M+); Anal. Calc'd.

104

for: C, 62.16; H, 6.43; N, 12.08. Found: C, 61.86; H, 6.29; N, 12.06.

Preparation 48

5

10

15

20

To a solution of a compound of Preparation 47 (0.7 g, 1.2 mmol), N-methylamine hydrochloride (0.08 g, 1.2 mmol), triethylamine (0.5 mL, 3.6 mmol), and 1-hydroxybenzotriazole hydrate (0.18 g, 1.32 mmol) stirring in N,N-dimethylformamide (50 mL) at room temperature was added 1,3-dicyclohexylcarbodiimide (0.27 g, 1.32 mmol). After 16 h, the mixture was concentrated and the resulting residue extracted with ethyl acetate. The combined organic extracts were washed with water, brine, dried over sodium sulfate, filtered, and concentrated. Purification by silica gel chromatography (methanol /chloroform) gave 0.25 g (35%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 592.4 (M+); Anal. Calc'd for 0.32 mol hydrate: C, 62.21; H, 6.76; N, 14.04. Found: C, 62.17; H, 6.74; N, 14.19.

105

Example 11

2HCl

5 To a slurry of a compound of Preparation 48 (0.2 g, 0.34 mmol) stirring in dichloromethane (12 mL) at room temperature was added trifluoroacetic acid (4 mL, 52 mmol). After 2 h, additional trifluoroacetic acid (4 mL, 52 mmol) was added and the reaction was heated to reflux. After 7 h, 10 the mixture was cooled to room temperature, water (40 mL) added, followed excess solid sodium bicarbonate. mixture was extracted with ethyl acetate and the combined organic extracts washed with brine, dried over sodium sulfate, filtered and concentrate. The resulting crude 15 product was dissolved in ethyl acetate (40 mL) and a saturated solution of HCl in diethyl ether was added (40 mL). After 15 min, this slurry was concentrated to give 0.13 g (68%) of the desired product as a white solid: 1H-NMR is consistent with structure; MS (FD) 492 (M+); Anal. 20 Calc'd for: C, 55.22; H, 6.06; N, 14.86. Found: 55.33; H, 6.28; N, 13.24; IR (KBr) 3224, 3061, 3032, 2962, 2936, 2873, 1678, 1636, 1538, 1498, 1454, 1101 cm⁻¹.

106

Preparation 49

Reaction of a compound of Preparation 47 (1.00 g, 580 mmol), hexamethyleneimine (0.2 mL, 1.7 mmol), 1-hydroxybenzotriazole hydrate (0.25 g, 1.9 mmol) and 1,3-dicyclohexylcarbodiimide (0.4 g, 1.9 mmol) in N,N-dimethylformamide (50 mL) as described in Preparation 4 gave 0.76 g (68%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 660.2 (M+); Anal. Calc'd for: C, 65.43; H, 7.32; N, 12.02. Found: C, 65.92; H, 7.86; N, 11.71.

15

20

107

Example 12

2.67 HC1

Reaction of a compound of Preparation 49 (0.67 g, 1.0 mmol) and trifluoroacetic acid (4 mL, 52 mmol) in dichloromethane (12 mL) for 1 h at ambient temperature, followed by acidification with HCl in ethyl acetate, according to Preparation 4 gave 0.3 g (48%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 560.4 (M); Anal. Calc'd for: C, 58.77; H, 6.56; N, 12.01. Found: C, 56.48; H, 6.41; N, 12.06.

Preparation 50

15

20

To a solution of m-nitroaniline (1.0 g, 7.24 mmol) stirred in anhydrous N,N-dimethylformamide (40 mL) at room temperature was added a solution of a compound of

108

Preparation 5 (2.11 g, 8.69 mmol) in anhydrous N, Ndimethylformamide (10 mL). After 2.5 h, the reaction mixture was diluted with H2O (70 mL) and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried (Na₂SO₄), and concentrated to give a yellow oil. Purification by radial chromatography (silica gel, 10%-75% ethyl acetate/hexanes) provided 1.65 g (76%) of the product (1:1 mixture of diastereomers) as an orange solid. ¹H NMR (300 MHz, CDCl₃) d 7.47-7.53 (m, 3H), 7.33-7.41 (m, 4H), 7.20-7.25 (app. t, 1H, J = 8.1Hz), 6.81-6.8510 (dd, 1H, J = 8.0Hz; 2.1Hz), 5.10 (s, 1H), 4.12-4.26 (m, 2H),1.20-1.25 (t, 3H, J = 7.1Hz); ^{13}C NMR (75.5 MHz, CDCl₃) d 171.0, 149.2, 146.5, 136.4, 129.6, 128.9, 128.5, 127.0, 119.1, 112.5, 107.2, 62.1, 60.3, 13.9; FD+ MS for C16H16N2O4 = 300; Anal. calcd. for C16H16N2O4: C, 63.99; H, 5.37; N, 9.33; Found: C, 64.77; H, 5.26; N, 9.17.

Preparation 51

20

25

To a slurry of sodium hydride (0.15 g of a 60% dispersion in mineral oil, 3.86 mmol) stirring in N,N-dimethylformamide (30 mL) at room temperature, was added a solution of 6-nitrobenzimidazole (0.60 g, 3.68 mmol) in N,N-dimethylformamide (10 mL). After 10 min, a solution of α -bromophenylacetic acid ethylester in N,N-dimethylformamide (10 mL) was added and the solution stirred for 4 h at room temperature, quenched with water, and extracted with ethyl

109

acetate. The combined organic extracts were washed with water, brine, dried (Na₂SO₄) and concentrated. Purification by silica gel chromatography (25%-75% ethyl acetate/hexanes) gave 0.580 g (50%) of the product (mixture of diastereomers)

5 as a yellow oil: ¹H NMR (300 MHz, CDCl₃) d 8.72-8.73 (d, 1H, J = 2.1Hz), 8.22-8.27 (dd, 1H, J = 9.2Hz; 2.1Hz), 8.15 (s, 1H), 7.44-7.50 (app. t, 3H, J = 6.9Hz), 7.34-7.41 (m, 3H), 6.19 (s, 1H), 4.26-4.39 (m, 2H), 1.27-1.33 (t, 3H, J = 6.9Hz); FD+ MS for C₁7H₁5N₃O₄ = 325; Anal. calcd. for

10 C₁7H₁5N₃O₄: C, 62.76; H, 4.65; N, 12,92; Found: C, 62.89; H, 4.92; N, 12.92.

Preparation 52

15

20

25

To a solution of a compound of Preparation 50 (0.81 g, 2.73 mmol) stirring in dioxane (30 mL) at room temperature was added LiOH·H₂O (0.57 g, 13.6 mmol) and H₂O (15 mL). After 45 min, the mixture was concentrated to a volume of approximately 20 mL. The resulting aqueous solution was diluted with H₂O (75 mL) and extracted with diethyl ether. The aqueous layer was acidified with 1N HCl and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried (Na₂SO₄) and concentrated to give 0.71 g (95%) of the product (1:1 mixture of diastereomers) as a yellow solid: 1 H NMR (300 MHz, CDCl₃) δ 7.48-7.55 (m,

110

3H), 7.35-7.43 (m, 4H), 7.21-7.27 (app. t, 1H, J = 8.1Hz), 6.81-6.85 (dd, 1H, J = 8.2Hz; 2.0Hz), 5.16 (s, 1H); FD+ MS for $C_{14}H_{12}N_{2}O_{4} = 272$; Anal. calcd. for $C_{14}H_{12}N_{2}O_{4}$: C, 61.76; H, 4.44; N, 10.29; Found: C, 62.15; H, 4.52; N, 9.63.

Preparation 53

10

5

To a solution of a compound of Preparation 51 (0.48 g, 1.48 mmol) stirring in dioxane (20 mL) at room temperature was added $\text{LiOH} \cdot \text{H}_2\text{O}$ (0.31 g, 7.38 mmol) and H_2O (10 mL). After 45 min, the reaction mixture was concentrated to a 15 volume of approximately 15 mL. The resulting aqueous solution was diluted with H2O (75 mL) and extracted with diethyl ether. The aqueous layer was acidified with 1N HCl and extracted with ethyl acetate. The combined organic 20 extracts were washed with brine, dried (Na2SO4) and concentrated to give 0.450 g (>95%) of the product (1:1 mixture of diastereomers) as a light yellow solid: 1H NMR (300 MHz, DMSO) δ 8.63 (s, 1H), 8.56-8.57 (d, 1H, J = 2.1Hz), 8.14-8.20 (dd, 1H, J = 9.2Hz; 2.1Hz), 7.82-7.86 (d, 1H, J = 9.2Hz), 7.52-7.58 (m, 2H), 7.38-7.49 (m, 3H), 6.88(s, 1H); FD+ MS for C15H11N3O4 = 297; Anal. calcd. for C₁5H₁1N₃O₄: C, 60.61; H, 3.73; N, 14.14; Found: C, 59.59; H, 4.16; N, 12.78.

111

Preparation 54

5

To a solution of a compound of Preparation 52 (0.75 g, 2.78 mmol), L-proline methyl ester hydrochloride (0.46 g, 2.78 mmol), 1-hydroxybenzotriazole hydrate (0.38 g, 2.78 mmol) and N,N-diisopropylethylamine (1.26 g, 9.72 mmol) in 10 anhydrous 1,2-dichlormethane (30 mL) stirring at room temperature, was added 1-(3-dimethylaminopropyl)-3ethylcarbodiimide (0.585 g, 3.05 mmol). After 18 h, the reaction mixture was diluted with H2O (50 mL) extracted with ethyl acetate. The combined organic extracts were washed 15 with 10% citric acid, sat'd aqueous sodium bicarbonate, water, brine, dried (Na₂SO₄) and concentrated. Purification by radial chromatography (silica gel, 40%-75% ethyl acetate/hexanes) gave 0.56 g (53%) of the product (1:1 mixture of diastereomers) as a yellow solid: ¹H NMR (300 20 MHz, CDCl₃) δ 7.43-7.50 (m, 3H), 7.27-7.43 (m, 4H), 7.13-7.20 (app. t, 1H, J = 7.5Hz), 6.83-6.91 (t, 1H, J = 5.8Hz), 5.14 (s, 1H), 4.52-4.58 (m, 0.5H), 4.41-4.47 (m, 0.5H), 3.89-3.97 (m, 1H), 3.71 (s, 1.5H), 3.62 (s, 1.5H), 3.23-3.36(m, 1H), 1.82-2,24 (m, 5H); ¹³C NMR (75.5 MHz, CDCl₃) d172.2, 171.7, 168.7, 168.5, 149.0, 146.9, 146.5, 136.4, 25 135.9, 129.5, 129.4, 129.0, 128.8, 128.5, 128.2, 128.0,

112

127.8, 119.9, 119.6, 112.2, 112.0, 106.5, 106.5, 59.5, 59.4, 59.3, 59.3, 52.2, 52.0, 46.7, 46.7, 28.7, 28.6, 24.9, 24.5; FD+ MS for C₂₀H₂₁N₃O₅ = 383; Anal. calcd. for C₂₀H₂₁N₃O₅: C, 62.65; H, 5.52; N, 10.96; Found: C, 61.93; H, 5.62; N, 10.46.

Preparation 55

10

15

20

5

To a solution of a compound of Preparation 53 (0.43 g, 1.46 mmol), L-proline methyl ester hydrochloride (0.24 g, 1.46 mmol), 1-hydroxybenzotriazole hydrate (0.20 g, 1.46 mmol) and N,N-diisopropylethylamine (0.66 g, 5.10 mmol) stirring in anhydrous 1,2-dichlormethane (30 mL) at room temperature was added 1-(3-dimethylaminopropyl)-3ethylcarbodiimide (0.31 g, 1.60 mmol). After 18 h, the reaction mixture was quenched with H2O (50 mL) and extracted with ethyl acetate. The combined organic extracts were washed with 10% citric acid, saturated aqueous sodium bicarbonate, H2O, brine, dried (Na2SO4) and concentrated. Purification by radial chromatography (silica gel, 50% ethyl acetate/hexanes to 100% ethyl acetate gradient) gave 0.25 g (42%) of the a single diastereomer as a white foam solid: ¹H NMR (300 MHz, CDCl₃) d 8.75-8.76 (δ , 1H, J = 2.1Hz), 8.28-8.32 (dd, 1H, J = 8.9Hz; 2.1Hz), 7.91 (s, 1H), 7.45-7.58 (m, 6H), 6.26 (s, 1H), 4.65-4.70 (m, 1H), 3.83-3.92 (m,1H), 3.78 (s, 3H), 3.30-3.39 (m, 1H), 1.95-2.30 (m, 5H); FD+

113

MS for $C_{21}H_{20}N_{4}O_{5} = 408$; Anal. calcd. for $C_{21}H_{20}N_{4}O_{5}$: C, 61.76; H, 4.94; N, 13.72; Found: C, 61.24; H, 5.16; N, 13.10.

5

Preparation 56

To a slurry of 5% Pd/C (0.07 g) in ethanol (30 mL) was added a solution of a compound of Preparation 54 (0.15 g, 10 0.39 mmol) in ethyl acetate (30 mL). The mixture was treated with hydrogen gas (32 psi) at room temperature for 4 h on a Parr apparatus then carefully filtered through celite. The resulting filtrate was evaporated to provide an off-white solid foam which was dissolved in N,N-15 dimethylformamide (30 mL). To this solution was added a compound of Preparation 4 (0.16 g, 0.41 mmol), 1hydroxybenzotriazole hydrate (0.06 g, 0.41 mmol) and 1,3dicyclohexylcarbodiimide (0.09 g, 0.45 mmol). This solution 20 was stirred overnight at room temperature and subsequently diluted with water (50 mL) then extracted with ethyl acetate. The combined organic extracts were washed with water, brine, dried (Na2SO4) and evaporated to provide a tan foam. Purification by radial chromatography (silica gel, 25 50% ethyl acetate/hexanes to 100% ethyl acetate gradient) yielded 0.23 g (82%) of the product (mixture of diastereomers) as an off-white solid foam. ¹H NMR (300 MHz, $CDC1_3$) d 8.83-8.91 (m, 1H), 7.44-7.51 (m, 2H), 7.20-7.36 (m,

114

8H), 6.88-7.06 (m, 3H), 6.32-6.38 (app. t, 1H, J = 6.9Hz), 5.28 (s, 1H), 5.12-5.19 (m, 1H), 4.88-4.91 (br. s, 1H), 4.48-4.60 (m, 3H), 4.17-4.24 (m, 1H), 3.64-3.72 (app. q, 2H, J = 8.0Hz), 3.62 (s, 3H), 3.39-3.52 (m, 1H), 3.28-3.39 (m, 1H), 1.81-2.15 (m, 5H), 1.53-1.57 (app. d, 3H, J = 7.9Hz), 1.38 (s, 3H), 1.39 (s, 9H); FD+ MS for C39H49N5O8 = 716; Anal. calcd. for C39H49N5O8: C, 65.44; H, 6.90; N, 9.78; Found: C, 65.23; H, 7.43; N, 10.34.

10

Preparation 57

To a slurry of 5% Pd/C (0.042 g) in ethanol (30 mL) was 15 added a solution of a compound of Preparation 55 (0.08 g, 0.20 mmol) in ethyl acetate (30 mL). The mixture was treated with hydrogen gas (32 psi) at room temperature for 4 h (Parr apparatus) then carefully filtered through celite. The resulting filtrate was evaporated to provide a white 20 solid foam which was dissolved in N,N-dimethylformamide (20 mL). To this solution was added a compound of Preparation 4 (0.08 g, 0.20 mmol), 1-hydroxybenzotriazole hydrate (0.03 g, 0.22 mmol) and 1,3-dicyclohexylcarbodiimide (0.05 g, 0.22 mmol). This solution was stirred overnight at room 25 temperature and subsequently diluted with water (50 mL) then extracted with ethyl acetate. The combined organic extracts were washed with water and brine, dried (Na2SO4) and

115

concentrated. Purification by radial chromatography (silica gel, 75% ethyl acetate/hexanes to 100% ethyl acetate gradient) yielded 0.10 g (66%) of the product (one diastereomer) as an off-white solid foam: ^{1}H NMR was consistent with structure; FD+ MS for C40H48N6O8 = 740.

Example 13

10

15

20

To a solution of a compound of Preparation 56 (0.17 g, 0.24 mmol) and anisole (0.03 g, 0.26 mmol) stirring in anhydrous dichloromethane (5 mL) at 0 °C was added trifluoroacetic acid (1 mL). After 4 h, the reaction mixture was quenched carefully with saturated aqueous sodium bicarbonate extracted with ethyl acetate. The combined organic extracts were washed with sat'd aqueous sodium bicarbonate, water, brine, dried (NaS2O4) and evaporated to yield the desired product (1:1 mixture of diastereomers) as an off-white foam: 0.100 g (67%). ¹H NMR was consistent with structure; FD+ MS for C34H41N5O6 = 615; Anal. calcd for C34H41N5O6: C, 66.32; H, 6.71; N, 11.37; Found: C, 65.83; H, 6.50; N, 6.50.

116

Example 14

5 To a solution of a compound of Preparation 57 (0.080 g, 0.11 mmol) and anisole (0.0123 g, 0.114 mmol) stirring in anhydrous dichloromethane (5 mL) at 0 °C was added trifluoroacetic acid (1 mL). After 4 h, the mixture was quenched carefully with saturated sodium bicarbonate and extracted with ethyl acetate. The combined organic extracts 10 were washed with saturated aqueous sodium bicarbonate, water, brine, dried (NaS2O4) and concentrated to yield foam 0.09 g (95%) of the desired product (one diastereomer) as an off-white solid: ¹H NMR was consistent with structure; FD+ 15 MS for $C35H40N6O6 \cdot 2CF3COOH = 640 (M-2CF3COOH)$; Anal. calcd. for C39H42N6O10F6: C, 53.92; H, 4.87; N, 9.67; Found: C, 51.86; H, 4.74; N, 9.54.

20

117

EXAMPLES PART 2A

Preparation 1a

tert-Butyloxycarbonyl-O-benzyl-D-serine methyl ester.

To a solution of t-butyloxycarbonyl-O-benzyl-D-serine (25.0 g, 84.7 mmol) stirring in dimethylformamide (500 mL) at room temperature was added sodium bicarbonate (14.2 g, 169 mmol) followed by methyl iodide (26.4 mL, After 18 h, the reaction mixture was 424 mmol). concentrated to approximately 100 mL. Ethyl acetate was added and the mixture washed with aqueous sodium bicarbonate and brine. The organic extract was dried and concentrated to give the desired compound (25 g, 96%) as a light yellow oil: ¹H NMR (300 MHz, CDCl₃) d 1.45 (s, 9H), 3.70 (m, 1H), 3.75 (s, 3H), 3.85 (m, 1H), 4.50 (m, 3H), 7.30 (m, 5H); MS (FD) m/e 310; Anal. calc'd for C16H23NO5: C, 62.12; H, 7.49; N, 4.53. Found: C, 62.31; H, 7.49; N, 4.43.

Preparation 1b

O-benzyl-D-serine methyl ester.

To a solution of tert-butyloxycarbonyl-O-benzyl-D-serine methyl ester (BF8-EZ0-275) (5.0 g, 16 mmol) stirring in dichloromethane (40 mL) and anisole (1 mL) at 0 °C was added trifluoroacetic acid (10 mL). After 4 h at room temperature, a saturated aqueous solution of sodium bicarbonate was added and the resulting mixture extracted with ethyl acetate. The combined organic extracts were

118

washed with brine, dried over sodium sufate, and concentrated. The crude product was used in the next step without further purification.

Preparation 1c

To a solution of O-benzyl-D-serine methyl ester (the product of Preparation 1b) (65.4 mmol), boc-ag, aminoisobutyric acid (13.2 65.4 mmol), 1hydroxybenzotriazole (8.8 65.4 mmol), g, and diisopropylethylamine (22.8 mL, 130.7 mmol) stirring in dichloromethane (500 mL) at 0 °C was added 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (12.3 71.9 mmol). After 18 h, ethyl acetate and ammonium chloride (saturated aqueous solution) were added and the resulting mixture extracted with aqueous ammonium chloride, aqueous sodium bicarbonate, and brine. The organic extracts were dried over sodium sulfate and concentrated. Purification by flash chromatography (25% ethyl acetate/hexanes) yielded the desired compound (21.6 g, 83%) as a white solid: 1 H NMR (300 MHz, CDCl₃) d 1.39 (s, 9H), 1.48 (s, 6H), 3.62 (dd, J = 3.4, 9.1 Hz, 1H), 3.70 (s, 3H), 3.85(dd, J = 3.4, 9.1 Hz, 1H), 4.48 (dd, J = 12.5, 22.7 Hz,2H), 4.75 (m, 1H), 4.92 (s, 1H), 7.11 (d, J = 8.6 Hz, 1H), 7.35 (m, 5H); MS (FD) m/e 395; Anal. calc'd for C20H30N2O6: C, 60.90; H, 7.67; N, 7.10. Found: C, 61.02; H, 7.78; N, 7.10.

119

Preparation 1d

To a solution of the product of Preparation 1c (5.30 g, 13.4) stirring in dioxane (100 mL)/water (50 mL) at room temperature was added lithium hydroxide (2.80 g, 67.3 After 18 h, water was added and the solution concentrated. The resulting mixture was extracted with diethyl ether. Brine was added to the aqueous layer and the pH adjusted to 3.5 with 1 N HCl. The resulting mixture was extracted with ethyl acetate and the combined organic extracts dried over sodium sulfate concentrated to yield the title compound (4.40 g, 86%) as a white foam: ^{1}H NMR (300 MHz, CDCl₃) d 1.39 (s, 9H), 1.45 (s, 3H), 1.47 (s, 3H), 3.68 (m, 1H), 3.95 (m, 1H), 4.54 (s, 2H), 4.70 (m, 1H), 5.51 (bs, 1H), 7.18 (d, J =9.1 Hz, 1H), 7.25 (m, 5H), 9.90 (bs, 1H); MS (FD) m/e 381; Anal. calc'd for C19H28N2O6: C, 59.99; H, 7.42; N, 7.36. Found: C, 59.74; H, 7.26; N, 7.30.

Preparation 1e

A solution of sodium ethoxide was generated by the addition of sodium metal (52.89 grams, 2.3007 mol) over 3 hours to ethanol (1500 mL). To the sodium ethoxide solution at ambient temperature was added a solution of diethylacetamidomalonate (499.75 grams, 2.3007 mol) dissolved in ethanol (225 mL). The reaction mixture was stirred for 1.5 hours at ambient temperature. 1-bromo-3-phenylpropane (458.07 grams, 2.3007 mol) was added over 15 minutes and the reaction mixture was refluxed until

complete as determined by hplc (16 hours). The reaction mixture was concentrated to dryness and the residue partitioned between ethyl acetate (1 x 1500 mL and 2 x 500 mL) and water (1500 mL). The ethyl acetate layers were combined, washed with saturated sodium chloride solution (4 \times 500 mL), dried using sodium sulfate, and concentrated to give 752.1 grams (98%) of the desired compound as a light yellow solid. A 1.0 gram sample was recrystallized from hexane:ethyl acetate (19:1, v:v) to give a mp 84-86°C. ¹H nmr (CDCl₃): δ 1.18-1.23 (t, 6H), 1.37-1.50 (m, 2H), 2.02 (s, 3H), 2.34-2.41 (m, 2H), 2.58-2.62 (t, 2H), 4.16-4.24 (q, 4H), 6.76 (s, broad, 1H), 7.11-7.28 (m, 5H). ¹³C nmr (CDCl₃): δ 23.03, 25.67, 31.85, 35.45, 62.46, 66.49, 125.40, 125.90, 128.27, 128.35, 141.77, 168.11, 168.94. MS (FIA) m/z 336.3 ($[M+H]^+$). IR (KBr, cm^{-1}) 1645.98 (amide), 1744.76 (C=O). Anal. Calcd. for $C_{18}H_{25}NO_5$: C, 64.46; H, 7.51; N, 4.17. Found: C, 64.60; H, 7.37; N, 4.39.

Preparation 1f

(DL)-N-Acetyl-2-amino-5-phenylpentanoic Acid. A slurry consisting of the product of Preparation 1e (249.15 grams, 0.7428 mol) and 2.5 N sodium hydroxide solution was heated at 100°C for three hours. The reaction mixture was cooled to 30°C and the pH adjusted to 5.0 using concentrated hydrochloric acid. The solution was heated to 100 °C and the pH was held at 5.0 using concentrated hydrochloric acid as needed until the reaction was complete as determined by hplc. The solution was filtered while hot through diatomaceous earth. The

filtrate was cooled to 5-10 °C and the pH adjusted to 1.0 using concentrated hydrochloric acid. The resulting slurry was stirred for 1 hour at 5 °C, filtered, and dried in vacuum at 50 °C to give 160.34 grams (92%) of (DL)-Nacety1-2-amino-5-phenylpentanoic acid as a white powder, mp 145-148 °C. ¹H nmr (DMSO-d₆): δ 1.60-1.71 (m, 4H), 1.86 (s, 3H), 2.56-2.59 (m, 2H), 4.19-4.23 (m, 1H), 7.16-7.30 (m, 5H), 8.14 (d, 1H). 13 C nmr (DMSO-d₆): δ 23.17, 28.25, 31.55, 35.51, 52.55, 126.60, 129.14, 142.64, 170.25, 174.65. MS (FIA) m/z 236.2 (M⁺). IR (KBr, cm⁻ 1609.17 (amide), 1741.12 (C=O). Anal. Calcd. for $C_{13}H_{17}NO_3$: C, 66.36; H, 7.28; N, 5.95. Found: C, 66.41; H, 7.15; N, 5.96.

Preparation 1g

(D) - N-Acetyl-2-amino-5-phenylpentanoic Acid. A solution consisting of (DL)-N-acety1-2-amino-5-phenylpentanoic (438.0 grams, 1.862 mol), cobalt chloride (1.10 acid grams), 2N potassium hydroxide solution (931 mL, 1.862 mol), and water (8000 mL) was adjusted to a pH of 8.0 by the addition of 2N potassium hydroxide solution. reaction mixture was added Acylase I (aspergillus melleus, 39.42 grams) and vigorously stirred for 24 hours at 40 °C while maintaining a pH of 8.0 by addition of 2N potassium hydroxide. The resulting slurry was filtered. The filtrate was adjusted to a pH of 2.0 giving a thick slurry. The product was isolated by filtration, washed with hexane (2000 mL) and dried in vacuum at 50 °C to give 188.52 grams (43%) of (D)-N-acety1-2-amino-5phenylpentanoic acid. ¹H nmr (DMSO-d₆): δ 1.59-1.74

122

(m, 4H), 1.86 (s, 3H), 2.57-2.60 (m, 2H), 4.22-4.26 (m, 1H), 7.16-7.30 (m, 5H), 8.02 (d, 1H), 12.39 (s, broad, 1H). 13 C nmr (DMSO-d₆): δ 23.18, 28.13, 31.66, 35.54, 52.58, 126.56, 129.10, 142.67, 170.12, 174.48. MS (FIA) m/z 236.1 (M⁺). IR (KBr, cm⁻¹) 1625.08 (amide), 1700.24 (C=O). Anal. Calcd. for $C_{13}H_{17}NO_3$: C, 66.36; H, 7.28; N, 5.95. Found: C, 66.49; H, 7.00; N, 6.03.

Preparation 1h

(D)-2-Amino-5-phenylpentanoic Acid, Ethyl Ester Hydrochloride. A solution consisting of (D)-N-acety1-2amino-5-phenylpentanoic acid (188.8 grams, 0.8024 mol), ethanol (535 mL), and concentrated hydrochloric acid (268 mL, 3.21 mol) was warmed to 85 °C and monitored by hplc. The reaction was determined to be incomplete by hplc at 14.5 hours and additional concentrated hydrochloric acid (50 mL) was added. The reaction was determined to be complete by hplc after 22.5 hours. Water azeotropically distilled from the reaction by continuous addition and distillation of 8000 mL of ethanol. ethanol was azeotropically distilled from the reaction by the continuous addition and distillation of ethyl acetate (2000 mL). Upon cooling the solution to 0 °C the product crystallized. The solution containing the product was stirred for 1 hour at 0 °C, filtered, and the cake dried in vacuum at 40 °C to give 199.0 grams (96%) of 2-amino-5phenylpentanoic acid, ethyl ester hydrochloride, mp 117-¹H nmr (DMSO-d₆): δ 1.15-1.21 (t, 3H), 1.50-1.89 (m, 4H), 2.48-2.67 (m, 2H), 3.92-3.98 (t, 1H), 4.08-4.25 (m, 2H), 7.12-7.29 (m, 5H), 8.76 (s, broad, 3H). ¹³C nmr $(DMSO-d_6): \delta 13.90, 25.97, 29.52, 34.41, 51.71,$

123

61.56, 124.91, 125.81, 128.24, 141.27, 169.35. MS (FIA) m/z 222.3 (M $^{+}$). IR (KBr, cm $^{-1}$) 1741.14 (C=O). [α] $^{20}_{D}$ = -11.17(c = 30.62 mg / 3mL, MeOH). Anal. Calcd. for C₁₃H₂₀NO₂Cl: C, 60.58; H, 7.82; N, 5.43. Found: C, 60.45; H, 7.67; N, 5.55.

Preparation 1i

A slurry consisting of N-t-BOC- α -aminoisobutyric acid (90.64 grams, 0.446 mol), 2-chloro-4,6-dimethoxy-1,3,5triazine (75.90 grams, 0.425 mol), N-methyl morpholine (88.13 grams, 0.871 mol), and diethyl ether (1000 mL) was stirred at ambient temperature until complete determined by hplc (3 hours). The D-2-amino-5phenylpentanoic acid, ethyl ester hydrochloride (109.55 grams, 0.425 mol) was added and the reaction mixture stirred for 16 hours at ambient temperature. reaction mixture was partitioned between 10% citric acid solution (1000 mL) and ethyl acetate (3 \times 500 mL). organic phase was washed with 10% citric acid solution (3 \times 500 mL), saturated sodium bicarbonate solution (3 \times 500 mL), water (1 x 500 mL), dried using sodium sulfate, and concentrated to dryness. The residue was recrystallized from hexane (3000 mL) to give 155.11 grams of the desired compound: mp 97-99 °C. 1 H nmr (CDCl₃): δ 1.25-1.28 (t, 3H), 1.43 (s, 9H), 1.48 (s, 3H), 1.50 (s, 3H), 1.70-1.73 (m, 3H), 1.87-1.93 (m, 1H), 2.62-2.67 (m, 2H), 4.16-4.21(m, 2H), 4.57-4.62 (m, 1H), 4.95 (s, 1H), 6.96 (s, broad, 1H), 7.16-7.19 (m, 3H), 7.26-7.33 (m, 2H). ¹³C nmr ($CDCl_3$): δ 14.53, 26.32, 27.17, 28.67, 32.47, 35.73,

124

52.54, 57.17, 61.62, 126.21, 128.69, 128.79, 142.12, 154.99, 172.81, 174.69. MS (FIA) m/z 407.5 ([M+H] †). IR (KBr, cm $^{-1}$) 1652.75, 1685.52 (amides), 1741.73 (C=O). [α] $^{20}_{D}$ = 7.83 (c = 10.22 mg / 1mL, MeOH). UV (0.1% trifluoroacetic acid in water : acetonitrile) λ_{max} 215.6 nm. Anal. Calcd. for $C_{22}H_{34}N_2O_5$: C, 65.00; H, 8.43; N, 6.89. Found: C, 65.23; H, 8.34; N, 6.94.

Preparation 1j


A solution consisting of the product of Preparation 1i (152.53 grams, 0.3752 mol) and tetrahydrofuran (884 mL) was cooled to 5 °C. A solution consisting of lithium hydroxide (26.96 grams, 1.126 mol) and water (1419 mL) was added to the reaction dropwise over 10 minutes maintaining a temperature of 5-10 °C. Ethanol (183 mL) was added and the reaction stirred at 5-10 °C until complete as determined by hplc (2 hours). The pH of the reaction mixture was adjusted to 2.0 using 6 N hydrochloric acid solution while maintaining 5-10 °C. The product was extracted from solution with ethyl acetate (3 \times 500 mL). The ethyl acetate extracts were combined, dried using sodium sulfate, and concentrated to dryness to give 141.51 grams (100%) of The desired compound: 1H nmr (DMSO-d₆): δ 1.32-1.37 (m, 15H), 1.57-1.75 (m, 4H), 2.51-2.58 (m, 2H), 4.23-4.27 (m, 1H), 6.85 (s, broad, 1H), 7.15-7.28 (m, 5H), 7.42 (d, 1H), 12.5 (s, broad, 1H). ¹³C nmr (DMSO- d_6): δ 26.31, 27.85, 29.00, 31.86. 35.60, 52.53, 56.60, 78.95, 126.52, 129.10, 142.69, 155.06, 174.40, 175.17. MS (FIA) m/z $379.5 ([M+H]^{+})$. IR (KBr, cm⁻¹) 1641.98, 1692.22 (amides),

1719.72 (C=O). $\left[\alpha\right]^{20}_{D} = -5.73$ (C = 10.48 mg / 1mL, MeOH). Anal. Calcd. for $C_{20}H_{30}N_{2}O_{5}$: C, 63.47; H, 7.99; N, 7.40. Found: C, 63.25; H, 7.84; N, 7.46.

Preparation 1L

 $N ext{-Methyl morpholine}$ (4.79 mL, 2 eq, 47.3 mm) was added to a stirred slurry of N-Boc-a-aminoisobutyric acid (4.43 g, 21.7 mm, 1 eq) and 3.89 g (21.7 mm, 1.0 eq) of 2-chloro-(4,6)-dimethoxy-1,3,5-triazine (CDMT) in 100 mL diethyl ether. After stirring the reaction mixture at ambient temperature for 1.5 hours, D-tryptophan ester hydrochloride was added. After stirring overnight, the reaction mixture was quenched by the addition of 150 mL of 10% aqueous citric acid solution. The layers were separated and the ether layer was washed with 50 mL of saturated sodium bicarbonate solution and 50 mL of water. Lithium hydroxide (2.43 g, 5 eq) was dissolved in 100 ml of water and the solution was added to the diethyl ether solution and stirred vigorously for 4 hours at room temperature. The layers were separated and the pH of the aqueous layers was adjusted to 5.6 with 1M HCl. was then adjusted to 3.95 with 10% citric acid solution and the aqueous layer was extracted with 100 mL of ethyl The ethyl acetate layers were washed with acetate. brine, dried over magnesium sulfate and filtered. volatiles were removed under vacuum to give 82 % yield of the desired product as a white foam. 1H-NMR consistent with structure.

Preparation 2A



To a solution of 4-methoxyphenylacetic acid (98 g, 590 mmol) in absolute ethanol (300 mL) was added of ptoluenesulfonic acid (20 g, 105 mmol). The reaction mixture was heated to reflux and maintained at that temperature for 5 h then cooled to room temperature and concentrated to dryness. The resulting oil was purified by flash chromatography (silica gel, 20% ethyl acetate/hexanes) to give 102 g (89%) of the desired product as a colorless oil: $^1\text{H-NMR}$ (d, DMSO) 1.17 (t, J = 8.7 Hz, 3H), 3.56 (s, 2H), 3.73 (s, 3H), 4.05 (q, J = 7.2 Hz, 2H), 6.87 (d, J = 8.7 Hz, 2H), 7.17 (d, 8.7 Hz, 2H); MS (ion spray) 195.3 (M+1); Anal. Calc'd for $C_{11}H_{14}O_{3}$: C, 68.02; H, 7.27. Found: C, 67.95, 7.17.

Preparation 2B

To a solution of the product of Preparation 2A (40 g, 200 mmol) in carbon tetrachloride (500 mL) was added N-bromosuccinimide (37 g, 206 mmol) and hydrobromic acid (4 drops of 48% aqueous solution). The resulting mixture was heated to reflux and maintained at that temperature for 5 h then cooled to room temperature, filtered, and concentrated. The resulting oil was purified by flash chromatography (silica gel, chloroform) to give 51.1 g (94%) of the desired product as a colorless oil: ¹H-NMR (d, DMSO) 1.19 (t, J = 8.4 Hz, 3H), 3.77 (s, 3H), 4.18 (m, 2H), 5.88 (s, 1H), 6.95 (d, J = 8.4 Hz, 2H), 7.50 (d, J = 8.4 Hz, 2H); MS (FD) 272, 274 (M+); Anal. Calc'd for C₁₁H₁₃BrO₃: C, 48.37; H, 4.80. Found: C, 48.52, 4.77.

Preparation 3

To a solution of the product of Preparation 2B (49.5 g, 181 mmol) stirring in dimethylformamide (500 mL) at room temperature was added 4-nitroimidazole (20.5 g, 181 mmol) and potassium carbonate (75 g, 543 mmol). After 16 h, the reaction was filtered and concentrated. The resulting oil was partitioned between ethyl acetate and water and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The resulting oil was purified by flash chromatography (silica gel, 30-70% ethyl acetates/hexanes gradient) to yield 33.6 g (61%) of

the desired product as an orange oil that solidifies upon standing: ^{1}H -NMR (d, DMSO) 1.17 (t, J = 7.2 Hz, 3H), 3.78 (s, 3H), 4.25 (q, J = 7.2 Hz, 2H), 6.57 (s, 1H), 7.02 (d, J = 8.7 Hz, 2H), 7.46 (d, J = 8.7 Hz, 2H), 7.92 (s, 1H), 8.38 (s, 1H); MS (ion spray) 306 (M+1); Anal. Calc'd for $C_{14}H_{15}N_{3}O_{5}$: C, 55.08; H, 4.95; N, 13.76. Found: C, 54.93; H, 4.89; N, 13.82.

Preparation 4

To a slurry of 10% palladium on carbon (6.0 g) in tetrahydrofuran (30 mL) was added a slurry of the product of Preparation 3 (8.4 g, 27.5 mmol) in tetrahydrofuran The reaction mixture placed under a hydrogen atmosphere (40 mm Hg) using a Parr apparatus until the reduction was complete then filtered through celite. the resulting solution stirring at room temperature was added the product of Preparation 1d (10.5 g, 27.5 mmol), 1-hydroxybenzotriazole (4.1 g, 30.3 mmol) dimethylaminopropyl)-3-ethylcarbodiimide (6.3 30.3 mmol). After 16 h, the reaction mixture was concentrated and the resulting oil was slurried in ethyl acetate and filtered. The solution was diluted with water and then extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The resultant crude material was purified by flash chromatography (silica gel, 3% methanol/chloroform) to give 14.4 g (83%) of the desired product as a tan foam: 1H-NMR (d, DMSO) 1.78 (t, J

= 7.2 Hz, 3H), 1.27-1.32 (m, 15H), 3.60 (m, 1H), 3.67 (m, 1H), 3.76 (s, 3H), 4.20 (d, J = 7.2 Hz, 2H), 4.44 (d, J = 3.0 Hz, 2H), 4.57 (m, 1H), 6.35 (s, 1H), 6.97 (d, J = 7.2 Hz, 2H), 7.20-7.35 (m, 10H), 7.40 (m, 1H), 7.52 (s, 1H); MS (ion spray) 638 (M+1); Anal. Calc'd for $C_{33}H_{43}N_5O_8$: C, 62.15; H, 6.80; N, 10.98. Found: C, 62.41; H, 6.85; N, 11.09.

Preparation 5

To a solution of the product of Preparation 4 (14.4 g, 23 mmol) stirring in dioxane (150 mL) at room temperature was added a solution of of lithium hydroxide (0.65 g, 27.6 mmol) in water (75 mL). After 20 min, the reaction mixture was acidified to pH = 2.9 with 1 N hydrochloric acid. To the resulting solution was added water and ethyl acetate the mixture was extracted with ethyl The combined organic extracts were washed with brine, dried over sodium sulfate and concentrated to yield 13.0 g (93%) of the desired product as a yellow foam: ¹H NMR (d, DMSO) 1.25-1.40 (m, 15H), 3.65-3.70 (m, 2H), 3.76 (s, 3H), 4.44 (d, J = 3.4 Hz, 2H), 4.57 (m, 1H), 6.20 (s, 1H), 6.97 (d, J = 3.4 Hz, 2H), 7.15-7.35(m, 10H), 7.42 (m, 1H), 7.53 (s, 1H), 10.2 (s, 1H); MS (ion spray) 610.7 (M+1); Anal. Calc'd for $C_{31}H_{39}N_5O_8$: C, 61.07; H, 6.45; H, 11.49. Found: C, 60.90; H, 6.43; N, 11.32.

Preparation 6

To a solution of the product of Preparation 5 g,13.0 mmol) stirring in dimethylformamide (150 mL) at room temperature was added 4-methylpiperidine (1.6 mL, 13.0 mmol), 1-hydroxybenzotriazole (2.0 g, 14.3 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide g, 14.3 mmol). After 16h, the reaction mixture was filtered and concentrated. The resulting material was partitioned between ethyl acetate and water and extracted with ethyl acetate. The combined organic extracts were washed with dried brine, over sodium sulfate, filtered concentrated to dryness. The resulting crude material was purified by flash chromatography (silica gel, methanol/ chloroform) to yield 7.65 g (85%) of desired product as a yellow foam: 1H-NMR (d, DMSO) 0.2 (m, 1H), 0.50 (d, J = 6.0 Hz, 1.5 H), 0.80 (d, J = 6.0 Hz, 1.5 H)Hz, 1.5 H), 1.05 (m, 1H), 1.22-1.45 (m, 15H), 1.50-1.65 (m, 4H), 2.65 (m, 1H), 3.00 (m, 1H), 3.55 (m, 1H), 3.65 (m, 1H), 3.75 (s, 3H), 4.37 (m, 1H), 4.40-4.50 (m, 2H),4.60 (m, 1H), 6.62 (d, J = 13 Hz, 1H), 6.98 (t, J = 9.4Hz, 2H), 7.10-7.45 (m, 11H), 10.15 (br s, 1H); MS (ion spray) 691.3 (M+1); Anal. Calc'd for $C_{37}H_{50}N_6O_7$ 0.6 H_2O : C, Found: C, 63.25; H, 7.03; 63.34; H, 7.35; N, 11.98. 11.87.

Examples 1 and 2

131


To a solution of the product of Preparation 6 (7.26 g, 10.5 mmol) stirring in dichloromethane (25 mL) at room temperature was added trifluoroacetic acid (10 mL). After 4 h, the reaction mixture was poured into a saturated solution of sodium bicarbonate extracted with chloroform. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated to yield 6.12 g (99%) of the free base as a tan foam. The diastereomeric material (3.0 g) was chromatographed on an 8 x 15 cm Prochrom column packed with Kromosil CHI-DMP chiral phase using an eluent mixture of 3A alcohol (13% by v), dimethylethylamine (0.2% by v) in heptane at a flow rate of 250 mL/min to provide the individual diastereomers in pure form:

Example 1. Isomer 1 To a solution of the purified isomer in ethyl acetate was added a saturated solution of hydrochloric acid in diethyl ether. The resulting slurry was concentrated to dryness to yield 1.1 g (37%) of the desired product as a white solid: ¹H NMR (d, DMSO) 0.50 (d, J = 6.0 Hz, 1.5 H), 0.80 (d, J = 6.0 Hz, 1.5 H), 1.16 (m, 1H), 1.35 (m, 1H), 1.50-1.70 (m, 8H), 2.60-2.70 (m, 2H), 3.03 (m, 1H), 3.65-3.80 (m, 6H), 4.40 (m, 1H), 4.53 (s, 2H), 4.75 (m, 1H), 6.90-7.08 (m, 3H), 7.25-7.45 (m, 9H), 8.20-8.40 (m, 4H), 8.61 (d, J = 7.5 Hz, 1H), 11.15 (br s, 1H); $t_R = 7.93$ min; MS (ion spray) 591.6 (M+1); Anal. Calc'd for $C_{12}H_{42}N_6O_5$ 2HCl: C, 57.92; H, 6.69; N, 12.66. Found: C, 57.72; H, 6.47; N, 12.42.

132

Example 2. Isomer 2 To a solution of the purified isomer in ethyl acetate was added a saturated solution of hydrochloric acid in diethyl ether. The resulting slurry was concentrated to yield 0.98 g (33%) of the desired product as a white solid: ^{1}H NMR (d, DMSO) 0.50 (d, J = 6.0 Hz, 1.5 H), 0.80 (d, J = 6.0 Hz, 1.5 H), 1.16 (m, 1H), 1.35 (m, 1H), 1.50-1.70 (m, 8H), 2.60-2.70 (m, 2H), 3.03 (m, 1H), 3.65-3.80 (m, 6H), 4.40 (m, 1H), 4.53 (s, 2H), 4.75 (m, 1H), 6.90-7.08 (m, 3H), 7.25-7.45 (m, 9H), 8.20-8.40 (m, 4H), 8.61 (d, J = 7.5 Hz, 1H), 11.15 (br s, 1H); $t_R = 11.78$ min; MS (ion spray) 591.6 (M+1); Anal. Calc'd for $C_{32}H_{42}N_6O_5$ 2.2HCl: C, 57.29; H, 6.64; N, 12.53. Found: C, 57.23; H, 6.29; N, 12.57.

Preparation 7




Reaction of the product of Preparation 5 (0.6 g, 1.0 mmol), 4-(4-fluorobenzoyl) piperidine hydrochloride (0.24 g, 1.0 mmol), triethylamine (0.15 mL, 1.1 mmol), 1-hydroxybenzotriazole (0.16 g, 1.1 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol) in dimethylformamide (40 mL) as described in Preparation 6 gave 0.58 g (73%) of the desired product as a tan foam: $^1\text{H-NMR}$ (d, DMSO) 1.20-1.40 (m, 18H), 1.40-1.90 (m, 3H), 2.83 (m, 1H), 3.55-3.73 (m, 3H), 3.75 (s, 3H), 3.85 (m, 1H), 4.45 (d, J = 3.8 Hz, 2H), 4.60 (m, 1H), 6.65 (d, J = 10.93 Hz, 1H), 6.95-7.05 (m, 2H), 7.10-7.20 (m, 2H), 7.20-7.50 (m, 11H), 8.00-8.10 (m, 2H), 10.15 (br s, 1H); MS (FD) 798.7 (M+); Anal. Calc'd for

133

 $C_{43}H_{51}FN_6O_8$: C, 64.65; H, 6.43; N, 10.53. Found: C, 64.38; H, 6.48; N, 10.61.

Examples 3 and 4



Reaction of the product of Preparation 7 (0.53 g, 0.66 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.34 g (74%) of the desired mixture of diastereomers as a tan foam. This material (0.11 g) was purified by HPLC $(8 \times 15 \text{ cm})$ Prochrom column packed with Kromosil CHI-DMP of phase, eluent mixture 3A alcohol and dimethylethylamine in heptane) to provide the individual diastereomers which were converted to their repsective hydrochloride salts as desribed in Example 1.

Example 3. Isomer 1. 1 H-NMR (d, DMSO) 1.15-1.20 (m, 6H), 1.20-1.60 (m, 3H), 1.70 (m, 1H), 2.90 (m, 1H), 3.55-3.70 (m, 4H), 3.75 (s, 3H), 3.85 (m, 1H), 4.40 (m, 1H), 4.40-4.55 (m, 2H), 4.60 (m, 1H), 6.65 (d, J = 11 Hz, 1H), 7.00-7.05 (m 2H), 7.20 (m, 1H), 7.20-7.40 (m, 13H), 8.00-8.10 (m, 2H), 10.40 (br s, 1H); $t_R = 6.4$ min; MS (ion spray) 699.7 (M+1); Anal. Calc'd for $C_{38}H_{43}FN_6O_6$: C, 65.31; H, 6.20; N, 12.03. Found: C, 65.08; H, 6.18; N, 11.87.

Example 4. Isomer 2 ¹H-NMR (d, DMSO) 1.15-1.20 (m, 6H), 1.20-1.60 (m, 3H), 1.70 (m, 1H), 2.90 (m, 1H), 3.55-3.70 (m, 4H), 3.75 (s, 3H), 3.85 (m, 1H), 4.40 (m, 1H), 4.40-4.55 (m, 2H), 4.60 (m, 1H), 6.65 (d, J = 11 Hz, 1H), 7.00-7.05 (m 2H), 7.20 (m, 1H), 7.20-7.40 (m, 13H), 8.00-

134

8.10 (m, 2H), 10.40 (br s, 1H); $t_R = 8.0$ min; MS (high res) calc'd for $C_{38}H_{44}FN_6O_6$: 699.3306. Found: 699.3313.

Preparation 8

Reaction of the product of Preparation 7 (1.0 g, mmol), mmol), piperidine (0.17 mL, 1.7 1-1.9 hydroxybenzotriazole (0.25 mmol), 1-(3g, dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), and dimethylformamide (15 mL) as desribed in Preparation 6 gave 0.7 g (60%) of the desired product as a tan foam: $^{1}H-NMR$ (d, DMSO) 0.97 (m, 1H), 1.25-1.40 (m, 15H), 1.40-1.55 (m, 7H), 3.30-3.45 (m, 2H), 3.60 (m, 1H), 3.67 (m, 1H), 3.75 (s, 3H), 4.45 (d, J = 3.4 Hz, 2H), 4.57 (m, 1H), 6.62 (s, 1H), 6.98 (d, J = 8.7 Hz, 2H), 7.13 (m, 1H), 7.25-7.45 (m, 10H), 10.15 (br s, 1H); MS (ion spray) 677.5 (M+1); Anal. Calc'd for $C_{36}H_{48}N_6O_7$: C, 63.89; H, 7.15; N, 12.42. Found: C, 63.97; H, 6.99; N, 12.44.

135

Example 5

Reaction of the product of Preparation 8 (0.68 g, 1.0 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.6 g (93%) of the desired product as a white solid: $^{1}\text{H-NMR}$ (d, DMSO) 0.95 (m, 1H), 1.30-1.60 (m, 11H), 3.20-3.40 (m, 3H), 3.60-3.75 (m, 3H), 3.78 (s, 3H), 4.50-4.55 (m, 2H), 4.75 (m, 1H), 6.80 (s, 1H), 7.05 (d, J = 9.0 Hz, 2H), 7.25-7.35 (m, 7H), 7.37 (d, J = 8.7 Hz, 2H), 8.10 (m, 1H), 8.20-8.30 (m, 3H), 8.58 (d, J = 7.6 Hz, 1H), 11.00 (br s, 1H); MS (ion spray) 577.4 (M+1); Anal. Calc'd for $C_{31}H_{40}N_6O_5$ ·2.2HCl: C, 56.68; H, 6.48; N, 12.79. Found: C, 56.70; H, 6.64; N, 12.37.

Preparation 9

Reaction of the product of Preparation 5 (1.42 g, 2.3 mmol), d-proline methyl ester (0.3 g, 2.3 mmol), 1-hydroxybenzotriazole (0.35 g, 2.5 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.53 g, 2.5 mmol) in tetrahydrofuran (15 mL) as described in Preparation 6 gave 0.99 g (60%) of the desired product as a white foam: $^{1}\text{H-NMR}$ (d, DMSO) 1.25-1.40 (m, 18H), 1.75-1.90 (m, 2H), 2.40 (m, 1H), 3.30 (m, 1H), 3.60-3.80 (m, 7H), 4.40 (m, 1H), 4.45-4.50 (m, 2H), 4.57 (m, 1H), 6.50 (m, 1H), 6.95-7.05 (m, 2H), 7.10-7.40 (m, 11H), 10.20 (br s, 1H); MS (ion spray) 721.3 (M+1); Anal. Calc'd for $\text{C}_{37}\text{H}_{48}\text{N}_{6}\text{O}_{9}$: C, 61.65; H, 6.71; N, 11.66. Found: C, 61.42; H, 6.43; N, 11.65.

Example 6

Reaction of the product of preparation 9 (0.87 g, 1.2 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.58 g (70%) of the desired product: $^{1}\text{H-NMR}$ (d, DMSO) 1.40-1.60 (m, 6H), 1.75-1.95 (m, 3H), 2.20 (m, 1H), 2.95 (m, 1H), 3.60-3.80 (m, 9H), 4.40 (m, 1H), 4.50-4.55 (m, 2H), 4.75 (m, 1H), 6.70 (s, 1H), 7.00 (t, J = 8.7 Hz, 2H), 7.40-7.45 (m,

137

9H), 8.05 (m, 1H), 8.20-8.30 (m, 3H), 8.55 (m, 1H), 10.95 (m, 1H); MS (ion spray) 621.5 (M+1); Anal. Calc'd for $C_{32}H_{40}N_6O_7$ 2.3HCl: C, 54.55; H, 6.05; N, 11.93. Found: C, 54.46; H, 5.81; N, 11.79.

Preparation 10

To a suspension of 5% palladium on carbon (1.75 g) and tetrahydrofuran (120 mL) was added the product Preparation 3 (3.51 g, 11.5 mmol). The reaction mixture was placed under a hydrogen atmosphere (40 mm Hg) on a Parr apparatus for 2 h then filtered through celite. filtrate was subsequently added to a solution of the product of Preparation 1j (4.33 g, 11.5 mmol), 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (2.60 12.6mmol) and 1-hydroxybenzotriazole (1.72 g, 12.6 mmol) stirring in tetrahydrofuran (50 mL) at 0°C. After 16 h at room temperature, the reaction mixture was concentrated. The resulting residue was dissolved in ethyl acetate, filtered and the resulting filtrate concentrated. crude residue was purified by flash chromatography (silica gel, 90 용 ethyl acete/hexanes to methanol/ethyl acetate gradient) to give 4.5 g (62 %) the desired product as a light orange foam: 1H NMR consistent with structure; MS (IS) m/e 636 (M + 1). $(C_{34}H_{45}N_5O_7)$ C, H, N.

138

Pr paration 11

To a solution of the product of Preparation 10 (1.01 g, 1.59 mmol) stirring in tetrahydrofuran (30 mL) and water (15 mL) at room temperature was added lithium hydroxide (0.26 g, 6.30 mmol). After 25 min, the reaction mixture was concentrated and the resulting residue was diluted with water and extracted with diethyl ether. The aqueous extracts were acidified to pH 2-3 with 1N hydrochloric acid and then extracted with ethyl acetate. The combined extracts were washed with brine, dried with organic sodium sulfate and concentrated to provide 0.96 g (99 %) of the desired compound as a light tan foam that was used without further purification: ¹H NMR consistent with structure; MS (IS) m/e 608 (M + 1). Anal. (C₃₂H₄₁N₅O₇) C: calcd, 63.25; found, 62.68, H, N.

Preparation 12

To a solution of the product of Preparation 11 (0.93 g, 1.53 mmol) stirring in dichloromethane (25 mL) at room temperature was added N-methylmorpholine (0.20 mL, 1.83 mmol) and of 2-chloro-(4,6)-dimethoxy-1,3,5-triazine (0.35 g, 1.99 mmol). After 1 h, 4-methylpiperidine (0.20

139

mL, 1.68 mmol) was added and the resulting mixture was stirred room temperature for 2 h at which time 2-chloro-(4,6)-dimethoxy-1,3,5-triazine (0.10 g, 0.70 mmol) was After 1 h, the reaction mixture was concentrated and the resulting residue purified chromatography (silica gel, ethyl acetate/methanol gradient) to give the desired compound as a light yellow ¹H NMR consistent with solid foam (0.875 g, 83%): structure; MS (IS) m/e 689 (M + 1). Anal. ($C_{38}H_{52}N_6O_6$) C, H, N.


Example 7

To a solution of the product of Preparation 12 (0.77 g, 1.12 mmol) and anisole (0.13 mL, 1.13 mmol) stirring in dichloromethane (20 mL) at 0 C was added trifluoroacetic After 3-4 h, the reaction mixture was warmed to room temperature and then quenched by pouring over cold saturated aqueous sodium bicarbonate. The organic layer was collected and the aqueous layer was extracted twice with dichloromethane. The combined organic extracts were washed with aqueous sodium bicarbonate, water, brine, then dried over sodium sulfate and concentrated. resulting material was purified by flash chromatography (silica gel, 5% methanol/ 95% ethyl acetate gradient to 5% triethylamine/10% methanol/ 85% ethyl acetate) to (95 %) of the desired mixture provide 0.63 g diastereomers as an off-white solid foam. The mixture (190 mg) was resolved by chiral HPLC [Kromasil packing

140

material, 15% 3A alcohol/ 85% heptane 0.2% (w/ dimethylamine)] to provide the two desired diastereomers. To a solution of diastereomer 2 (65 mg) (retention time = 9.00 min) stirring in ethyl acetate (5 mL) was added saturated solution of hydrochloric acid in diethyl ether. The resulting white precipitate was collected by vacuum filtration and rinsed with diethyl ether to provide the desired compound (60 mg) as a white amorphous solid: NMR consistent with structure; MS (IS) m/e 589 (M + 1). Anal. $(C_{33}H_{44}N_6O_4^{\circ}2HC1)$ C, H, N.

Preparation 13



To a solution of Preparation 3 (3.00 g, 9.84 mmol) stirring in tetrahydrofuran (10 mL) and ethanol (5 mL) was added to sodium hydroxide (20 mL of a 5 N aqueous solution). The resulting smixture was stirred at ambient until temperature hydrolysis was complete subsequently acidified 2.0 with to Нq aqueous hydrochloric acid. The reaction mixture was extracted with ethyl acetate, dried over sodium sulfate, concentrated. The resulting carboxylic acid was combined with pyrrolidine (0.710)g, 10 mmol), hydroxybenzotriazole hydrate (1.35 g, 10 mmol) and 1,3dicyclohexylcarbodiimide (2.06 g, 10.0 mmol) stirring in tetrahydrofuran (100 mL) at room temperature. h, the mixture was concentrated, the residue slurried in ethyl acetate then filtered and concentrated. Purification by flash chromatography (silica chloroform/methanol) provided afford 2.74 g (84%) of the desired product: MS: (M+H) * 331.2; ¹H NMR (300 MHz,

WO 99/08697

141

DMSO- d_6) δ 8.19 (d, 1H, J = 1.51 Hz), 7.80(d, 1H, J = 1.51 Hz), 7.45 (d, 2H, J = 8.67 Hz), 7.02(d, 2H, J = 8.67 Hz), 6.58 (s, 1H), 3.77 (s, 3H), 3.75-3.60 (m, 1H) 3.45-3.30 (m, 2H), 2.90-2.75 (m, 1H)1.95-1.60 (m, 4H); Anal. Calcd. for $C_{16}H_{18}N_4O_4$: C, 58.18; H, 5.49;N, 16.96. Found: C, 58.44; H, 5.45; N, 16.87.

Preparation 14

The product of Preparation 13 (1.13 g, 3.42 mmol) was added to a mixture of 10% palladium/carbon (0.65 g) and palladium/black (0.15 g) in tetrahydrofuran (40 mL) the mixture shaken under hydrogen (38 psi) in a Parr After reduction was complete, the reaction apparatus. mixture was filtrated through celite and the filtrate immediately combined with 1-(3-dimethylaminopropyl)-3ethylcarbodiimide (0.71)3.45mmol), g, hydroxybenzotriazole (0.46 g, 3.40 mmol), the product of Preparation 1j (1.30 g, 3.44 mmol) and additional tetrahydrofuran (60 mL). After stirring overnight at ambient temperature, the mixture was concentrated and the residue slurried in ethyl acetate then filtered. filtrate was concentrated and the residue purified by flash chromatography(silica gel, chloroform/methanol) which afforded 1.50g (66%) of the desired product which was used without further purification.

Example 8

142

To a solution of the product of Preparation 14 (1.45 g, in dichloromethane (30 mL) was 2.20 mmol) triflouroacetic acid (10 mL). After 2 h, the mixture was concentrated and the residue treated with excess aqueous sodium bicarbonate and extracted. The combined organic extracts were concentrated and the resulting residue was flash chromatography purified by (silica chloroform/methanol) to provide 0.68 g of the desired product as a yellow solid: MS: (M+H)⁺ 561.3. ¹H NMR was consistent with product. Anal. Calcd. for C31H40N6O4'0.2 CHCl3: C, 64.11; H, 6.93; N, 14.38. Found: C, 64.19; H, 7.19; N, 14.50. The isomeric mixture (1.72 g) was separated as previously described in Example 7 to provide 0.64 g of isomer 1 (t_R = 7.50 min) and 0.49 g of isomer 2 $(t_R = 10.15 \text{ min})$. Isomer 2 (486 mg, 0.87 mmol) was dissolved in a minimal amount of ehtyl acetate and treated with an excess of saturated hydrochloric acid in ethyl acetate. Concentration and subsequent evaporation from diethyl ether allowed for recovery of 580 mg of an off-white solid: MS: $(M+H)^+$ 561.3, 562.4. ¹H NMR was consistent with product. Anal. Calcd. for C31H40N6O43.0 HCl: C, 55.57; H, 6.47; N, 12.54. Found: C, 56.40; H, 6.43; N, 12.20.

143

Preparation 15

The product of Preparation 13 (0.85 g, 2.57 mmol) was with 10% palladium/carbon (0.50 combined palladium/black (0.15 g) in tetrahydrofuran (40 mL)and the mixture shaken under a hydrogen atmosphere (38 mm Hg) in a Parr apparatus. After reduction was complete, the catalyst was removed by filtration through celite and the amine/tetrahydrofuran solution was immediately combined 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide(0.53 g, 2.57mmol), 1-hydroxybenzotriazole (0.35 g, 2.57 mmol), the product of Preparation 1L (1.00 g, 2.57 mmol) and additional tetrahydrofuran (60 mL). After stirring overnight at ambient temperature, the mixture concentrated and the residue slurried in ethyl acetate and filtered. The filtrate was concentrated and the residue purified by flash chromatography(silica gel, chloroform/methanol) which gave 1.62 g of the desired product which was used without further purification.

Example 9

The compound of Preparation 15 (1.57 g, 2.34 mmol) was dissolved in dichloromethane (25 mL) and triflouroacetic acid (10 mL) added. The resulting mixture was stirred at

ambient temperature for 2.5 h, concentrated, and the residue treated with excess aqueous sodium bicarbonate. The aqueous mixture was extracted with ethyl acetate and the combined organic extracts concentrated and dried. The silica gel chromatographed over residue was (chloroform/methanol) to provide 0.71 g (53 %) of the desired product: MS: (M+H) + 572.5. H NMR was consistent with product. Anal. Calcd. for C31H37N7O4.0.35 CHCl3: 61.38; H, 6.14; N, 15.98. Found: C, 61.36; H, 6.11; N, 16.08. The isomeric mixture (2.16 g) was separated as previously described in Example 7 to provide 1.10 g of isomer 1 (t_R = 10.34 min) and 0.80 g of isomer 2 (t_R = 13.70 min). The product derived from isomer 2 (0.80 g, 1.40 mmol) was dissolved in a minimal amount of ethyl acetate and the resulting solution treated with an excess of hydrochloric acid in ethyl acetate. The solution was then concentrated to provide 0.88 g (82 %) of the desired product as an off white solid: MS: (M+H) 572.3, 573.4. ¹H NMR was consistent with product. Anal. Calcd. for $C_{31}H_{37}N_7O_4$ 3.0 HCl: C, 54.67; H, 5.92; N, 14.40. Found: C, 54.25; H, 5.89; N, 13.35.

Preparation 16

To a solution of the product of Preparation 3 -411159-(5.75 g,18.9 mmol) stirring at room temperature in tetrahydrofuran (10 mL) was added sodium hydroxide (25 mL of a 5 N aqueous solution) along with water (15 mL) and ethanol (10 mL). After hydrolysis was complete, the mixture was acidified to pH 2.0 with aqueous hydrochloric

145

acid and extracted. The combined organic extracts were dried, filtered, and concentrated to give the desired product in quantitative yield as a tan solid: ^{1}H NMR (300 MHz,DMSO- d_{6}) δ 14.05-13.60 (bs, 1H), 8.34 (s, 1H) 7.90 (s, 1H), 7.45 (d, 2H, J = 8.67 Hz), 7.00 (d, 2H, J = 8.67 Hz), 6.42 (s, 1H), 3.77 (s, 3H). FDMS: 277 (M) $^{+}$ Anal. Calcd. for $C_{12}H_{11}N_{3}O_{5}$ 0.67 $H_{2}O$: C, 49.82; H, 4.30;N, 14.52. Found: C, 50.05; H, 4.01; N, 14.12.

146

Preparation 17

The compound of Preparation 16 (2.50 g,9.0 mmol) with aqueous dimethylamine (40%, 1.15 1-hydroxy-benzotriazole hydrate(1.22 mmol), g, mmol) and 1,3-dicyclohexylcarbodiimide (1.86 g, 9.0 mmol) in tetrahydrofuran (60 mL) and the mixture stirred at After 18 h, the mixture was ambient temperature. concentrated and the residue slurried in ethyl acetate and filtered. The filtrate was concentrated and the resulting residue purified by flash chromatography (silica gel, chloroform/methanol) to afford 1.83 g (67%) of the desired product: 1 H NMR (300 MHz, DMSO- d_{6}) δ 8.14 (s, 1H) 7.76 (s, 1H), 7.42 (d, 2H, J = 8.67 Hz), 7.00 (d, 2H, J = 8.67 Hz)2H, J = 8.67 Hz), 6.78 (s, 1H), 3.77 (s, 3H), 2.91 (2, 3H)3H), 2.85 (s, 3H). ESMS: $(M+H)^{+}$ 305.2.

Preparation 18

The compound of Preparation 17 (1.26 g, 4.14 mmol) was combined with 10% palladium/carbon (0.70)and palladium/black (0.15 g) in tetrahydrofuran(40 mL) and the mixture shaken under a hydrogen atmosphere (38 mm Hg) in a Parr apparatus. After reduction was complete, the catalyst was removed by filtration through celite and the immediately combined solution with 1-(3was

147

dimethylaminopropyl)-3-ethylcarbodiimide(0.82 g, mmol), 1-hydroxybenzotriazole mono-hydrate (0.54 g, 4.0 mmol), the product of Preparation 1j, (1.50 g, 3.97 mmol), and additional tetrahydrofuran (60 mL). After stirring ambient temperature, the mixture overnight at concentrated and the resulting residue slurried in ethyl acetate and filtered. The filtrate was concentrated and the residue purified by silica gel chromatography (chloroform/methanol) which provided 1.50g (57%) of the 635.6. ¹H NMR was desired product. MS: $(M+H)^{+}$ consistent with product. Anal. Calcd. for C34H46N6O6: C, 64.33; H, 7.30; N, 13.24. Found: C, 64.09; H, 7.09; N, 13.01.

Example 10

To a solution of the compound of preparation 18 (1.45 g, 2.29 mmol) stirring in dichloromethane (50 mL) at room temperature was added triflouroacetic acid (15 mL). After 3 h, the reaction mixture was concentrated and the residue treated with excess aqueous sodium bicarbonate. The mixture was extracted with ethyl acetate and the combined organic extracts were dried over sodium sulfate and concentrated. The residue was purified by flash chromatography (silica gel, chloroform/methanol) to give 0.73 g (60 %) if the desired product as a yellow solid: (60%. ESMS: (M+H)⁺ 535.4. ¹H NMR was consistent with product. Anal. Calcd. for C₂₉H₃₈N₆O₄·0.05 CHCl3: C, 64.54; H, 7.09; N, 15.54. Found: C, 64.28; H, 6.70; N, 15.35.

WO 99/08697

148

PCT/US98/17201

The diastereomeric mixture (2.35 g) was resolved by HPLC (8 x 15 cm Prochrom column packed with Kromosil CHI-DMP chiral phase using an eluent mixture of 3A alcohol and dimethylethylamine in heptane to provide the individual diastereomers in pure form (isomer 1, $t_R = 7.84$ min), isomer 2 (1.03 g, $t_R = 10.27$ min). To a solution of isomer 2 (1.03 g, 1.93 mmol) in ethyl acetate was added a saturated solution of hydrochloric acid in ethyl acetate. The resulting solution was concentrated, treated with diethyl ether and concentrated to provide 1.23 g of the desired product as an off white solid: ESMS: $(M+H)^+$ 535.3, 536.4. ¹H NMR was consistent with product. Anal. Calcd. for $C_{29}H_{38}N_6O_4$ 3.0 HCl: C, 54.08; H, 6.42; N, 13.05. Found: C, 54.12; H, 6.38; N, 12.86.

Preparation 19

The compound of preparation 17 (0.73 g, 2.38 mmol) was combined with 10% palladium/carbon (0.50)g) and palladium/black (0.10 g) in tetradyrofuran(40 mL)and the mixture shaken under hydrogen (38 mm Hg) in a Parr After reduction was complete, the catalyst removed by filtration through celite resulting solution was immediately combined dicyclohexylcarbodiimide (0.49 g, 2.38 mmol), hydroxybenzotriazole mono-hydrate (0.32 g, 2.37 mmol), the product of Preparation 1L (0.93 g, 2.39 mmol) and additional tetrahydrofuran (60 mL). After ambient temperature, the mixture overnight at concentrated and the residue slurried in ethyl acetate

149

and filtered. The filtrate was concentrated and the residue purified by silica gel chromatography(chloroform/methanol) to provide 0.76 g (50%) of the desired product as an off white solid which was used without further purification.

Example 11

To a solution of the compound of preparation 19 (0.74 g, stirring room temperature at dichloromethane (30 mL) was added triflouroacetic acid (10 mL). After 2 h, the mixture was concentrated and the residue treated with excess aqueous sodium bicarbonate. The resulting mixture was extracted with ethyl acetate and the combined organic extracts were concentrated. residue was purified by flash chromatography (silica gel, chloroform/methanol) to provide 0.23 g (37%) of 546.6. ¹H NMR was ESMS: (M+H) desired product: consistent with product. Anal. Calcd. for C29H35N7O4 0.25 CHCl3: C, 61.05; H, 6.17; N, 17.04. Found: C, 61.41; H, 16.52. The isomeric mixture (2.00 g) was separated as described in Example 10 to provide 0.73 g of isomer 1 (t_R = 9.85 min) and 0.82 g of isomer 2 (t_R = 12.87 min). To a solution of isomer 2 (0.82 g, 1.50 mmol) stirring in ethyl acetate and methanol was added a saturated solution of hydrochloric acid in ethyl acetate. The resulting mixture was concentrated to provide 0.84 g of the desired product: ESMS: (M+H) 546.2, 547.3. H NMR was consistent with product. Anal. Calcd. for C29H35N7O4'3.0

150

HCl: C, 53.18; H, 5.85; N, 14.97. Found: C, 53.73; H, 6.03; N, 14.04.

Preparation 20

Reaction of (3,4-dimethoxyphenyl)acetic acid (30.0 g, 153 mmol) and p-toluenesulfonic acid (6.5 g, 33.8 mmol) in absolute ethanol (200 mL) according to Preparation 1 gave 31.6 g (92%) of the desired product as a yellow oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 225 (M+1); Anal. Calc'd for $C_{12}H_{16}O_{4}$: C, 64.27; H, 7.19. Found: C, 64.08; H, 7.07.

Preparation 21

Reaction of the compound of Preparation 20 (1.5 g, 6.7 mmol), N-bromosuccinimide (1.3 g, 7.4 mmol), 2,2'-azobis(2-methylpropionitrile) (0.2 g) in carbon tetrachloride (30 mL)as decribed in Preparation 2 provided 2.03 g (100%) of the desired product as a clear oil: 1 H-NMR is consistent with structure; Anal. Calc'd for $C_{12}H_{15}BrO_4$: C, 47.54; H, 4.99. Found: C, 47.64; H, 5.17.

Reaction of the product of Preparation 21 (13.3 g, 44 mmol), 4-nitroimidazole (5.0 g, 44 mmol) and sodium hydride (2.1 g, 53 mmol) in tetrahydrofuran (400 mL) as desribed in Preparation 3 provided 22.6 g (85%) of the desired product as a tan oil. $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 334.1 (M-1); Anal. Calc'd for $C_{15}H_{17}N_3O_6$ 0.1CHCl3: C, 52.23; H, 4.96; N, 12.10. Found: C, 52.55; H, 4.81; N, 11.85.

Preparation 23

Hydrogenation of the compound of Preparation 22 (2.1 g, 6.3 mmol) with 10% palladium on carbon (1.5 g) in tetrahydrofuran (100 mL) followed by reaction with the product of Preparation 1d g, (2.4 6.3 mmol), hydroxybenzotriazole (0.97 6.9 1-(3g, mmol), dimethylaminopropyl)-3-ethylcarbodiimide (1.43 g, mmol) as described in Preparation 4 gave 2.08 g (49%) of the desired product as a red foam: 1H-NMR is consistent with structure; MS (ion spray) 668.4 (M+1).

Preparation 24

Reaction of the product of Preparation 23 (426814) (1.93 g, 2.9 mmol) and lithium hydroxide (0.08 g, 3.5 mmol) in

dioxane (50 mL) and water (25 mL) as described in Preparation 5 provided 1.68 g (91%) of the desired product as a tan foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 640.3 (M+1); Anal. Calc'd for $C_{32}H_{41}N_{5}O_{9}$: C, 60.08; H, 6.46; N, 10.95. Found: C, 60.31; H, 6.75; N, 10.65.

Preparation 25

Reaction of the product of Preparation 25 (426815) (0.8 g, 1.3 mmol), 4-methylpiperidine (0.16 mL, 1.3 mmol), 1-hydroxybenzotriazole (0.2 g, 1.43 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.3 g, 1.43 mmol) in dimethylformamide (20 mL) as desribed in Preparation 6 provided 0.56 g (60%) of the desired product as a tan foam: 1 H-NMR is consistent with structure; MS (ion spray) 721.5 (M+1); Anal. Calc'd for $C_{38}H_{52}N_6O_8$: C, 63.31; H, 7.27; N, 11.66. Found: C, 63.18; H, 7.30; N, 11.60.

Example 12

Reaction of the compound of Preparation 25 (0.5 g, 0.7 mmol and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.4 g (83%) of the

desired mixture of isomers as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 621.6 (M+1); Anal. Calc'd for $C_{33}H_{44}N_{6}O_{6}$ 2.3hydrochloric acid: C, 56.25; H, 6.62; N, 11.93. Found: C, 56.39; H, 6.33; N, 11.83.

Preparation 26

Reaction of the the product of Preparation 24 (0.8 g, 1.3 mmol), dimethylamine hydrochloride (0.11 g, 1.3 mmol), triethylamine (0.2 mL, 1.43 mmol), 1-hydroxybenzotriazole (0.2 g, 1.43 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.3 g, 1.43 mmol) in dimethylformamide (20 mL) as described in Preparation 6 gave 0.3 g (35%) of the desired product as a tan foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 667.4 (M+1); Anal. Calc'd for $C_{34}\text{H}_{46}\text{N}_{6}\text{O}_{8}$: C, 61.25; H, 6.95; N, 12.60. Found: C, 60.83; H, 6.48; N, 12.45.

Example 13


Reaction of the product of Preparation 26 (0.28 g, 0.42 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.21 g (78%) of the desired mixture of isomers as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (high res) calc'd for $C_{29}H_{39}N_6O_6$: 567.2931. Found: 567.2938. Anal. Calc'd for

WO 99/08697

154

 $C_{29}H_{36}N_6O_6$ 2hydrochloric acid: C, 54.46; H, 6.30; N, 13.14. Found: C, 54.67; H, 6.08; N, 13.00.

Preparation 27




Reaction of 4-trifluoromethylphenyl acetic acid (15.0 g, 73.4 mmol) and p-toluenesulfonic acid (2.8 g, 14.7 mmol) in absolute ethanol (100 mL) as described in Preparation 1 gave 16.3 g (95%) of the desired product as colorless oil: $^{1}\text{H-NMR}$ (d, DMSO) 1.18 (t, J = 7.0 Hz, 3H), 3.80 (s, 2H), 4.10 (q, J = 7.0 Hz, 2H), 7.49 (d, J = 7.9 Hz, 2H), 7.69 (d, J = 7.9 Hz, 2H); MS (FD) 232 (M+); Anal. Calc'd for $C_{11}H_{11}F_{3}O_{2}$: C, 56.90; H, 4.77. Found: C, 56.81; H, 4.85.

Preparation 28

Reaction of the product of Preparation of 27 (15.8 g, 68.0 mmol), N-bromosuccinimide (12.5 g, 70 mmol) and 48%-HBr (3 drops) in carbon tetrachloride (80 mL) as described in Preparation 2 gave 19.8 g (94%) of the desired product as a colorless oil: 1 H-NMR (d, DMSO) 1.19 (t, J = 7.2 Hz, 3H), 4.15-4.25 (m, 2H), 6.07 (s, 1H), 7.78 (s, 4H); MS (FD) 309, 311 (M+); Anal. Calc'd for $C_{11}H_{10}BrF_{3}O_{2}$: C, 42.47; H, 3.24. Found: C, 42.38; H, 3.13.

Preparation 29



Reaction of the product of Preparation 28 (51.8 g, 167 mmol), 4-nitroimidazole (18.8 g, 167 mmol), and potassium carbonate (51 g, 368 mmol) in N,N-dimethylformamide (600 mL) as described Preparation 3 gave 21.7 g (38%) of the desired product as a viscous orange oil: 1 H-NMR (d, DMSO) 1.19 (t, J = 7.2 Hz, 3H), 4.26 (q, J = 7.2 Hz, 2H), 6.80 (s, 1H), 7.76 (d, J = 8.3 Hz, 2H), 7.83 (d, J = 8.3 Hz, 2H), 8.01 (s, 1H), 8.51 (s, 1H); MS (ion spray) 344 (M+1); Anal. Calc'd for $C_{14}H_{12}F_{3}N_{3}O_{4}$: C, 48.99; H, 3.52; N, 12.24. Found: C, 49.03; H, 3.74; N, 11.96.

156

Pr paration 30

Hydrogenation of the product of Preparation 29 (8.5 g, 24.8 mmol) with 10% palladium on carbon (6.0 g) tetrahydrofuran (70 mL) followed by coupling with the product of Preparation 1d (9.5 g, 24.8 mmol), 1hydroxybenzotriazole (3.7 g, 27.3 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (5.6 g, mmol) as described in Preparation 4 gave 12.8 g (77%) of the desired product as a tan foam: ¹H-NMR (d, DMSO) 1.17 (t, J = 7.2 Hz, 3H), 1.25-1.35 (m, 15H), 3.60 (m, 1H),3.70 (m, 1H), 4.27 (q, J = 7.2 Hz, 2H), 4.44 (d, J = 2.6)Hz, 2H), 4.60 (m, 1H), 6.63 (s, 1H), 7.23-7.30 (m, 7H), 7.45 (m, 1H), 7.58-7.65 (m, 3H), 7.81 (d, J = 8.3 Hz, 2H), 10.25 (br s, 1H); MS (ion spray) 676.5 (M+1); Anal. Calc'd for $C_{33}H_{40}F_3N_5O_7$: C, 58.66; H, 5.97; N, 10.36. Found: C, 58.58; H, 6.17; N, 10.27.

Preparation 31

Reaction of the product of Preparation 30 (12.3 g, 18.2 mmol) and lithium hydroxide (0.52 g, 21.8 mmol) in dioxane (100 mL) and water (75 mL) as described in Preparation 5 gave 11.8 g (100%) of the desired product as tan foam: 1 H-NMR (d, DMSO) 1.20-1.35 (m, 15 H), 3.60 (m, 1H), 3.65 (m, 1H), 4.45 (d, J = 2.6 Hz, 2H), 4.60 (m,

1H), 6.46 (s, 1H), 7.15 (m, 1H), 7.20-7.35 (m, 6H), 7.42 (m, 1H), 7.57-7.65 (m, 3H), 7.79 (d, J=8.3 Hz, 2H), 10.25 (br s, 1H); MS (ion spray) 648.9 (M+1); Anal. Calc'd for $C_{31}H_{36}F_{3}N_{5}O_{7}$: C, 57.41; H, 5.60; N, 10.81. Found: C, 57.31; H, 5.59; N, 10.53.

Preparation 32

Reaction of the product of Preparation 31 (8.0 g, mmol), 4-methylpiperidine (1.5 mL, 12.3 mmol), 1hydroxybenzotriazole (1.83 g, 13.5 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (2.8 a, mmol) in N,N-dimethylformamide (150 mL) as described in Preparation 6 gave 7.33 g (81%) of the desired product as a tan foam: $^{1}H-NMR$ (d, DMSO) 0.78 (d, J = 6.0 Hz, 1.5H), 0.84 (d, J = 6.0 Hz, 1.5H), 0.95 (m, 1H), 1.25-1.35 (m, 16H), 1.50-1.70 (m, 4H), 2.65 (m, 1H), 3.60 (m, 1H), 3.67 (m, 1H), 3.80 (m, 1H), 4.35-4.50 (m, 3H), 4.60 (m, 1H),6.88 (d, J = 9.8 Hz, 1H), 7.20-7.30 (m, 7H), 7.45 (m, 1H), 7.48-7.55 (m, 2H), 7.60 (m, 1H), 7.75-7.85 (m, 2H), 10.25 (br s, 1H); MS (ion spray) 729 (M+1); Anal. Calc'd for $C_{37}H_{47}F_3N_6O_6$: C, 60.98; H, 6.50; N, 11.53. Found: C, 61.24; H, 6.44; N, 11.77.

158

Examples 14 and 15

Reaction of the product of Preparation 32 (7.0 g, 10.0 mmol) and trifluoroacetic acid (10 mL) in dichloromethane (25 mL) as described in Example 1 gave 5.62 g (93%) of the desired product (3.0 g) as a tan foam which was purified by HPLC (8 x 15 cm Prochrom column packed with Kromosil CHI-DMP chiral phase with an eluent mixture of 3A alcohol and dimethylethylamine in heptane) to give 1.5 g (45 %) of isomer 1 and 1.1 g (30 %) of isomer 2.

Example 14 (isomer 1): 1 H-NMR (d, DMSO) 0.25 (m, 1H), 0.76 (d, J = 6.4 Hz, 1.5H), 0.86 (d, J = 6.4 Hz, 1.5H), 1.00 (m, 1H), 1.45-1.70 (m, 8H), 2.65-2.75 (m, 2H), 3.15 (m, 1H), 3.65-3.80 (m, 3H), 4.40 (m, 1H), 4.51 (s, 2H), 4.75 (m, 2H), 7.10 (d, J = 12.8 Hz, 1H), 7.20-7.40 (m, 6H), 7.58 (d, J = 8.0 Hz, 1H), 7.67 (d, J = 8.0 Hz, 1H), 7.80-7.90 (m, 2H), 8.10 (br s, 1H), 8.20-8.35 (m, 3H), 8.55 (d, J = 7.5 Hz, 1H), 10.95 (br s, 1H); $t_R = 8.23$ min; MS (ion spray) 629.3 (M+1); Anal. Calc.d for $C_{32}H_{39}F_3N_6O_4$ 2HCl: C, 54.78; H, 5.89; N, 11.98. Found: C, 54.85; H, 5.71; N, 11.70.

Example 15 (isomer 2): 1 H-NMR (d, DMSO) 0.25 (m, 1H), 0.76 (d, J = 6.4 Hz, 1.5H), 0.86 (d, J = 6.4 Hz, 1.5H), 1.00 (m, 1H), 1.45-1.70 (m, 8H), 2.65-2.75 (m, 2H), 3.15 (m, 1H), 3.65-3.80 (m, 3H), 4.40 (m, 1H), 4.51 (s, 2H), 4.75 (m, 2H), 7.10 (d, J = 12.8 Hz, 1H), 7.20-7.40 (m, 6H), 7.58 (d, J = 8.0 Hz, 1H), 7.67 (d, J = 8.0 Hz, 1H), 7.80-7.90 (m, 2H), 8.10 (br s, 1H), 8.20-8.35 (m, 3H), 8.55 (d, J = 7.5 Hz, 1H), 10.95 (br s, 1H); $t_R = 10.77$ min; MS (ion spray) 629.3 (M+1); Anal. Calc.d for

159

 $C_{32}H_{39}F_{3}N_{6}O_{4}$ 2.2HCl: C, 54.22;H, 5.86; N, 11.85. Found: C, 54.15; H, 5.84; N, 11.64.

Preparation 33

Reaction of the product of Preparation 31 (0.6 g, 0.93 mmol), 4-(4-fluorobenzoyl)piperidine hydrochloride (0.23 g, 0.93 mmol), triethylamine (0.15 mL, 10.2 mmol), 1hydroxybenzotriazole (0.14 g, 1.02 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.21 g, 1.02 mmol)in dimethylformamide (30 mL) as described in Preparation 6 provided in 0.35 g (45%) of the desired product as a yellow foam: 1H-NMR (d, DMSO) 1.25-1.35 (m, 1.40-1.50 (m, 2H), 1.75 (m, 1H), 1.85 (m, 1H), 2.85-3.00 (m, 2H), 3.55-3.75 (m, 3H), 3.90 (m, 1H), 4.40-4.50 (m, 3H), 4.60 (m, 1H), 6.90 (m, 1H), 7.25-7.40 (m, 12H), 7.50-7.60 (m, 3H), 8.03-8.10 (m, 2H), 10.20 (br s, 1H); MS (ion spray) 837.4 (M+1); Anal. Calc'd for $C_{43}H_{48}F_{4}N_{6}O_{7}$: C, 61.71; H, 5.78; N, 10.04. Found: C, 61.53; H, 5.98; N, 9.95.

Example 16

Reaction of the product of Preparation 33 (0.34 g, 0.4 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.2 g (63%) of the

desired product as a yellow solid: $^{1}\text{H-NMR}$ (d, DMSO) 1.45-1.65 (m, 6H), 1.75 (m, 1H), 1.85 (m, 1H), 2.85-3.05 (m, 2H), 3.25 (m, 1H), 3.60-4.00 (m, 7H), 4.40-4.55 (m, 3H), 4.75 (m, 1H), 7.05 (d, J = 10.6 Hz, 1H), 7.25-7.40 (m, 8H), 7.55-7.70 (m, 2H), 7.75-7.85 (m, 2H), 8.00-8.10 (m, 2H), 8.15-8.25 (m, 3H), 8.50 (d, J = 7.2 Hz, 1H), 10.75 (br s, 1H); MS (ion spray) 737.0 (M+1); Anal. Calc'd for $C_{38}H_{40}F_{4}N_{6}O_{5}$ 2.4HCl: C, 55.37; H, 5.18; N, 10.20. Found: C, 55.39; H, 5.45; N, 10.07.

Preparation 34

Hydrogenation of the product of Preparation 29 (1.75 g, 5.1 mmol) with 10% palladium on carbon (1.4 g) tetrahydrofuran (60 mL) followed by reaction with the product of Preparation 1L (2.0 g, 5.1 mmol), 1g, 5.6 mmol) hydroxybenzotriazole (0.76 and 1 - (3 dimethylaminopropyl)-3-ethylcarbodiimide (1.16 g, mmol) as described in Preparation 4 gave 2.51 g (72%) of the desired product as a tan foam: 1H-NMR (d, DMSO) 1.15-1.35 (m, 18H), 3.05-3.15 (m, 2H), 4.25 (m, 2H), 4.65 (br s, 1H), 6.62 (s, 1H), 6.85 (m, 1H), 6.95-7.08 (m, 2H), 7.20-7.30 (m, 2H), 7.40-7.55 (m, 2H), 7.55-7.65 (m, 3H), 7.82 (d, J = 8.3 Hz, 2H), 10.20 (br s, 1H), 10.75 (br s, (ion spray) 685 (M+1); Anal. Calc'd for $C_{34}H_{39}F_{3}N_{6}O_{6}^{-1}H_{2}0: C, 58.11; H, 5.88; N, 11.96.$ 58.15; H, 5.59; N, 11.92.

161

Reaction of the product of Preparation 34 (2.2 g, 3.2 mmol) and lithium hydroxide (0.1 g, 3.9 mmol) in dioxane (50 mL) and water (25 mL) as described in Preparation 5 gave 2.1 g (100%) of the desired product as a tan foam: $^{1}\text{H-NMR}$ (d, DMSO), 1.15-1.35 (m, 15H), 3.05-3.15 (m, 2H), 4.65 (br s, 1H), 6.97 (s, 1H), 6.90 (m, 1H), 6.98-7.10 (m, 2H), 7.20-7.30 (m, 2H), 7.40-7.55 (m, 2H), 7.57-7.64 (m, 3H), 7.80 (d, J = 8.3 Hz, 2H), 10.20 (br s, 1H), 10.75 (br s, 1H), 13.80 (br s, 1H); MS (ion spray) 657.4 (M+1); Anal. Calc'd for $C_{32}H_{35}F_{3}N_{6}O_{6}$: C, 58.53; H, 5.37; N, 12.80. Found: C, 59.28; H, 5.17; N, 12.65.

Preparation 36

Reaction of the product of Preparation 35 (0.7 g, 1.1 mmol), 4-methylpiperidine (0.13 mL, 1.1 mmol), hydroxybenzotriazole (0.17 g, 1.2 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.26 mmol) in N,N-dimethylformamide (30 mL) as described in Preparation 6 provided 0.47 g (58%) of the desired product as a tan foam: $^{1}H-NMR$ (d, DMSO) 0.78 (d, J = 6.4) Hz, 1.5H), 0.86 (d, J = 6.3 Hz, 1.5H), 1.15-1.35 (m, 18H), 1.50-1.70 (m, 3H), 2.60-2.70 (m, 2H), 3.00-3.15 (m, 2H), 3.30 (m, 1H), 4.40 (m, 1H), 4.65 (m, 1H), 6.85-6.95 (m, 2H), 7.00-7.10 (m, 2H), 7.17-7.30 (m, 2H), 7.40-7.60(m, 4H), 7.75-7.85 (m, 2H), 10.20 (br s, 1H), 10.75 (br

162

s, 1H); MS (ion spray) 738.5 (M+1); Anal. Calc'd for $C_{38}H_{46}F_{3}N_{7}O_{5}$ 1H₂0: C, 60.39; H, 6.40; N, 12.97. Found: C, 60.18; H, 6.21; N, 12.99.

Examples 17 and 18

Reaction of the product of Preparation 36 (4.8 g, 6.5 mmol) and trifluoroacetic acid (16 mL) in dichloromethane (40 mL) as described in Example 1 gave 2.0 g (44%) of the desired mixture as a tan foam. Purification by HPLC (8 x 15 cm Prochrom column packed with Kromosil CHI-DMP chiral phase with an eluent mixture of 3A alcohol (13% by v), dimethylethylamine (0.2% by v) in heptane at a flow rate of 250 mL/min) gave 0.5 g (12%) of isomer 1 and 0.4 g (9%) of isomer 2.

Example 17. (isomer 1) 1 H-NMR (d, DMSO) 0.77 (d, J = 6.5 Hz, 1.5H), 0.87 (d, J = 6.0 Hz, 1.5H), 1.00 (m, 1H), 1.32 (s, 3H), 1.50 (s, 3H), 1.50-1.70 (m, 2H), 2.72 (m, 1H), 3.00-3.30 (m, 4H), 3.75 (m, 1H), 4.05-4.33 (m, 3H), 4.20 (m, 1H), 4.78 (m, 1H), 6.94 (m, 3H), 7.20 (s, 1H), 7.30-7.40 (m, 2H), 7.55-7.70 (m, 2H), 7.75-8.00 (m, 4H), 8.05-8.15 (m, 2H), 8.50 (m, 1H), 10.86 (s, 1H), 11.05 (s, 1H); $t_R = 6.01 \text{ min}$; MS (ion spray) 638.2 (M+1).

Example 18. (isomer 2) 1 H-NMR (d, DMSO) 0.77 (d, J = 6.5 Hz, 1.5H), 0.87 (d, J = 6.0 Hz, 1.5H), 1.00 (m, 1H), 1.32 (s, 3H), 1.50 (s, 3H), 1.50-1.70 (m, 2H), 2.72 (m, 1H), 3.00-3.30 (m, 4H), 3.75 (m, 1H), 4.05-4.33 (m, 3H), 4.20 (m, 1H), 4.78 (m, 1H), 6.94 (m, 3H), 7.20 (s, 1H), 7.30-7.40 (m, 2H), 7.55-7.70 (m, 2H), 7.75-8.00 (m, 4H), 8.05-

163

8.15 (m, 2H), 8.50 (m, 1H), 10.86 (s, 1H), 11.05 (s, 1H); $t_R = 7.5 \text{ min}$; MS (ion spray) 638.2 (M+1).

Preparation 37

A mixture of the product of Preparation of 29 (11.1 g, 32.3 mmol) and 5% palladium on carbon (1.7 tetrahydrofuran (100 mL) was hydrogenated at 60 psi at room temperature using a Parr apparatus. After 1.5 h, the resulting brown solution was filtered through celite and concentrated to give 8.8 g (87%) crude oil which was used without purification. To a mixture of the amine stirring at O oC in tetrahydrofuran (20 mL) was added the product of Preparation 1j (10.6 g, 28.1 mmol) To this mixture was added 1tetrahydrofuran (30 mL). hydroxy-7-azobenzotriazole (4.0 g, 29.5 mmol) and 1,3dicyclohexylcarbodiimide (6.1 g, 29.5 mmol). The solution was allowed to warm to room temperature and the resulting mixture filtered after 3 days. The filtrate was concentrated and subsequently purified by flash chromatography (silica gel, 3.5% methanol/dichloromethane) to provide 12.1 g (64%) of the desired product as an orange solid: H-NMR (d, DMSO) 1.15 (t, J = 7 Hz, 3H), 1.18-1.32 (m, 15H), 1.35-1.70 (m, 4H),3.23 (m, 2H), 4.19 (q, J = 7 Hz, 2H), 4.31 (m, 1H), 6.58(s, 1H), 7.00 (br s, 1H), 7.05-7.22 (m, 6H), 7.41 (m, 1H), 7.52-7.58 (m, 3H), 7.75 (d, J = 8 Hz, 2H), 10.19 (br s, 1H); MS (ion spray) 674.7 (M+1); Anal. Calc'd for


164

 $C_{34}H_{42}F_3N_5O_6$: C, 60.61; H, 6.28; N, 10.39. Found: C, 60.44; H, 6.48; N, 10.36.

Preparation 38

To a solution of the product of Preparation 37 (12.0 g, 17.8 mmol)stirring in dioxane (20 ml)and water (20 ml) at room temperature was added lithium hydroxide (0.84 g, 35.6 mmol). After 90 min with intermittent sonication, the reaction was poured into a solution of sodium bisulfate(12 g/50 mL H₂O) and brine (20 mL) then extracted with ethyl acetate. The combined organic extracts were dried over sodium sulfate, filtered, and concentrated to provide 11.5 g (100%) of the desired product as a tan solid: $^{1}H-NMR$ (d, DMSO) 1.17-1.31 (m, 15 H), 1.40-1.70 (m, 4H), 2.45 (m, 2H), 4.33 (m, 1H), 6.40 (s, 1H), 7.00 (m, 1H), 7.05-7.23 (m, 6H), 7.40 (m, 1H), 7.55-7.71 (m, 1H)3H), 7.76 (d, J = 8 Hz, 2H), 10.25 (br s, 1H); MS (ion spray) 646.6 (M+1); Anal. Calc'd for C₃₂H₃₈F₃N₅O₆•0.7 H₂O: C, 58.39; H, 6.03; N, 10.64. Found: C, 58.52; H, 6.01; N, 9.87*.


165



To a solution of the product of Preparation 38 (6.0 g, 9.3 mmol)stirring at 0°C in dimethylformamide was added 9.3 dimethylamine hydrochloride (0.76 g, diethylcyanophosphonate mL, 9.3 mmol), (1.41)triethylamine (1.29 mL, 9.3 mmol). After 30 min, a second equivalent of dimethylamine hydrochloride, DECP and triethylamine were added. After 30 min, the reaction mixture was diluted with ethyl acetate (300 mL) washed with aqueous sodium bisulfate and brine. The organic extract was dried over sodium sulfate, filtered, and concentrated. The resulting crude material purified by radial chromatography (silica gel, methanol in dichloromethane) to give 4.7 g (75%) of the desired product as a tan foam: 1H-NMR (d, CDCl3) 1.25(s, 9H), 1.42(s, 6H), 1.60-1.80 (m, 4H), 1.90 (br s, 1H), 2.57 (m, 2H), 2.98 (s, 6H), 4.48 (m, 1H), 7.05-7.21 (m, 6H), 7.50 (m, 1H), 7.62-7.76 (m, 5H), 8.93 (br s, 1H), 10.93 (br s, 1H); MS (ion spray) 673.7 (M+1).

166

Examples 19 and 20



To the product of Preparation 39 (4.7 g, 7.0 mmol) was stirred at room temperature in a saturated solution of hydrochloric acid in glacial acetic acid (30 mL). After 90 min, the mixture was concentrated. The resulting material diluted with ethyl acetate and extracted with aqueous sodium bicarbonate. The organic extract was dried over sodium sulfate, filtered, and concentrated to give 3.7 g (93%) of an orange solid. MS (ion spray) 573.4 (M+1). The diastereomers (3.4 g) were separated by chiral chromatography using a Kromasil-CHI normal phase column to provide 1.40 g (41%) of isomer 1 and 1.26 g (37%) of isomer 2. The individual isomers were dissolved in a saturated solution of hydrochloric acid in glacial acetic acid (4 mL) and subsequently concentrated to provide the desired products as tan solids:


Example 19 (isomer 1) 1 H-NMR (d, DMSO) 1.41 (s, 3H), 1.42 (s, 3H), 1.51-1.73 (m, 4H), 2.53 (m, 2H), 2.82 (s, 3H), 2.84 (s, 3H), 4.39 (m, 1H), 6.91 (s, 1H), 7.10 (m, 3H), 7.18-7.29 (m, 3H), 7.55 (d, J = 8 Hz, 2H), 7.78 (d, J = 8 Hz, 2H), 7.91 (br s, 1H), 8.15 (br s, 3H), 8.38 (d, J = 7.5 Hz, 1H), 10.78 (br s, 1H); MS (ion spray) 573.4 (M+1); Anal. Calc.d for $C_{29}H_{35}F_{3}N_{6}O_{3}$ 2.3HCl: C, 53.06; H, 5.73; N, 12.80. Found: C, 52.90; H, 5.66; N, 12.70.

Example 20. (isomer 2) ¹H-NMR (d, DMSO) 1.42 (s, 6H), 1.51-1.73 (m, 4H), 2.53 (m, 2H), 2.82 (s, 3H), 2.84 (s, 3H), 4.39 (m, 1H), 6.91 (s, 1H), 7.10 (m, 3H), 7.18-7.29 (m, 3H), 7.55 (d, J = 8 Hz, 2H), 7.78 (d, J = 8 Hz, 2H), 7.91 (br s, 1H), 8.15 (br s, 3H), 8.38 (d, J = 7.5 Hz,

167


1H), 10.78 (br s, 1H); MS (ion spray) 573.4 (M+1); Anal. Calc.d for $C_{29}H_{35}F_{3}N_{6}O_{3}$ 2HCl: C, 53.96;H, 5.78; N, 13.02. Found: C, 53.84; H, 5.71; N, 12.93.

Preparation 40



Reaction of (2-fluoro-4-trifluoromethyl)phenylacetic acid (20.0 g, 90 mmol) and p-toluenesulfonic acid (5.0, 26 mmol) in absolute ethanol (200 mL) as described in Preparation 1 provided 22.5 g (100%) of the desired product as a colorless oil: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 250 (M+); Anal. Calc'd for $C_{11}H_{10}F_4O_2$: C, 52.81; H, 4.03. Found: C, 52.94; H, 3.94.

Preparation 41



Reaction of the product of Preparation 40 (16.8 g, 67 mmol), N-bromosuccinimide (12.3 g, 69 mmol) and 48% HBr (3 drops) in carbon tetrachloride (170 mL) as described in Preparation 2 gave 22.05 g (100%) of the desired product as a colorless oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 328, 330 (M+); Anal. Calc'd for $C_{11}H_{9}BrF_{4}O_{2}$: C, 40.15; H, 2.76. Found: C, 40.00; H, 2.77.

Preparation 42



Reaction of the product of Preparation 41 (21.4 g, 65 mmol), 4-nitroimidazole (8.8 g, 78 mmol) and potassium

168

carbonate (26.8 g, 195 mmol) in dimethylformamide (300 mL) as described in Preparation 3 gave 3.75 g (16%) of the desired product as a tan oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 362.2 (M+1); Anal. Calc'd for $C_{14}H_{11}F_{4}N_{3}O_{4}$: C, 46.55; H, 3.07; N, 11.63. Found: C, 47.13; H, 3.49; N, 11.37.

Preparation 43

Reduction of the product of Preparation 42 (2.88 g, 8.0 mmol) with 10% palladium on carbon (1.45 g) in tetrahydrofuran (60 mL) follwed by coupling with the product of Preparation 1d gave (3.0 g, 8.0 mmol), 1-hydroxybenzotriazole (1.2 g, 8.8 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (8.8 mmol) as described in Preparation 4 gave 2.85 g (51%) of the desired product as a tan foam: $^1\text{H-NMR}$ is consistent with structure; MS (ion spray) 694.4 (M+1); Anal. Calc'd for $C_{33}H_{39}F_4N_5O_7$: C, 57.14; H, 5.67; N, 10.10. Found: C, 57.28; H, 5.59; N, 10.09.

Preparation 44

Reaction of the product of Preparation 43 (2.64 g, 3.8 mmol) and lithium hydroxide (0.11 g, 4.6 mmol) in dioxane (50 mL) and water (25 mL) as described in Preparation 5

169


gave 2.53 g (100%) of the desired product as a tan foam. $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 664.4 (M+1).

Preparation 45

Reaction of the product of Preparation 44 (0.8 g, 1.2 mmol), 4-methylpiperidine (0.14 mL, 1.2 mmol), 1-hydroxybenzotriazole (0.18 g, 1.32 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.27 g, 1.32 mmol) in dimethylformamide (40 mL) as described in Preparation 6 gave 0.63 g (70%) of the desired product as a yellow foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 747.4 (M+1).

170

Example 21



Reaction of the product of Preparation 46 (0.54 g, 0.72 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.4 g (77%) of the desired mixture of isomers as a white solid. $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 647.6 (M+1); Anal. Calc'd for $C_{32}H_{38}F_{4}N_{6}O_{4}$ 2HCl: C, 53.41; H, 5.60; N, 11.68. Found: C, 53.34; H, 5.84; N, 11.65.

Preparation 46


Reaction of the product of Preparation 44 (0.8 g, 1.2 mmol), dimethylamine hydrochloride (0.1 g, 1.2 mmol), triethylamine (0.19 mL, 1.3 mmol), 1-hydroxybenzotriazole (0.18 g, 1.3 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.27 g, 1.3 mmol) in dimethylformamide (40 mL) as described in Preparation 6 gave 0.48 g (58%) of the desired product: $^1\text{H-NMR}$ is consistent with structure; MS (ion spray) 693.4 (M+1); Anal. Calc'd for $C_{33}H_{40}F_4N_6O_6$: C, 57.22; H, 5.82; N, 12.13. Found: C, 57.48; H, 5.74; N, 12.02.

Example 22

171

Reaction of the product of Preparation 46 (0.43 g, 0.62 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.25 g (60%) of the desired product as a mixture of diastereoisomers. $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 593.9 (M+1); Anal. Calc'd for $C_{28}H_{32}F_{4}N_{6}O_{4}$ 2hydrochloric acid: C, 50.53; H, 5.15; N, 12.62. Found: C, 50.25; H, 5.20; N, 12.35.

Preparation 47




Reaction of 4-fluorophenylacetic acid (15.0 g, 97.0 mmol), p-toluenesulfonic acid (2.0 g, 10.5 mmol) and absolute ethanol (100 mL) as described in Preparation 1 gave 15.4 g (87%) of the desired product as a colorless oil: $^{1}\text{H-NMR}$ (d, DMSO) 1.17 (t, J = 7.2 Hz, 3H), 3.66 (s, 2H), 4.06 (q, J = 7.2 Hz, 2H), 7.10-7.20 (m, 2H), 7.25-7.35 (m, 2H); MS (FD) 182 (M+); Anal. Calc'd for $C_{10}H_{11}FO_{2}$: C, 65.92; H, 6.09. Found: C, 65.67; H, 5.96.

172

Preparation 48

Reaction of the product of Preparation 47 (14.9 g, 82 mmol), N-bromosuccinimide (14.9 g, 84.5 mmol) and 48% HBr (4 drops) in carbon tetrachloride (80 mL) as described in Preparation 2 gave 18.3 g (85%) of the desired product as a colorless oil: 1 H-NMR (d, DMSO) 1.19 (t, J = 7.2 Hz, 3H), 4.15-4.25 (m, 2H), 5.95 (s, 1H), 7.15-7.30 (m, 2H), 7.56-7.70 (m, 2H); MS (FD) 260, 262 (M+); Anal. Calc'd for C_{10} H₁₀BrFO₂: C, 46.00; H, 3.96. Found: C, 46.10; H, 3.95.

Preparation 49



Reaction of the product of Preparation 48 (68 g, 260 mmol), 4-nitroimidazole (35.0 g, 312 mmol) and potassium carbonate (108 g, 780 mmol) in dimethylformamide (300 mL) as described in Preparation 3 gasve 39.8 g (52%) of the desired product as an orange oil: $^{1}\text{H-NMR}$ (d, DMSO) 1.83 (t, J = 7.2 Hz, 3H), 4.25 (q, J = 7.2 Hz, 2H), 6.66 (s, 1H), 7.25-7.35 (m, 2H), 7.55-7.65 (m, 2H), 7.95 (d, 1.13 Hz, 1H), 8.44 (d, J = 1.5 Hz, 1H); MS (ion spray) 294.2 (M+1); Anal. Calc'd for $C_{13}H_{12}FN_3O_4$: C, 53.24; H, 4.12; N, 14.33. Found: C, 53.51; H, 4.07; N, 14.42.

173

Preparation 50

Reduction of the product of Preparation 49 (8.9 g, 30.3 10% palladium on carbon (6.0 tetrahydrofuran (120 mL) follwed by coupling with the of Preparation 1d (11.4 g, 30 mmol), hydroxybenzotriazole (4.5 g, 33 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (6.8 g, 33 mmol) as described in Preparation 4 gave 10.8 g (58%) of the desired product as a tan foam: 1H-NMR (d, DMSO) 1.18 (t, J = 7.2 Hz, 3H, 1.25-1.35 (m, 15H), 3.60 (m, 1H), 3.70 (m,1H), 4.25 (q, J = 7.2 Hz, 2H), 4.44 (d, J = 2.6 Hz, 2H), 4.60 (m, 1H), 6.47 (s, 1H), 7.20-7.40 (m, 9H), 7.40-7.50 (m, 3H), 7.56 (s, 1H), 10.25 (br s, 1H); MS (ion spray) 626.1 (M+1); Anal. Calc'd for $C_{32}H_{40}FN_5O_7$: C, 61.43; H, 6.44; N, 11.19. Found: C, 61.63; H, 6.42; N, 11.26.

Preparation 51

Reaction of the product of Preparation 50 (10.5 g, 17.0 mmol) and lithium hydroxide (0.48 g, 20.4 mmol) in dioxane (200 mL) and water (100 mL) as described in Preparation 5 gave 10.1 g (100%) of the desired product as a tan foam: 1 H-NMR (d, DMSO) 1.25-1.40 (m, 15H), 3.35 (br s, 1H), 3.60 (m, 1H), 3.70 (m, 1H), 4.44 (d, J = 2.6

174

Hz, 2H), 4.60 (m, 1H), 6.33 (s, 1H), 7.20-7.35 (m, 9H), 7.40-7.50 (m, 3H), 7.56 (s, 1H), 10.20 (br s, 1H); MS (ion spray) 598.5 (M+1); Anal. Calc'd for $C_{30}H_{36}FN_5O_7$: C, 60.29; H, 6.07; N, 11.72. Found: C, 60.38; H, 6.29; N, 11.49.

Preparation 52

Reaction of product of Preparation 51 (9.2 g, 15.4 mmol), 1-4-methylpiperidine (1.83 mL, 15.4 mmol), 17 1-(3hydroxybenzotriazole (2.3)g, mmol) and dimethylaminopropyl)-3-ethylcarbodiimide (3.5 g, 17 mmol) in dimethylformamide (100 mL) as described in Preparation 6 gave 9.7 g (93%) of the desired product as a tan foam: 1 H-NMR (d, DMSO) 0.76 (d, J = 6.1 Hz, 1.5H), 0.86 (d, J = 6.1 Hz, 1.5H), 1.00 (m, 1H), 1.20-1.40 (m, 15H), 1.45-1.70 (m, 3H), 2.55-2.70 (m, 2H), 3.05 (m, 1H), 3.60 (m, 1H), 3.65-3.75 (m, 2H), 4.40 (m, 1H), 4.44 (d, J = 2.6Hz, 2H), 4.60 (m, 1H), 6.73 (d, J = 11.3 Hz, 1H), 7.15-7.35 (m, 9H), 7.35-7.50 (m, 4H), 10.20 (br s, 1H). (ion spray) 679.6 (M+1); Anal. Calc'd for $C_{36}H_{47}FN_6O_6$: C, 63.70; H, 6.98; N, 12.38. Found: C, 63.44; H, 6.86; N, 12.22.

175

Example 23 and 24

Reaction of the product of Preparation 52 (9.7 g, 14.3 mmol) with trifluoroacetic acid (16 mL) in dichloromethane (40 mL) as described in Example 1 gave 6.8 g (73%) of the desired product as a mixture of diastereoisomers. The mixture (3.2 g) was purified by HPLC (8 x 15 cm Prochrom column packed with Kromosil CHI-DMP chiral phase with an eluent mixture of 3A alcohol and dimethylethylamine in heptane) to give 0.8 g (24%) of isomer 1 and 0.9 g (26%) of isomer 2 as white solids:

Example 23 (Isomer 1). $^{1}H-NMR$ (d, DMSO) 0.75 (d, J = 6.4) Hz, 1.5H), 0.88 (d, J = 6.4 Hz, 1.5H), 1.10 (m, 1H), 1.35 (m, 1H), 1.45-1.70 (m, 8H), 2.60-2.75 (m, 2H), 3.15 (m, 1H)1H), 3.65-3.85 (m, 3H), 4.35 (m, 1H), 4.52 (s, 2H), 4.75(m, 1H), 6.95 (d, J = 11.3 Hz, 1H), 7.20-7.49 (m, 9H), 7.45 (m, 1H), 7.52 (m, 1H), 8.05 (br s, 1H), 8.25 (m,3H), 8.56 (m, 1H), 10.95 (br s, 1H); $t_R = 6.73$ min; MS 579.4 (M+1);Anal. Calc'd for (ion spray) $C_{31}H_{39}FN_6O_4$ 2HCl 0.2CHCl3: C, 56.29; H, 6.24; N, 12.67. Found: C, 56.47; H, 6.17; N, 12.24.

Example 24. (Isomer 2) 1 H-NMR (d, DMSO) 0.75 (d, J = 6.4 Hz, 1.5H), 0.88 (d, J = 6.4 Hz, 1.5H), 1.10 (m, 1H), 1.35 (m, 1H), 1.45-1.70 (m, 8H), 2.60-2.75 (m, 2H), 3.15 (m, 1H), 3.65-3.85 (m, 3H), 4.35 (m, 1H), 4.52 (s, 2H), 4.75 (m, 1H), 6.95 (d, J = 11.3 Hz, 1H), 7.20-7.49 (m, 9H), 7.45 (m, 1H), 7.52 (m, 1H), 8.05 (br s, 1H), 8.25 (m, 3H), 8.56 (m, 1H), 10.95 (br s, 1H); $t_R = 9.09$ min; MS (ion spray) 579.4 (M+1); Anal. Calc'd for $C_{31}H_{39}FN_6O_4$ 2HCl:

176

C, 57.14; H, 6.34; N, 12.90. Found: C, 57.17; H, 6.18; N, 12.79.

Preparation 53

Reaction of the product of Preparation 51 (0.6 g, 1.0 pyrrolidine (0.08 mL, 1.0 mmol), 1hydroxybenzotriazole (0.15 g, 1.1 mmol) and 1 - (3 dimethylaminopropyl)-3-ethylcarbodiimide (0.23 mmol) in dimethylformamide (20 mL)as described in Preparation 6 gave 0.27 g (41%) of the desired product as a white foam: 1H-NMR is consistent with structure; MS (FD) 650.5 (M+); Anal. Calc'd for C₃₄H₄₃FN₆O₆ 0.6H₂0: C, 61.73; H, 6.73; N, 12.70. Found: C, 61.98; H, 6.43; N, 12.66.

Example 25

Reaction of the product of Preparation 53(0.2 g, 0.3 mmol) and trifluoroacetic acid (4 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.16 g (84%) of the desired mixture of isomers as a yellow solid: ¹H-NMR is consistent with structure. MS (high res) calc'd for $C_{29}H_{36}FN_6O_4$: 551.2782. Found: 551.2790.

177

Reaction of the product of Preparation 51 (1.0 g, 1.7 mmol), dimethylamine hydrochloride (0.14 g, 1.7 mmol), triethylamine (0.26 mL, 1.9 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol) in dimethylformamide (30 mL) as described in Preparation 6 gave 0.55 g (52%) of the desired product as a tan foam: $^1\text{H-NMR}$ is consistent with structure; MS (ion spray) 625.4 (M+1); Anal. Calc'd for $C_{32}\text{H}_{41}\text{FN}_6\text{O}_6$: C, 61.53; H, 6.61; N, 13.45. Found: C, 61.22; H, 6.33; N, 13.44.

Example 26

Reaction of the product of Preparation 54 (0.54 g, 0.86 mmol) and trifluoroacetic acid (2 mL), dichloromethane (6 mL) as described in Example 1 gave 0.4 g (77%) of the desired product as a mixture of isomers: $^{1}\text{H-NMR}$ is consistent with structure. MS (ion spray) 525.4 (M+1); Anal. Calc'd for $C_{27}H_{33}FN_6O_6$ 2HCl: C, 54.27; H, 5.90; N, 14.06. Found: C, 53.11; H, 5.70; N, 13.58.

178

Reaction of the product of Preparation 51 (0.6 g, 1.0 mmol), 4-(4-fluorobenzoyl)piperidine hydrochloride (0.25 g, 1.0 mmol), triethylamine (0.16 mL, 1.1 mmol), hydroxybenzotriazole (0.15 g, 1.1 mmol) dimethylaminopropyl)-3-ethylcarbodiimide (0.23)g, mmol) in dimethylformamide (40 mL) as Preparation 6 gave 0.42 g (53%) of the desired product as a tan foam: 1H-NMR is consistent with structure; MS (ion spray) 787.4 (M+); Anal. Calc'd for C₄₂H₄₈F₂N₆O₇: C, 63.83; H, 6.17; N, 10.63. Found: C, 63.95; H, 5.90; N, 10.44.

Example 27

Reaction of the product of Preparation 55(0.4 g, 0.5 mmol) with trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.32 g (82%) of the desired product as a yellow foam: $^1\text{H-NMR}$ is consistent with structure. MS (high res) calc'd for $C_{37}H_{41}F_2N_6O_5$: 687.3106. Found: 687.3103. Anal. Calc'd for $C_{37}H_{40}F_2N_6O_5$: 2.4 HCl: C, 57.40; H, 5.52; N, 10.85. Found: C, 57.56; H, 5.53; N, 10.50.

179

Reduction of the product of Preparation 4 (4.8 g, 16.0 mmol) with 10% palladium on carbon (5.0 tetrahydrofuran (160 mL) followed by coupling with the product of Preparation 1j (6.0 g, 16.0 mmol), hydroxybenzotriazole (2.4 g, 17.6 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (3.6 g, 17.6 mmol) as described in Preparation 4 gave 15.4 g (77%) of the desired product as a tan foam: 1H-NMR (d, DMSO) 1.17 (t, J = 7.2 Hz, 3H), 1.23-1.45 (m, 15H), 1.45-1.57 (m,6H), 7.16 (q, J = 6.8 Hz, 2H), 4.40 (m, 1H), 6.45 (s, 1H), 7.05 (m, 1H), 7.10-7.30 (m, 8H), 7.40-7.48 (m, 3H), 7.54 (s, 1H), 10.20 (br s, 1H); MS (ion spray) 624.4 (M+1); Anal. Calc'd for $C_{33}H_{42}FN_5O_6$: C, 63.55; H, 6.79; N, 11.23. Found: C, 63.83; H, 6.78; N, 11.38.

180

Preparation 57

Reaction of the product of Preparation 56 (14.8 g, 24.0 mmol) with lithium hydroxide (0.66 g, 29.0 mmol) in dioxane (200 mL) and water (100 mL) as in described in Preparation 5 gave 14.3 g (100%) of the desired product as a tan foam: $^{1}\text{H-NMR}$ (d, DMSO) 1.25-1.40 (m, 15H), 1.50-1.75 (m, 6H), 4.40 (s, 1H), 6.60 (s, 1H), 7.05 (s, 1H), 7.10-7.30 (m, 8H), 7.40-7.50 (m, 3H), 7.55 (s, 1H), 10.2 (br s, 1H), 13.63 (br s, 1H); MS (ion spray) 596.5 (M+1); Anal. Calc'd for $C_{31}H_{30}FN_{5}O_{6}$ 0.1dioxane: C, 62.39; H, 6.47; N, 11.59. Found: C, 62.16; H, 6.56; N, 11.28.

Preparation 58

Reaction of the product of Preparation 57 (13.3 g, 23.1 mmol), 4-methylpiperidine (3 mL, 23.1 mmol), 1-hydroxybenzotriazole (3.4 g, 25.4 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (5.2 g, 25.4 mmol) in dimethylformamide (100 mL) as described in Preparation 6 gave 14.4 g (93%) of the desired product as a tan foam: 1 H-NMR (d, DMSO) 0.76 (d, J = 6.4 Hz, 1.5 H), 0.86 (d, J = 4.9 Hz, 1.5H), 1.00 (m, 1H), 1.25-1.45 (m, 17H), 1.45-1.75 (m, 8H), 2.60-2.80 (m, 2H), 3.75 (m, 1H), 4.30-4.45 (m, 2H), 6.71 (d J = 11.7 Hz, 1H), 7.05 (m,

181

1H), 7.10-7.30 (m, 9H), 7.30-7.45 (m, 3H), 10.15 (m, 1H); MS (ion spray) 677.5 (M+1); Anal. Calc'd for $C_{37}H_{49}FN_6O_5$: C, 65.66; H, 7.30; N, 12.42. Found: C, 65.78; H, 7.19; N, 12.44.

Examples 28 and 29

Reaction of the product of Preparation 58(13.8 g, 20.4 mmol) with trifluoroacetic acid (16 mL) in dichloromethane (40 mL) as described in Example 1 gave 10.5 g (89%) of the desired mixture as a tan foam. The mixture (4.0 g) was purified by HPLC (8 x 15 cm Prochrom column packed with Kromosil CHI-DMP chiral phase with an eluent mixture of 3A alcohol and dimethylethylamine in heptane) to give 1.5 g (38 %) of isomer 1 and 0.77 g (20%) of isomer 2 as white solids:


Example 28. (isomer 1) 1 H-NMR (d, DMSO) 0.75 (t, J = 6.4 Hz, 1.5 H), 0.87 (t, J = 6.0 Hz, 1.5 H), 1.15 (m, 1H), 1.35 (m, 1H), 1.45-1.80 (m, 12H), 2.55-2.75 (m, 3H), 3.05 (m, 1H), 3.65-3.75 (m, 2H), 4.30-4.50 (m, 2H), 6.94 (d, J = 12 Hz, 1H), 7.10-7.20 (m, 2H), 7.20-7.40 (m, 7H), 7.45 (m, 1H), 7.55 (m, 1H), 8.08 (m, 1H), 8.15-8.30 (m, 3H), 8.44 (t, J = 7.2 Hz, 1H), 10.90 (br s, 1H); $t_R = 6.62$ min; MS (ion spray) 578.3 (M+1); Anal. Calc'd for $C_{32}H_{41}FN_{6}O_{3}$ 2.3HCl: C, 58.81; H, 6.61; N, 12.72. Found: C, 57.91; H, 6.55; N, 12.72.

Example 29. (isomer 2) 1 H-NMR (d, DMSO) 0.75 (t, J = 6.4 Hz, 1.5 H), 0.87 (t, J = 6.0 Hz, 1.5 H), 1.15 (m, 1H), 1.35 (m, 1H), 1.45-1.80 (m, 12H), 2.55-2.75 (m, 3H), 3.05 (m, 1H), 3.65-3.75 (m, 2H), 4.30-4.50 (m, 2H), 6.94 (d, J

182


= 12 Hz, 1H), 7.10-7.20 (m, 2H), 7.20-7.40 (m, 7H), 7.45 (m, 1H), 7.55 (m, 1H), 8.08 (m, 1H), 8.15-8.30 (m, 3H), 8.44 (t, J = 7.2 Hz, 1H), 10.90 (br s, 1H); $t_R = 8.95$ min; MS (ion spray) 578.3 (M+1); Anal. Calc'd for $C_{32}H_{41}FN_6O_3$ 2.3HCl: C, 58.81; H, 6.61; N, 12.72. Found: C, 58.05; H, 6.64; N, 12.43.

Preparation 59



Reaction of 3,4-difluorophenylacetic acid (25.0 g, 145 mmol) with p-toluenesulfonic acid (9.5 g, 49.5 mmol) in absolute ethanol (150 mL) as described in Preparation 1 gave 28.7 g (99%) of the desired product as a colorless oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 201 (M+).


Preparation 60



Reaction of the product of Preparation 59 (10.0 g, 50.0 mmol, N-bromosuccinimde (9.17 g, 51.5 mmol) and 48% HBr (4 drops) in carbon tetrachloride (40 mL) as in Preparation 2 gave 12.0 g (86%) of the desired product as a colorless oil which was used without further purification: ¹H-NMR is consistent with structure; MS (ion spray) 278, 280 (M+1).

183

Preparation 61



Reaction of the product of Preparation 60 (10.5 g, 38 mmol), 4-nitroimidazole (5.2 g, 45.6 mmol) and potassium carbonate (15.1 g, 114 mmol) in dimethylformamide (400 mL) as described in Preparation 3 gave 4.54 g (39%) of the desired product as an orange oil: $^1\text{H-NMR}$ is consistent with structure; MS (ion spray) 312.0 (M+1); Anal. Calc'd for $C_{13}H_{11}F_2N_3O_4\cdot 0.2H_2O$: C, 49.59; H, 3.65; N, 13.35. Found: C, 49.58; H, 3.62; N, 13.09.

Preparation 62

Reduction of the product of Preparation 61 (1.35 g, 4.3 mmol) with 10% palladium on carbon (0.8 g) in tetrahydrofuran (40 mL) followed by coupling with the product of Preparation 1d (1.64 g, 4.3 mmol), 1-hydroxybenzotriazole (0.7 g, 4.7 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.04 g, 4.7 mmol) as described in Preparation 4 gave 1.9 g (69%) of the desired product as a tan foam: $^1\text{H-NMR}$ is consistent with structure; MS (ion spray) 644 (M+1); Anal. Calc'd for $C_{32}H_{39}F_2N_5O_7$: C, 59.71; H, 6.11; N, 10.80. Found: C, 59.72; H, 6.04; N, 10.63.

Preparation 63

Reaction of the product of preparation 62 (1.9 g, 3:0 mmol) with lithium hydroxide (0.09 g, 3.6 mmol) in dioxane (50 mL) and water (25 mL) as described in Preparation 5 gave 1.6 g (87%) of the desired product as a tan foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 616.4 (M+1).

Preparation 64

Reaction of the product of Preparation 63 (0.5 g, 0.8 mmol), 4-methylpiperidine (0.1 mL, 0.8 mmol), 1-hydroxybenzotriazole (0.12 g, 0.88 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.18 g, 0.88 mmol) in dimethylformamide (40 mL) as described in Preparation 6 gave 0.3 g (54%) of the desired product as a white foam: $^1\text{H-NMR}$ is consistent with structure; MS (ion spray) 697 (M+1); Anal. Calc'd for $C_{36}H_{46}F_2N_6O_6$; C, 62.06; H, 6.65; N, 12.06. Found: C, 61.82; H, 6.57; N, 11.96.

185

Example 30

Reaction of the product of Preparation 64 (0.22 g, 0.3 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.2 g (100%) of the desired mixture of isomers as a yellow foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 597.5 (M+1); Anal. Calc'd for $C_{31}H_{38}F_{2}N_{6}O_{4}$ 2.2HCl; C, 55.01; H, 5.99; N, 12.42. Found: C, 55.16; H, 5.96; N, 12.20.

Preparation 65

Reaction of the product of Preparation 63 (0.5 g, 0.8 (0.07 mL, 0.8 pyrrolidine mmol), 1mmol), hydroxybenzotriazole (0.12 mL, 0.88 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.18 g, mmol) in dimethylformamide (40 mL) as described in Preparation 6 gave 0.25 g (42%) of the desired product as a tan foam: 1H-NMR is consistent with structure. MS (ion spray) 669.4 (M+1); Anal. Calc'd for $C_{34}H_{42}F_2N_6O_6 \cdot 0.7H_20$: C, 59.94; H, 6.42; N, 12.33. Found: C, 59.96; H, 6.28; N, 11.97.

Example 31

186

Reaction of the product of Preparation 65 (0.2 g, 0.3 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.14 g (74%) of the desired product as a yellow foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 569.4 (M+1); Anal. Calc'd for $C_{29}H_{54}F_{2}N_{6}O_{4}$ 2.2HCl: C, 53.68; H, 5.62; N, 12.95. Found: C, 53.83; H, 5.57; N, 12.37.

Preparation 66

Reaction of the product of preparation 63 (0.5 g, 0.8 mmol), 4-(4-fluorobenzoyl)piperidine hydrochloride (0.2 g, 0.8 mmol), triethylamine (0.13 mL, 0.88 mmol), 1-hydroxybenzotriazole (0.12 g, 0.88 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.18 g, 0.88 mmol) in dimethylformamide (40 mL) as described in Preparation 5 gave 0.14 g (22%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (ion spray) 805.6 (M+1); Anal. Calc'd for C₄₂H₄₇F₃N₆O₇: C, 62.68; H, 5.89; N, 10.44. Found: C, 62.45; H, 5.82; N, 10.40.

Example 32

187

Reaction of the product of Preparation 66 (0.14 g, 0.17 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.1 g (77%) of the desired mixture of isomers as a yellow solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 705.5 (M+1); Anal. Calc'd for $C_{37}H_{39}F_{3}N_{6}O_{5}$ 2.1HCl: C, 56.88; H, 5.30; N, 10.76. Found: C, 56.64; H, 5.31; N, 10.30.

Preparation 67



Reaction of 2,4-difluorophenylacetic acid (20 g, 116 mmol) and p-toluenesulfonic acid (6.0 g, 31 mmol) in absolute ethanol (150 mL) as described in Preparation 1 gave 22.1 g (95%) of the desired product as a colorless oil which solidifies upon setting: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 200 (M+).

Preparation 68

Reaction of the product of Preparation 67 (21.4 g, 100 mmol), N-bromosuccinimide (19.6 g, 103 mmol) and 48% HBr (6 drops) in carbon tetrachloride (100 mL) as described in Preparation 2 gave 27.9 g (100%) of the desired product as a colorless oil: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 278, 280 (M+); Anal. Calc'd for $C_{10}H_9\text{Br}F_2O_2$: C, 43.04; H, 3.25. Found: C, 42.92; H, 3.15.

Preparation 69

Reaction of the product of Preparation 68 (26.9 g, 96 mmol), 4-nitroimidazole (13.0 g, 115 mmol) and potassium carbonate (40 g, 288 mmol) in dimethylformamide (150 mL) as described in Preparation 3 gave 14.3 g (48%) of the desired product as an orange oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 312 (M+1); Anal. Calc'd for $C_{13}H_{11}F_{2}N_{3}O_{4}$: C, 50.17; H, 3.56; N, 13.50. Found: C, 49.90; H, 3.56; N, 13.26.

Preparation 70

Reduction of the product of preparation 69(1.35 g, 4.3 mmol) with 10% palladium on carbon (0.8 g) in tetrahydrofuran (40 mL) followed by coupling with the product of Preparation 1d (1.64 g, 4.3 mmol), hydroxybenzotriazole (0.7 g, 4.7 mmol) and dimethylaminopropyl)-3-ethylcarbodiimide (1.04 mmol) as desribed in Preparation 4 gave 1.52 g (55%) of the desired product as a tan foam: 1H-NMR is consistent with structure; MS (ion spray) 644.5 (M+1); Anal. Calc'd for $C_{32}H_{39}F_2N_5O_7$: C, 59.71; H, 6.11; N, 10.88. Found: C, 59.43; H, 5.97; N, 10.91.

189

Preparation 71

Reaction of the product of preparation 70 (1.42 g, 2.2 mmol) with lithium hydroxide (0.07 g, 2.64 mmol) in dioxane (50 mL) and water (25 mL) as described in Preparation 5 gave 1.35 g (100%) of the desired product as a tan foam: 1H -NMR is consistent with structure; MS (ion spray) 616.3 (M+1); Anal. Calc'd for $C_{30}H_{35}F_2N_5O_7$; C, 58.33; H, 5.73; N, 11.38. Found: C, 57.71; H, 5.86; N, 10.80.

Preparation 72

Reaction of the product of Preparation 71 (0.6 g, 1.0 mmol), 4-methylpiperidine (0.12 mL, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol) in dimethylformamide (30 mL) as described in Preparation 6 gave 0.66 g (94%) of the desired product as a tan foam: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 696 (M+); Anal. Calc'd for $C_{36}H_{46}F_2N_6O_6$: C, 62.05; H, 6.65; N, 12.06. Found: C, 62.21; H, 6.48; N, 12.17.

Example 33

190

Reaction of the product of Preparation 72 (0.51 g, 0.73 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as in described in Example 1 gave 0.25 g (51%) of the desired product as tan foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 597.5 (M+1); Anal. Calc'd for $C_{31}H_{38}F_{2}N_{6}O_{4}$ 2.2HCl: C, 55.01; H, 5.99; H, 12.42. Found: C, 56.92; H, 5.98; N, 12.36.

Preparation 73

Reaction of the product of Preparation 71 (0.6 g, 1.0 mmol), pyrrolidine (0.8 mL, 1.0 mmol), hydroxybenzotriazole (0.15 g, 1.1 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, mmol); dimethylformamide (30 mL) as described Preparation 6 gave 0.4 g (58%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (ion spray) 669.5 (M+1); Anal. Calc'd for $C_{32}H_{42}F_2N_6O_6$: C, 61.07; H, 6.33; N, 12.57. Found: C, 60.84; H, 6.31; N, 12.32.

Example 34

191

Reaction of the product of Preparation 73 (0.3 g, 0.45 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.21 g (70%) of the desired product as a white foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 569.4 (M+1); Anal. Calc'd for $C_{29}H_{34}F_{2}N_{6}O_{4}$ 2.3HCl: C, 53.38; H, 5.61; N, 12.88. Found: C, 53.59; H, 5.58; N, 12.42.

Preparation 74



To a solution of the compound of Preparation 49 (17.0 g, 58.0 mmol) stirring at room temperature was added to sodium hydroxide (125 mL of a 2N aqueous solution) along with tetrahydrofuran (10 mL) and ethanol (10 mL). hydrolysis was complete, the mixture was cooled in an bath and acidified to pH 2.75 with aqueous hydrochloric acid and extracted with ethyl acetate. The combined organic extracts were washed with water, dried over sodium sulfate and concentrated to provide 15.0 g (99%) of the desired carboxylic acid. The crude material was combined with aqueous N, N-dimethyl amine (40%, 9.0 mL, 71.8 mmol), 1-hydroxybenzotriazole hydrate (7.64 g, 56.6 mmol) and 1,3-dicyclohexylcarbodiimide (11.7 g, 56.6 mmol) in tetrahydrofuran (150 mL). After 18 h, the mixture was concentrated and the residue slurried in ethyl acetate, filtered, and the filtrate concentrated.

192

Purification of the concentrate by flash chromatography (silica gel, chloroform/methanol) provided 10.2 g (62%) of the desired product: ESMS: $(M+H)^+$ 293.1. 1H NMR (300 MHz, DMSO- d_6) δ 8.21 (d, 1H, J = 1.51 Hz) 7.80(d, 1H, J = 1.13 Hz), 7.60-7.50 (m, 2H,), 7.38-7.25 (m, 2H), 6.88 (s, 1H), 2.92 (s, 3H), 2.86 (s, 3H). Anal. Calcd. for $C_{13}H_{13}N_4O_3$: C, 53.43; H, 4.48; N, 19.17. Found: C, 53.43; H, 4.71; N, 19.07.

Preparation 75

The product of Preparation 74 (2.0 g (6.85 mmol) was combined with 10% palladium/carbon (1.80 g) and palladium/black (0.20 g) in tetrahydrofuran(75 mL)and the mixture shaken under a hydrogen atmosphere (38 mm Hg) in a Parr apparatus. After reduction was complete, the catalyst was removed by filtration through celite and the resulting solution was immediately added to a solution of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide 7.3 mmol), 1-hydroxybenzotriazole (1.0 g, 7.3 mmol), the product of Preparation 1j (2.77 g, 7.3 mmol) in tetrahydrofuran (50 mL) at room temperature. After 16 h, the mixture was concentrated and the residue slurried in acetate then filtered. The filtrate concentrated and resulting crude product purified by flash chromatography (silica gel, chloroform/methanol) which afforded 3.47 g (81%) of the desired product: ESMS: $(M+H)^+$ 623.5, 624.6. ¹H NMR was consistent with product. Anal. Calcd. for C₃₃H₄₃N₆O₄F.0.02 CHCl3:

193

63.44; H, 6.94; N, 13.44. Found: C, 63.04; H, 7.41; N, 11.93.

Example 35

To a solution of the product of Preparation 75 (1.45 g, mmol) stirring at room temperature dichloromethane (50 mL) was added triflouroacetic acid (15 mL). After 3 hours, the mixture was concentrated and the material treated with excess aqueous sodium The aqueous mixture was extracted with bicarbonate. ethyl acetate and the combined organic concentrate. The resulting residue was purified by flash chromatography (silica gel, chloroform/methanol) provide 1.55 g of the desired product: ESMS: (M+H) 523.3. The isomeric mixture (3.44 g) was separated as previously described in Example 7 to provide 0.98 g of pure isomer 1 ($t_R = 7.94 \text{ min}$) and 0.81 g of isomer 2 ($t_R =$ 10.57 min). For isomer 2, 0.80 g (1.53 mmol) was dissolved in ethyl acetate/methanol and treated with a saturated solution of hydrochloric acid in diethyl ether. The resulting mixture was concentrated to provide 0.90 g (92%) of the desired product as a light tan solid: ESMS: (M+H) 523.4, 524.5. H NMR was consistent with product. Anal. Calcd. for $C_{28}H_{35}N_6O_3F^3.25$ HCl: C, 52.46; H, 6.01; N, 13.11. Found: C, 52.49; H, 6.23; N, 11.80.

Preparation 75

WO 99/08697


The product of Preparation 74 (2.00 g, 6.85 mmol) 10% palladium/carbon combined with (1.80)g) and palladium/black (0.20 g) in tetrahydrofuran 75 mL) and the mixture shaken under hydrogen atmosphere (39 mm Hg) in a Parr apparatus. After reduction was complete, the catalyst was removed by filtration through celite and the amine/tetrahydrofuran solution was immediately combined 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide(1.41 g, 6.85 mmol), 1-hydroxybenzotriazole mono-hydrate (0.93 g, 6.85 mmol), the product of Preparation 1j (2.60 g, 6.84 mmol) and additional tetrahydrofuran (75 mL). After stirring overnight at ambient temperature, the mixture was concentrated and the residue slurried in ethyl acetate. The filtrate was concentrated and the residue chromatography(silica purified by flash chloroform/methanol) to provide 3.65 g (85%)of the 625.4. ¹H desired product as a tan solid: ESMS: (M+H) NMR was consistent with product. Anal. Calcd. for $C_{32}H_{41}N_6O_6$ 0.03 chloroform: C, 61.17; H, 6.60; N, 13.34. Found: C, 61.25; H, 6.90; N, 12.69.

Example 36

WO 99/08697

To a solution of the product of Preparation 75 (3.30 g, 5.3 mmol) stirring in dichloromethane (30 mL) at room temperature was added triflouroacetic acid (10 m). After 3 h, the mixture was concentrated and the residue treated with excess aqueous sodium bicarbonate. The resulting mixture was extracted with ethyl acetate and the combined organic extracts were washed with 1N aqueous hydroxide, dried over sodium sulfate, and concentrated. The residue was purified by flash chromatography (silica gel, chloroform/methanol) to provide 1.40 g (51%) of the desired product as a light tan solid: ESMS: (M+H) + 525.3. ¹H NMR was consistent with product. Anal. Calcd. for $C_{27}H_{33}N_6O_4F$: 1.3 methanol: C, 60.03; H, 6.80; N, 14.84. C, 60.19; H, 6.81; N, 14.56. Found: The isomeric mixture (3.20 g) was separated as previously described in Example 7 to give 1.57 g of isomer 1 ($t_R = 7.57$ min) and 0.88 g of isomer 2 ($t_R = 10.43 \text{ min}$). For isomer 2, 0.88 g (1.68 mmol) was dissolved in ethyl acetate and treated with a saturated solution of hydrochloric acid in diethyl ether. The resulting mixture was concentrated, washed with diethyl ether to give 0.97 g of the desired product: 525.4, 526.7: ¹H NMR was consistent with ESMS: (M+H) + product. Anal. Calcd. for $C_{25}H_{33}N_6O_4F^2$ 2.75 HCl: C, 51.73; H, 6.07; N, 13.41. Found: C, 51.62; H, 5.74; N, 13.34.

Preparation 76



Reaction of 4-ethoxyphenylacetic acid (23.5 g, 130 mmol) and p-toluenesulfonic acid (4.0 g, 21 mmol) in absolute ethanol (150 mL) as described in Preparation 1 gave 23.2 g (86%) of the desired product as a colorless oil: 1 H-NMR (d, DMSO) 1.17 (t, J = 7.2 Hz, 3H), 1.31 (t, J = 7.2 Hz, 3H), 3.56 (s, 2H), 3.99 (q, J = 7.2 Hz, 2H), 4.05 (q, J =


196

7.2 Hz, 2H), 6.85 (d, J = 8.7 Hz, 2H), 7.14 (d, J = 8.7 Hz, 2H); MS (ion spray) 209 (M+1); Anal. Calc'd for $C_{12}H_{16}O_3$: C, 69.21; H, 7.74. Found: C, 68.91; H, 7.55.

Preparation 77

To a solution of the product of Preparation 76 (53 g, 255 mmol) stirring in carbon tetrachloride (600 mL) at room temperature was added 46.6 g (262 mmol) of N-bromosuccinimide and 3.0 g (18.3 mmol) of 2,2'-azobis(2-methylpropionitrile). The resulting reaction mixture was heated to reflux. After 3.5 h, the solution was cooled to room temperarure, filtered and concentrated. The resulting oil was chromatographed on silica gel using chloroform as eluant to afford 70.9 g (97%) of the desired product as a colorless oil: 1 H-NMR (d, DMSO) 1.17 (t, J = 7.2 Hz, 3H), 1.25-1.35 (m, 3H), 4.00-4.10 (m, 2H), 4.13-4.25 (m, 2H), 5.86 (s, 1H), 6.92 (d, J = 8.7 Hz, 2H), 7.47 (d, J = 9.0 Hz, 2H); MS (FD) 287, 289 (M+).

Preparation 78



Reaction of the product of Preparation 77 (11.4 g, 40 mmol), 4-nitroimidazole (4.5 g, 40 mmol) and potassium carbonate (16.6 g, 120 mmol) in dimethylformamide (100 mL) as described in Preparation 3 gave 5.47 g (43%) of the desired product as a yellow oil: 1 H-NMR (d, DMSO) 1.18 (t, J = 7.2 Hz, 3H), 1.29 (t, J = 7.2 Hz, 3H), 4.03 (q, J

197

= 7.2 Hz, 2H), 4.23 (q, J = 7.2 Hz, 2H), 6.54 (s, 1H), 6.70 (d, J = 8.7 Hz, 2H), 7.42 (d, J = 8.7 Hz, 2H), 7.90 (s, 1H), 8.34 (s, 1H); MS (ion spray) 320.2 (M+1); Anal. Calc'd for $C_{15}H_{17}N_3O_5$: C, 56.42; H, 5.37; N, 13.16. Found: C, 56.29; H, 5.17; N, 13.15.

Preparation 79

Reduction of the product of Preparation 78 (9.6 g, 30 with 10% palladium on carbon (7.0 g) tetrahydrofuran (100 mL) followed by coupling with the product of Preparation 1d (11.5 g, 30 mmol), hydroxybenzotriazole (4.5 a, 33 mmol) and dimethylaminopropy1)-3-ethylcarbodiimide (6.8 g, 33 mmol) as described in Preparation 4 gave 9.9 g (50%) of the desired product as a tan foam: 1H-NMR (d, DMSO) 1.17 (t, J = 7.2 Hz, 3H, 1.25-1.40 (m, 18H), 3.58 (m, 1H), 3.70 (m,1H), 4.02 (q, J = 7.2 Hz, 2H), 4.20 (q, J = 7.2 Hz, 2H), 4.44 (d, J = 3.4 Hz, 2H), 4.60 (m, 1H), 6.33 (s, 1H), 6.95 (d, J = 8.7 Hz, 2H), 7.15-7.35 (m, 9H), 7.43 (m, 1H), 7.51 (m, 1H), 10.2 (br s, 1H); MS (ion spray) 652.4 (M+1); Anal. Calc'd for $C_{34}H_{45}N_5O_8$: C, 62.66; H, 6.96; N, 10.74. Found: C, 62.92; H, 7.00; N, 10.98.


Preparation 80

198

Reaction of the product of Preparation 80 (9.7 g, 15.0 mmol) and lithium hydroxide (0.42 g, 18.0 mmol) in dioxane (200 mL) and water (100 mL) as described in Preparation 5 gave 9.4 g (100%) of the desired product as a tan foam: 1 H-NMR (d, DMSO) 1.25-1.40 (m, 18H), 3.60 (m, 1H), 3.68 (m, 1H), 4.02 (q, J = 7.2 Hz, 2H), 4.44 (d, J = 3.0 Hz, 2H), 4.60 (m, 1H), 6.19 (m, 1H), 6.95 (d, J = 8.7 Hz, 2H), 7.28-7.35 (m, 9H), 7.40 (m, 1H), 7.51 (s, 1H), 10.2 (br s, 1H), 13.5 (br s, 1H); MS (ion spray) 624.5 (M+1); Anal. Calc'd for $C_{43}H_{41}N_{5}O_{8}$: C, 61.62; H, 6.63; N, 11.23. Found: C, 61.58; H, 6.92; N, 10.99.

199

Preparation 81



Reaction of the product of Preparation 80 (7.43 g, 12.0 mmol), 4-methylpiperidine (1.42 mL, 12.0 mmol), hydroxybenzotriazole (1.78 g, 13.2 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (2.72 13.2 g, mmol) in dimethylformamide (100 mL) as described in Preparation 6 gave 6.4 g (76%) of the desired product as a tan foam: $^{1}H-NMR$ (d, DMSO) 0.74 (d, J = 6.4 Hz, 1.5 H), 0.87 (d, J = 6.0 Hz, 1.5H), 1.05 (m, 1H), 1.25-1.40 (m, 18H), 1.50-1.70 (m, 3H), 2.55-2.70 (m, 2H), 3.00 (m, 1H), 3.57 (m, 1H), 3.65-3.85 (m, 2H), 4.00-4.20 (m, 2H), 4.38 (m, 1H), 4.44 (d, J = 3.4 Hz, 2H), 4.60 (m, 1H), 6.61 (d,J = 12.0 Hz, 1H), 6.95-7.00 (m, 2H), 7.15-7.20 (m, 2H),7.20-7.45 (m, 9H), 10.15 (br s, 1H); MS (ion spray) 705.5 (M+1); Anal. Calc'd for $C_{38}H_{52}N_6O_7$: C, 64.75; H, 7.44; N, 11.92. Found: C, 64.59; H, 7.21; N, 11.87.

Example 37 and 38

Reaction of the product of Preparation 81 (6.4, 9.1 mmol) and trifluoroacetic acid (10 mL) in dichloromethane (25 mL) as described in Example 1 gave 4.71 g (77%) of the desired mixture of diastereomers as a tan foam. Resolution of the diastereomers (2.4 g) by HPLC (Kromsil

200

CHI-DMP chiral stationary phase, 3A alcohol/dimethylethylamine/heptane) provided 200 mg (8 %) of isomer 1 and 0.8 g (31 %) of isomer 2, both isolated as white solids after acidification with hydrochloric acid as described in Example 7:

Example 37. (Isomer 1) 1 H-NMR (d, DMSO) 0.74 (d, J = 6.4 Hz, 1.5H), 0.88 (d, J = 6.0 Hz, 1.5H), 1.20 (m, 1H), 1.31 (t, J = 6.8 Hz, 3H), 1.45-1.70 (m, 8H), 2.60-2.70 (m, 2H), 3.05 (m, 1H), 3.65-3.80 (m, 3H), 4.00-4.20 (m, 3H), 4.37 (m, 1H), 4.52 (s, 2H), 4.75 (m, 1H), 6.80 (d, J = 13.2 Hz, 1H), 6.95-7.05 (m, 2H), 7.25-7.40 (m, 9H), 7.92 (br s, 1H), 8.20-8.30 (m, 3H), 8.53 (d, J = 7.2 Hz, 1H), 10.9 (br s, 1H); $t_R = 9.17$ min; MS (ion spray) 605 (M+1); Anal. Calc'd for $C_{33}H_{44}N_6O_5$ 2HCl 0.1 CHCl3: C, 58.45; H, 6.74; N, 12.74. Found: C, 58.64; H, 6.77; N, 12.36.

Example 38. (Isomer 2) 1 H-NMR (d, DMSO) 0.74 (d, J = 6.4 Hz, 1.5H), 0.88 (d, J = 6.0 Hz, 1.5H), 1.20 (m, 1H), 1.31 (t, J = 6.8 Hz, 3H), 1.45-1.70 (m, 8H), 2.60-2.70 (m, 2H), 3.05 (m, 1H), 3.65-3.80 (m, 3H), 4.00-4.20 (m, 3H), 4.37 (m, 1H), 4.52 (s, 2H), 4.75 (m, 1H), 6.80 (d, J = 13.2 Hz, 1H), 6.95-7.05 (m, 2H), 7.25-7.40 (m, 9H), 7.92 (br s, 1H), 8.20-8.30 (m, 3H), 8.53 (d, J = 7.2 Hz, 1H), 10.9 (br s, 1H); $t_R = 12.68 \text{ min}$; MS (ion spray) 605 (M+1); Anal. Calc'd for $C_{33}H_{44}N_6O_5$ HCl: C, 59.35; H, 6.85; N, 12.98. Found: C, 59.62; H, 7.01; N, 12.71.


201

Preparation 82

Reaction of the product of Preparation 80 (0.9 g, 1.5 mmol), dimethylamine hydrochloride (0.13 g, 1.5 mmol), mL, 1.65 mmol), (0.23 triethylamine hydroxybenzotriazole (0.23 g, 1.65 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.34 g, 1.65 mmol) in dimethylformamide (50 mL) as described in Preparation 6 gave 0.46 g (47%) of the desired product as a tan foam: $^{1}H-NMR$ (d, DMSO) 1.25-1.35 (m, 18H), 2.90 (m, 6H), 3.57 (m, 1H), 3.67 (m, 1H), 4.03 (q, J = 7.2 Hz, 2H), 4.43-4.47 (m, 2H), 4.57 (m, 1H), 6.55 (m, 1H), 6.97 (d, J = 8.7 Hz, 2H), 7.15-7.45 (m, 11H), 10.16 (br s,1H); MS (ion spray) 651.4 (M+1); Anal. Calc'd for $C_{34}H_{46}N_6O_7$: C, 62.75; H, 7.13; N, 12.91. Found: C, 62.55; H, 6.84; N, 12.84.

202

Examples 39 and 40



Reaction of the product of Preparation 82 (0.44 g, 0.68 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.19 g (45%) of the desired product as a tan foam. Resolution of the diastereomers (90 mg, 0.14 mmol) by HPLC (Kromsil CHI-DMP chiral stationary phase, 3A alcohol/dimethylethylamine /heptane) provided 50 mg (50 %) of isomer 1 and 27 mg (27%) of isomer 2, both isolated as white solids after acidification with hydrochloric acid as described in Example 7:

Example 39. (isomer 1): 1 H-NMR (d, DMSO) 1.32 (t, J = 6.8 Hz, 3H), 1.50 (s, 6H), 2.86 (s, 3H), 2.90 (s, 3H), 3.70-3.80 (m, 2H), 4.03 (q, J = 7.2 Hz, 2H), 4.52 (s, 2H), 4.75 (m, 1H), 6.76 (s, 1H), 7.00 (d, J = 8.7 Hz, 2H), 7.25-7.40 (m, 9H), 8.06 (m, 1H), 8.20-8.30 (m, 3H), 8.52-8.60 (m, 1H), 11.00 (br s, 1H); $t_R = 7.70$ min; MS (high res) calc'd for $C_{29}H_{39}N_6O_5$: 551.2982. Found: 551.2987. Anal. Calc'd for $C_{29}H_{38}N_6O_5$: 2.3 HCl'0.3ethyl acetate: C, 54.88; H, 6.51; N, 12.72. Found: C, 54.70; H, 6.49; N, 12.43.

Example 40. (isomer 2): 1 H-NMR (d, DMSO) 1.32 (t, J = 6.8 Hz, 3H), 1.50 (s, 6H), 2.86 (s, 3H), 2.90 (s, 3H), 3.70-3.80 (m, 2H), 4.03 (q, J = 7.2 Hz, 2H), 4.52 (s, 2H), 4.75 (m, 1H), 6.76 (s, 1H), 7.00 (d, J = 8.7 Hz, 2H), 7.25-7.40 (m, 9H), 8.06 (m, 1H), 8.20-8.30 (m, 3H), 8.52-8.60 (m, 1H), 11.00 (br s, 1H); t_{R} = 9.09 min; MS (high res) calc'd for $C_{29}H_{39}N_{6}O_{5}$: 551.2982. Found: 551.2976. Anal. Calc'd for $C_{29}H_{38}N_{6}O_{5}$: 2.HCl'0.3 ethyl acetate: C,

203

55.18; H, 6.53; N, 12.79. Found: C, 55.01; H, 6.33; N, 12.54.

Preparation 83

Reaction of the product of Preparation 80 (0.9 g, 1.5 mmol), pyrrolidine (0.13 mL, 1.5 mmol), hydroxybenzotriazole (0.23 g, 1.65 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.34 g, 1.65 mmol) in dimethylformamide (40 mL) as described in Preparation 6 gave 0.7 g (74%) of the desired product as a tan foam: ¹H-NMR (d, DMSO) 1.25-1.40 (m, 18H), 1.70-1.90 (m, 4H), 2.95 (m, 1H), 3.30-3.40 (m, 2H), 3.55-3.70 (m, 2H)3H), 4.03 (q, J = 7.2 Hz, 2H), 4.44 (d, J = 3.4 Hz, 2H), 4.57 (m, 1H), 6.34 (s, 1H), 6.97 (d, J = 8.7 Hz, 2H), 7.20-7.35 (m, 9H), 7.40-7.45 (m, 2H), 10.15 (br s, 1H); (ion spray) 677.6 (M+1); Anal. Calc'd for $C_{36}H_{48}N_6O_7$ 0.2 H_2 0: C, 63.55; H, 7.17; N, 12.35. Found: C, 63.32; H, 6.96; N, 12.24.


Example 41

Reaction of the product of Preparation 83 (0.59 g, 0.9 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.36 g (64%) of

204

the desired product as a mixture of isomers: $^{1}H-NMR$ (d, DMSO) 1.32 (t, J = 6.8 Hz, 3H), 1.45-1.60 (m, 6H), 1.65-1.90 (m, 4H), 2.90 (m, 1H), 3.25-3.45 (m, 2H), 3.65-3.75 (m, 3H), 4.02 (q, J = 6.8 Hz, 2H), 4.45-4.55 (m, 2H), 4.70-4.80 (m, 1H), 6.54 (s, 1H), 6.98 (d, J = 8.7 Hz, 2H), 7.20-7.40 (m, 9H), 8.05 (m, 1H), 8.20-8.30 (m, 3H), 8.54 (d, J = 7.2 Hz, 1H), 10.95 (br s, 1H); MS (high res) calc'd for $C_{31}H_{41}N_6O_5$: 577.3138. Found: 577.3132. Anal. Calc'd for $C_{31}H_{40}N_6O_5$: 2HCl: C, 57.32; H, 6.52; N, 12.94. Found: C, 57.46; H, 6.59; N, 12.91.


Preparation 84



Reaction of 4-butyloxyphenylacetic acid (10.0 g, 48 mmol) and p-toluenesulfonic acid (2.5 g, 13 mmol) in absolute ethanol (100 mL) as described in Preparation 1 gave 11.04 g (98%) of the desired product as a colorless oil: $^{1}\text{H-NMR}$ (d, DMSO) 0.94 (t, J = 7.4 Hz, 3H), 1.18 (t, J = 7.0 Hz, 3H), 1.40-1.50 (m, 2H), 1.60-1.80 (m, 2H), 3.57 (s, 2H), 3.93 (q, J = 6.5 Hz, 2H), 4.08 (q, J = 7.3 Hz, 2H), 6.86 (d, J = 8.4 Hz, 2H), 7.15 (d, J = 8.4 Hz, 2H); MS (ion spray) 237 (M+1); Anal. Calc'd for $C_{14}H_{20}O_{3}$: C, 71.16; H, 8.53. Found: C, 71.33; H, 8.55.


205

Preparation 85



To a solution of the product of Preparation 84, 6.0 g (25 mmol) in 100 mL of carbon tetrachloride was added 4.7 g (25.8 mmol) of N-bromosuccinimide and 0.6 g of 2,2'-azobis(2-methylpropionitrile). The reaction mixture was heated to reflux. After 3.5 h, the mixture was cooled to room temperature, filtered and concentrated. The resulting oil was purified by flash chromatography (silica gel, 3% methanol/chloroform) to proved 6.9 g (88%) of the desired product as a colorless oil: ¹H-NMR (d, DMSO) 0.93 (t, J = 7.35 H, 3H), 1.20 (t, J = 7.2 Hz, 3H), 1.40-1.50 (m, 2H), 1.60-1.80 (m, 2H), 3.95-4.05 (m, 2H), 4.10-4.15 (m, 2H), 5.87 (s, 1H), 6.93 (d, J = 8.7 Hz, 2H), 7.45 (d, J = 8.7 Hz, 2H); MS (FD) 314, 316 (M+); Anal. Calc'd for C14H19BrO3 0.5CHCl3: C, 52.54; H, 5.98. Found: C, 52.35; H, 5.84.

Preparation 86



Reaction of the product of Preparation 85(5.82 g, 19.0 mmol), 4-nitroimidazole (2.1 g, 19.0 mmol) and potassium carbonate (8.0 g, 57 mmol) in dimethylformamide (150 mL) as described in Preparation 3 gave 3.5 g (53%) of the desired product as a yellow oil: $^{1}\text{H-NMR}$ (d, DMSO) 0.93 (t, J = 7.3 Hz, 3H), 1.19 (t, J = 7.0 Hz, 3H), 1.35-1.50 (m, 2H), 1.60-1.80 (m, 2H), 3.92-4.06 (m, 2H), 4.20-4.30

206


(m, 2H), 6.56 (s, 1H), 6.99 (d, J = 8.6 Hz, 2H), 7.44 (d, J = 8.6 Hz, 2H), 7.92 (s, 1H), 8.37 (s, 1H); MS (ion spray) 348.3 (M+1); Anal. Calc'd for $C_{17}H_{21}N_3O_5$: C, 58.78; H, 6.09; N, 12.10. Found: C, 59.08; H, 6.21; N, 12.19.

Preparation 87

Reduction of the product of Preparation 86 (1.5 g, 4.3 10% palladium on carbon (0.8g)tetrahydrofuran (40 mL) followed by coupling with the product of Preparation 1d (1.64 g, 4.3 mmol), hydroxybenzotriazole (0.7 g, 4.7 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (1.04 g, mmol) as described in Preparation 4 gave 1.1 g (38%) of the desired product as a tan foam: ¹H-NMR (d, DMSO) 0.92 (t, J = 7.5 Hz, 3H), 1.18 (t, J = 7.2 Hz, 3H), 1.25-1.40(m, 15H), 1.40-1.50 (m, 2H), 1.60-1.75 (m, 2H), 3.60 (m, 2H)1H), 3.70 (m, 1H), 3.95-4.00 (m, 2H), 4.20-4.25 (m, 2H), 4.45-4.48 (m, 2H), 4.57 (m, 1H), 6.35 (s, 1H), 6.97 (t, J = 9.0 Hz, 2H), 7.15-7.35 (m, 9H), 7.40 (m, 1H), 7.50 (s,1H), 10.20 (br s, 1H); MS (ion spray) 680.5 (M+1); Anal. Calc'd for $C_{36}H_{49}N_5O_8$: C, 63.61; H, 7.27; N, 10.30. Found: C, 63.53; H, 6.99; N, 10.54.

207

Preparation 88



Reaction of the product of Preparation 87 (1.1 g, 1.6 mmol) and lithium hydroxide (0.5 g, 1.92 mmol) in dioxane (50 mL) and water (25 mL) as described in Preparation 5 gave 1.04 g (100%) of the desired product as a tan foam: 1 H-NMR (d, DMSO) 0.95 (t, J = 7.5 Hz, 3H), 1.25-1.35 (m, 15H), 1.35-1.50 (m, 2H), 1.65-1.75 (m, 2H), 3.57 (m, 1H), 3.65 (m, 1H), 3.95 (t, J = 6.4 Hz, 2H), 4.57 (m, 1H), 6.19 (d, J = 1.5 Hz, 2H), 6.20 (s, 1H), 6.96 (d, J = 8.7 Hz, 2H), 7.10-7.35 (m, 9H), 7.40 (m, 1H), 7.50 (s, 1H), 10.20 (br s, 1H), 13.45 (br s, 1H); MS (ion spray) 652.5 (M+1); Anal. Calc'd for $C_{32}H_{45}N_5O_8$: C. 62.66; H, 6.96; N, 10.75. Found: C, 62.45; H, 7.07; N, 10.72.

Preparation 89

Reaction of the product of Preparation 88 (1.0 g, 1.6 mmol), 4-methylpiperidine (0.19 mL, 1.6 mmol), 1-hydroxybenzotriazole (0.24 g, 1.8 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.35 g, 1.8 mmol) in dimethylformamide (60 mL) as described in Preparation 6 gave 0.57 g (48%) of the desired product as a tan foam: 1 H-NMR (d, DMSO) 0.75 (d, J = 6.0 Hz, 1H), 0.85-0.95 (m, 6H), 1.25-1.40 (m, 15H), 1.40-1.75 (m, 7H), 2.55-2.75 (m, 2H), 3.00 (m, 1H), 3.55 (m, 1H), 3.60-3.85

208

(m, 2H), 3.95-4.00 (m, 2H), 4.60 (m, 1H), 4.85-4.98 (m, 3H), 6.97 (d, J=8.7 Hz, 1H), 6.90-7.00 (m, 2H), 7.15 (m, 1H), 7.20-7.45 (m, 10H), 10.15 (br s, 1H); MS (ion spray) 733.5 (M+1); Anal. Calc'd for $C_{40}H_{56}N_6O_7$: C, 65.55; H, 7.70; N, 11.47. Found: C, 65.44; H, 7.49; N, 11.59.

Examples 42 and 43

Reaction of the product of Preparation 89 (0.55 g, 0.75 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.4 g (75%) of the desired product as a mixture diastereomers. This material was resolved by HPLC (Kromsil CHI-DMP chiral stationary phase, 3A alcohol/dimethylethylamine/heptane) to provide the desired diastereomers, both isolated as white solids after acidification with hydrochloric acid as described in Example 7:


Example 42. (isomer 1): 1 H-NMR (d, DMSO) 0.75 (d, J = 6.4 Hz, 1H), 0.85-1.00 (m, 5H), 1.25-1.40 (m, 2H), 1.40-1.50 (m, 2H), 1.50-1.60 (m, 6H), 1.60-1.75 (m, 4H), 2.60=2.70 (m, 2H), 3.00 (m, 1H), 3.60-3.75 (m, 3H), 3.95-4.00 (m, 2H), 4.52 (s, 2H), 4.75 (m, 1H), 4.88 (m, 1H), 6.89 (d, J = 14 Hz, 1H), 7.00-7.05 (m, 2H), 7.20-7.40 (m, 9H), 8.10 (m, 1H), 8.20-8.30 (m, 3H), 8.60 (m, 1H), 11.02 (br s, 1H); $t_R = 5.90$ min; MS (high res) calc'd for $C_{35}H_{49}N_6O_5$: 633.3764. Found: 633.3768. Anal. Calc'd for $C_{35}H_{48}N_6O_5$: 2.3HC1: C, 58.66; H, 7.07; N, 11.73. Found: C, 58.59; H, 6.99; N, 11.46.

Exampl 43. (isomer 2): $^{1}H-NMR$ (d, DMSO) 0.75 (d, J = 6.4 Hz, 1H), 0.85-1.00 (m, 5H), 1.25-1.40 (m, 2H), 1.40-1.50

209

(m, 2H), 1.50-1.60 (m, 6H), 1.60-1.75 (m, 4H), 2.60=2.70 (m, 2H), 3.00 (m, 1H), 3.60-3.75 (m, 3H), 3.95-4.00 (m, 2H), 4.52 (s, 2H), 4.75 (m, 1H), 4.88 (m, 1H), 6.89 (d, J = 14 Hz, 1H), 7.00-7.05 (m, 2H), 7.20-7.40 (m, 9H), 8.10 (m, 1H), 8.20-8.30 (m, 3H), 8.60 (m, 1H), 11.02 (br s, 1H); $t_R = 7.47$ min; MS (high res) calc'd for $C_{35}H_{49}N_6O_5$: 633.3764. Found: 633.3762. Anal. Calc'd for $C_{35}H_{49}N_6O_5$ HCl: C, 59.57; H, 7.14; N, 11.91. Found: C, 59.74; H, 7.30; N, 11.72.


Preparation 90



Reaction of 4-phenoxyphenylacetic acid (25.0 g, 110 mmol) and p-toluenesulfonic acid (5.0 g, 26 mmol) in absolute ethanol (150 mL) as in described in Preparation 1 gave 27.6 g (98%) of the desired product as a yellow oil: 1 H-NMR (d, DMSO) 1.18 (t, J = 7.2 Hz, 3H), 3.64 (s, 2H), 4.08 (q, J = 7.2 Hz, 2H), 6.90-7.00 (m, 4H), 7.13 (t, J = 7.5 Hz, 1H), 7.23 (d, J = 8.7 Hz, 2H), 7.40 (t, J = 5.7 Hz, 2H); MS (ion spray) 257.2 (M+1); Anal. Calc'd for $C_{15}H_{16}O_{3}$: C, 74.98; H, 6.29. Found: C, 74.88; H, 6.31.

210

Preparation 91



Reaction of the product of Preparation 90 (10.0 g, 39.0 mmol), N-bromosuccinimide (7.2 g, 40.2 mmol) and 48% HBr (4 drops) in carbon tetrachloride (40 mL) as described in Preparation 2 gave 11.9 g (92%) of the desired product as a colorless oil: $^{1}\text{H-NMR}$ (d, DMSO) 1.21 (t, J = 7.3 Hz, 3H), 4.15-4.30 (m, 2H), 5.94 (s, 1H), 6.95-7.15 (m, 4H), 7.20 (m, 1H), 7.40-7.50 (m, 2H), 7.52-7.70 (m, 2H); MS (FD) 334, 336 (M+); Anal. Calc'd for $C_{16}H_{15}BrO_{3}$ 0.05CHCl3: C, 56.51; H, 4.45. Found: C, 56.85; H, 4.27.

Preparation 92

Reaction of the product of Preparation 91 (10.9 g, 33.0 mmol), 4-nitroimidazole (4.5 g, 39.6 mmol) and potassium carbonate (13.4 g, 99.0 mmol) in dimethylformamide (150 mL) as described in Preparation 3 gave 5.92 g (49%) of the desired product as a yellow oil: $^{1}\text{H-NMR}$ (d, DMSO) 1.17 (t, J = 6.8 Hz, 3H), 4.25 (q, J = 7.2 Hz, 2H), 6.60 (s, 1H), 7.00-7.10 (m, 4H), 7.17 (t, J = 7.2 Hz, 1H), 7.43 (t, J = 6.0 Hz, 2H), 7.53 (d, J = 6.8 Hz, 2H), 7.94 (s, 1H), 8.41 (s, 1H); MS (ion spray) 368.2 (M+1); Anal. Calc'd for $C_{19}H_{17}N_3O_5$ 0.15CHCl3: C, 59.30; H, 4.49; N, 10.91. Found: C, 59.55; H, 4.73; N, 10.97.

Preparation 93

PCT/US98/17201

Reaction of the product of Preparation 92 (1.58 (4.3 carbon (0.8 with 10% palladium on tetrahydrofuran (70 mL) followed by coupling with the product of Preparation 1d (1.64 g, 4.3 mmol), hydroxybenzotriazole (0.7 g, 4.7 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (1.04)4.7 mmol) as described in Preparation 4 gave 1.92 g (62%) of the desired product as a tan foam: 1H-NMR (d, DMSO) 1.20 (t, J = 7.2 Hz, 3H), 1.25-1.35 (m, 15H), 3.57 (m, 1H),3.70 (m, 1H), 4.25 (q, J = 7.2 Hz, 2H), 4.45-4.47 (m, 2H), 4.60 (m, 1H), 6.43 (s, 1H), 7.00-7.10 (m, 4H), 7.20 (m, 1H), 7.25-7.35 (m, 6H), 7.35-7.45 (m, 6H), 7.55 (s, 6H)1H), 10.20 (br s, 1H); MS (ion spray) 700.7 (M+1); Anal. Calc'd for $C_{38}H_{45}N_5O_8$: C, 65.22; H, 6.48; N, 10.01. Found: C, 65.12; H, 6.43; N, 9.87.

Preparation 94

Reaction of the product of Preparation 93 (1.72 g, 2.5 mmol) and lithium hydroxide (0.07 g, 3.0 mmol) in dioxane (50 mL) and water (25 mL) as described in Preparation 5 gave 1.68 g (100%) of the desired product as a tan foam: 1 H-NMR (d, DMSO) 1.25-1.40 (m, 15H), 3.60 (m, 1H), 3.70 (m, 1H), 4.45-4.50 (m, 2H), 4.57 (m, 1H), 6.25 (s, 1H), 7.00-7.07 (m, 4H), 7.15-7.35 (m, 8H), 7.35-7.45 (m, 5H),

212

7.55 (s, 1H), 10.20 (br s, 1H), 13.55 (br s, 1H); MS (ion spray) 672.6 (M+1); Anal. Calc'd for $C_{36}H_{41}N_5O_8$: C, 64.37; H, 6.15; N, 10.43. Found: C, 64.56; H, 6.37; N, 10.23.

Preparation 95

Reaction of the product of Preparation 94 (0.45 g, 0.67 mmol), pyrrolidine (0.07 mL, 0.67 mmol), hydroxybenzotriazole (0.1 g, 0.74 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.15 g, mmol)in dimethylformamide (30 mL) as described in Preparation 6 gave 0.22 g (45%) of the desired product as a white foam: $^{1}H-NMR$ (d, DMSO) 1.20-1.40 (m, 15H), 1.65-1.90 (m, 4H), 3.05 (m, 1H), 3.25-3.45 (m, 2H), 3.55-3.75 (m, 3H), 4.45-4.50 (m, 2H), 4.60 (m, 1H), 6.43 (s, 1H), 7.05 (t, J = 8.7 Hz, 3H), 7.20 (m, 1H), 7.25-7.30(m, 7H), 7.35-7.50 (m, 7H), 10.20 (br s, 1H); MS (ion)spray) 725.7 (M+1); Anal. Calc'd for $C_{40}H_{48}N_6O_7$: C, 66.28; H, 6.68; N, 11.59. Found: C, 66.42; H, 6.68; N, 11.59.

213

Example 44


Reaction of the product of Preparation 95 (0.22 g, 0.3 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.2 g (100%) of the desired mixture of isomers: 1 H-NMR (d, DMSO) 1.45-1.55 (m, 6H), 1.70-1.90 (m, 4H), 2.95 (m, 1H), 3.25-3.45 (m, 2H), 3.50-3.90 (m, 3H), 4.45-4.55 (m, 2H), 4.75 (m, 1H), 6.60 (m, 1H), 7.00-(m, 3H), 7.20 (m, 1H), 7.25-7.50 (m, 12H), 7.98 (m, 1H), 8.15-8.30 (m, 3H), 8.52 (t, J = 7.6 Hz, 1H), 10.88 (br s, 1H); MS (ion spray) 625.4 (M+1); Anal. Calc'd for $C_{35}H_{40}N_6O_5$ ·2HCl; C, 60.26; H, 6.07; N, 12.05. Found: C, 60.02; H, 6.01; N, 11.81.

Preparation 96

Reaction of the product of Preparation 94 (0.6 g, 0.9 (0.1 mL,0.9 mmol), 4-methylpiperidine hydroxybenzotriazole (0.14 g, 1.0 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.2 g, mmol) in dimethylformamide (30 mL) as described Preparation 6 gave 0.4 g (59%) of the desired product as a white foam: 1H-NMR is consistent with structure; MS (ion spray) 753.5 (M+1); Anal Calc'd for $C_{42}H_{52}N_6O_7$: C, 67.00; H, 6.96; N, 11.16. Found: C, 66.73; H, 6.91; N, 11.04.


214

Example 45



Reaction of the product of Preparation 96 (0.34 g, 0.45 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.27 g (83%) of the desired mixture of isomers as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (high res) calc'd for $C_{37}H_{45}N_{6}O_{5}$: 653.3451. Found: 653.3446.


Preparation 97



Reaction of biphenylacetic acid (25.2 g, 119 mmol) and ptoluenesulfonic acid (3.3 g, 17 mmol) in absolute ethanol (250 mL) as described in Preparation 1 gave 25.4 g (89%) of the desired product as a yellow oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 240.1 (M+); Anal. Calc'd for $C_{16}H_{16}O_{2}$: C, 79.97; H, 6.71. Found: C, 79.75; H, 6.59.

215

Preparation 98



Reaction of the product of Preparation 97 (18.0 g, 75.0 mmol), N-bromosuccinimide (13.7 g, 77.25 mL) and 48% HBr (4 drops) in carbon tetrachloride (80 mL) as described in Preparation 2 gave 22.56 g (94%) of the desired product as a yellow oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 318, 320 (M+); Anal. Calc'd for $C_{16}H_{15}BrO_{2}$ 0.05Chydrochloric acid₃: C, 60.21; H, 4.74. Found: C, 59.50; H, 4.75.

Preparation 99

To a slurry of sodium hydride (2.42 g, 60.5 mmol) in dimethylformamide (200 mL) stirring at temperature was added 4-nitroimidazole (6.9 g, 60.5 After 10 min, the product of Preparation 98 (17.62 g, 55.0 mmol) was added. After 16 h, the reaction mixture was concentrated and the residue was slurried in ethyl acetate then filtered. The resulting oil was partitioned between ethyl acetate and water then extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The resulting oil was absorbed onto silica gel and purified by flash chromatography (silica gel, 30-50% ethyl acetate/hexanes) to yield 12.0 g (62%) of the desired product as a yellow viscous oil: ¹H-NMR is consistent with structure; MS (FD) 351 (M+).

PCT/US98/17201 WO 99/08697

216

Preparation 100

Reduction of the product of Preparation 99 (2.0 g, 5.8 mmol) under a hydrogen atmosphere with 10% palladium on carbon (0.8 g) and tetrahydrofuran (70 mL) followed by coupling with the product of Preparation 1d (2.2 g, 5.8 mmol), 1-hydroxybenzotriazole (0.86 g, 6.4 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.3 g, 6.4 mmol) as described in Preparation 4 gave 0.7 g (18%) of the desired product as a tan foam: 1H-NMR is consistent with structure; MS (FD) 683 (M+); Anal. Calc'd for $C_{38}H_{45}N_5O_7$: C, 66.75; H, 6.63; N, 10.34. Found: C, 66.79; H, 6.48; N, 10.32.

Preparation 101

Reaction of the product of Preparation 100 (0.7 g, 1.0 mmol) and lithium hydroxide (0.03 g, 1.2 mmol) in dioxane (20 mL) and water (10 mL) as described in Preparation 5gave 0.66 g (100%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 656 (M+); Anal. Calc'd for $C_{36}H_{41}N_5O_7$: C, 65.94; H, 6.30; N, 10.68. Found: C, 65.90; H, 6.37; N, 10.42.

WO 99/08697

217

Reaction of the product of Preparation 101 (0.7 g, 1.1 mmol) withy 4-methylpiperidine (0.13 mL, 1.1 mmol), 1hydroxybenzotriazole (0.17 g, 1.2 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.25 g, 1.2 in dimethylformamide (40 mL)as described Preparation 6 gave 0.52 g (65%) of the desired product as a tan foam: 1H-NMR is consistent with structure; MS (FD) 728.4 (M+); Anal. Calc'd for $C_{37}H_{47}F_3N_6O_6$: C, 60.98; H, 6.50; N, 11.53. Found: C, 61.18; H, 6.35; N, 11.44.

Examples 46 and 47

Reaction of the product of Preparation 102 (0.36 g, 0.49 mmol) and trifluoroacetic acid (4 mL) in dichloromethane (12 mL) as described in Example 1 gave 0.3 g (88%) of the isomers. Resolution of mixture of diastereomers (4 g, 3.6 mmol) by HPLC (Kromsil CHI-DMP chiral stationary phase, 3A alcohol/dimethylethylamine /heptane eluant) provided provided 0.6 (16 %) of isomer 1 and 0.5 mg (12 %) of isomer 2, both isolated as white solids after formation of their respective hydrochloride salts as described in Example 7:


Example 46. (isomer 1). H-NMR is consistent with structure; $t_R = 6.9 \text{ min}$; MS (ion spray) 637.4 (M+1); Anal.

218

Calc'd for $C_{37}H_{44}N_6O_4$ 2.5HCl: C, 61.05; H, 6.44; N, 11.54. Found: C, 60.89; H, 6.53; N, 11.25.


Example 47. (isomer 2) $^{1}\text{H-NMR}$ is consistent with structure; $t_R = 9.2 \text{ min}$; MS (ion spray) 637.4 (M+1); Anal. Calc'd for $C_{37}\text{H}_{44}\text{N}_6\text{O}_4$ 2.6HCl: C, 60.75; H, 6.42; N, 11.49. Found: C, 60.67; H, 6.63; N, 11.18.

Preparation 103



Reaction of 3-fluorophenylacetic acid (15.0 g, 97.0 mmol) and p-toluenesulfonic acid (3.0 g, 16 mmol) in absolute ethanol as described in Preparation 1 gave 16.5 g (94%) of the desired product as a colorless oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 182 (M+); Anal. Calc'd for $C_{10}H_{11}FO_{2}$: C, 65.92; H, 6.09. Found: C, 64.94; H, 5.99.

Preparation 104



Reaction of the product of Preparation 103 (15.0 g, 82 mmol), N-bromosuccinimide (15.0 g, 84.5 mmol) and 48% HBr (4 drops) in carbon tetrachloride (80 mL) as described in Preparation 2 gave 19.2 g (90%) of the desired product as a colorless oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 259, 261 (M+); Anal. Calc'd for $C_{10}\text{H}_{10}\text{BrFO}_2$: C, 46.00; H, 3.86. Found: C, 45.71; H, 3.90.

219

Reaction of the product of Preparation 104 (15.0 g, 58.0 mmol), 4-nitroimidazole (7.8 g, 63.8 mmol) and sodium hydride ((2.8 g, 63.8 mmol) in dimethylformamide (200 mL) as in desribed in Preparation 3 gave 11.13 g (65%) of the desired product as a yellow oil: ¹H-NMR is consistent with structure; MS (FD) 293 (M+); Anal. Calc'd for $C_{13}H_{12}FN_3O_4$: C, 53.24; H, 4.12; N, 14.33. Found: C, 53.12; H, 4.22; N, 14.47.

Preparation 106

Reaction of the product of Preparation 105 (1.7 g, 5.8 with 10% palladium on carbon (0.7 q) in tetrahydrofuran (40 mL) under a hydrogen atmosphere followed by coupling with the product of Preparation 1d (2.2 g, 5.8 mmol), 1-hydroxybenzotriazole (0.86 g, 6.4 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.3 g, 6.4 mmol) as described in Preparation 4 gave 2.05 g (60%) of the desired product as a yellow foam: 1H-NMR is consistent with structure; MS (FD) 625 (M+); Anal. Calc'd for $C_{32}H_{40}FN_5O_7$: C, 61.43; H, 6.44; N, 11.19. Found: C, 61.28; H, 6.64; N, 11.32.

220

Reaction of the product of Preparation 104 (0.12 g, 3.2 mmol) and lithium hydroxide (0.09 g, 3.84 mmol) in dioxane (40 mL) and water (20 mL) as described in Preparation 5 gave 1.91 g (100%) of the desired product as a tan foam: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 598 (M+); Anal. Calc'd for $C_{30}H_{36}FN_5O_7$: C, 60.29; H, 6.07; N, 11.72. Found: C, 60.21; H, 6.41; N, 11.06.

Preparation 107

Reaction of the product of Preparation 106 (0.7 g, 1.2 mmol), 4-methylpiperidine (0.14 mL, 1.2 mmol), 1-hydroxybenzotriazole (0.18 g, 1.3 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.27 g, 1.3 mmol) in dimethylformamide (100 mL) as described in Preparation 6 gave 0.52 g (64%) of the desired product as a white solid: 1 H-NMR is consistent with structure; MS (FD) 678 (M+); Anal. Calc'd for $C_{36}H_{47}FN_{6}O_{6}$: C, 63.70; H, 6.98; N, 12.38. Found: C, 63.62; H, 7.10; N, 12.31.

Example 48

221

Reaction of the product of Preparation 107 (0.51 g, 0.75 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.24 g (49%) of the desired mixture of isomers as a yellow solid: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 578 (M+); Anal. Calc'd for $C_{31}H_{39}FN_6O_4$ 2.7HCl: C, 54.99; H, 6.21; N, 12.41. Found: C, 54.97; H, 6.23; N, 12.40.

Preparation 108

Reaction of the product of Preparation 106 (0.7 g, 1.2 pyrrolidine 1.2 mmol), 1-(0.1 mL, mmol), hydroxybenzotriazole (0.18 g, 1.3 mmol) and dimethylaminopropyl)-3-ethylcarbodiimide (0.27 in dimethylformamide (40 mL) as desribed in mmol) Preparation 6 gave 0.54 g (69%) of the desired product as a yellow foam: 1H-NMR is consistent with structure; MS (FD) 650 (M+); Anal. Calc'd for C34H43FN6O6'0.2CHCl3: C, 60.89; H, 6.45; N, 12.46. Found: C, 60.91; H, 6.39; N, 12.36.

Example 49

222

Reaction of the product of Preparation 108 (0.4 g, 0.6 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as desribed in Example 1 gave 0.3 g (79%) of the desired mixture of isomers as a yellow solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 550 (M+); Anal. Calc'd for $C_{29}H_{35}FN_{6}O_{4}$ 2.2HCl: C, 55.21; H, 5.94; N, 13.32. Found: C, 55.07; H, 5.91; N, 12.53.

Preparation 109



Reaction of 2-fluorophenylacetic acid (15.0 g, 97.0 mmol) and p-toluenesulfonic acid (2.8 g, 14.5 mmol) in absolute ethanol (100 mL) as described in Preparation 1 gave 17.0 g (96%) of the desired product as a colorless oil: ¹H-NMR is consistent with structure; MS (FD) 182 (M+).

223

Preparation 110



Reaction of the product of Preparation 109 (15.0 g, 82 mmol), N-bromosuccinimde (15.0 g, 84.5 mmol) and 48% HBr (3 drops) in carbon tetrachloride (80 mL) as described in Preparation 2 gave 21 g (98%) of the desired product as a colorless oil: ¹H-NMR is consistent with structure; MS (FD) 260 (M+).

Preparation 111

Reaction of the product of Preparation 110 (15.0 g, 58 mmol), 4-nitroimidazole (7.8 g, 63.8 mmol) and sodium hydride (2.8 g, 63.8 mmol) in dimethylformamide (200 mL) as described in Preparation 3 gave 11.36 g (67%) of the desired product as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 293.1 (M+); Anal. Calc'd for $C_{13}H_{12}FN_3O_4$: C, 53.24; H, 4.12; N, 14.33. Found: C, 53.54; H, 4.18; N, 14.11.

224

Pr paration 112

Reaction of the product of Preparation 111 (1.7 g, 5.8 mmol) with 10% palladium on carbon (0.7 g) in tetrahydrofuran (50 mL) under a hydrogen atmosphere followed by coupling with the product of Preparation 1d (2.2 g, 5.8 mmol), 1-hydroxybenzotriazole (0.86 g, 6.4 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.3 g, 6.4 mmol) as described in Preparation 4 gave 2.4 g (67%) of the desired product as a tan foam: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 625 (M+); Anal. Calc'd for $C_{32}\text{H}_{40}\text{FN}_5\text{O}_7$: C, 61.43; H, 6.44; N, 11.19. Found: C, 61.51; H, 6.50; N, 11.34.

Preparation 113

Reaction of the product of Preparation 112 (2.35 g, 3.8 mmol) and lithium hydroxide (0.1 g, 4.6 mmol) in dioxane (40 mL) and water (20 mL) as desribed in Preparation 5 gave 2.27 g (100%) of the desired product as a tan foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 598 (M+); Anal. Calc'd for $C_{30}H_{36}FN_{5}O_{7}$: C, 60.29; H, 6.07; H, 11.72. Found: C, 60.08; H, 6.28; N, 11.42.

225

Reaction of the product of Preparation 113 (0.7 g, 1.2 mmol), 4-methylpiperidine (0.14 mL, 1.2 mmol), 1-hydroxybenzotriazole (0.18 g, 1.3 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.27 g, 1.3 mmol) in dimethylformamide (100 mL) as described in Preparation 6 gave 0.56 g (69%) of the desired product as a tan foam: 1 H-NMR is consistent with structure; MS (FD) 678.2 (M+); Anal. Calc'd for $C_{36}H_{47}FN_{6}O_{6}$: C, 63.70; H, 6.98; N, 12.38. Found: C, 63.44; H, 7.05; N, 12.10.

Example 50

Reaction of the product of Preparation 114 (0.53 g, 0.78 mmol) and trifluoroacetic acid (4 mL) in dichloromethane (12 mL) as described in Example 1 gave 0.38 g (75%) of the desired mixture of isomers as a yellow solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 578 (M+); Anal. Calc'd for $C_{31}H_{39}FN_{6}O_{4}$ 2.2HCl: C, 56.51; H, 6.30; N, 12.75. Found: C, 56.45; H, 6.10; N, 12.43.

226

Reaction of the product of Preparation 113 (0.7 g, 1.2 mmol), pyrrolidine (0.1 mL, 1.2 mmol), 1 mmol) (0.18 1.3 and 1-(3hydroxybenzotriazole g, dimethylaminopropyl)-3-ethylcarbodiimide (0.27 mmol) in dimethylformamide (100 mL) as described in Preparation 6 gave 0.6 g (77%) of the desired product as a tan foam. ¹H-NMR is consistent with structure; MS (FD) 650 (M+); Anal. Calc'd for C₃₄H₄₃FN₆O₆: C, 62.75; H, 6.66; N, 12.91. Found: C, 62.53; H, 6.58; N, 12.71.

Example 51

Reaction of the product of Preparation 115 (0.46 g, 0.7 mmol) and trifluoroacetic acid (4 mL) indichloromethane (12 mL) as described in Example 1 gave 0.44 g (100%) of the desired mixture of isomers as a white foam: $^1\text{H-NMR}$ is consistent with structure. MS (high res) calc'd for $C_{29}H_{36}FN_6O_4$: 551.2782. Found: C, 551.2779. Anal. Calc'd for $C_{29}H_{35}FN_6O_4$ 2HCl: C, 55.86; H, 5.98; N, 13.48. Found: C, 56.09; H, 5.91; N, 13.44.

Reaction of 3-trifluoromethylphenylacetic acid (15.0 g, 73.4 mmol) and p-toluenesulfonic acid (3 g, 15.6 mmol) in absolute ethanol (200 mL) as described in Preparation 1 gave 15.6 g (93%) of the desired product as a colorless oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 232 (M+); Anal. Calc'd for $C_{11}H_{11}F_{3}O_{2}$: C, 56.90; H, 4.77. Found: C, 56.93; H, 4.65.

Preparation 117

Reaction of the product of Preparation 116 (10.0 g, 44.0 mmol), N-bromosuccinimide (8.0 g, 45.3 mmol) and 48% HBr (4 drops) in carbon tetrachloride (70 mL), as desribed in Preparation 2 gave 11.2 g (82%) of the desired product as a colorless oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 264 (M+); Anal. Calc'd for $C_{11}\text{H}_{10}\text{BrF}_{3}\text{O}_{2}$: C, 42.47; H, 3.24. Found: C, 42.37; H, 3.26.

Preparation 118

Reaction of the product of Preparation 117 (11.2 g, 36.0 mmol), 4-nitroimidazole (4.9 g, 43.2 mmol) and sodium hydride (1.7 g, 43.2 mmol) in dimethylformamide (180 mL) as described in Preparation 3 gave 6.22 g (50%) of the desired product as a yellow oil: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 343.1 (M+); Anal. Calc'd for $C_{14}H_{12}F_3N_3O_4$: C, 48.99; H, 3.52; N, 12.24. Found: C, 48.74; H, 3.63; N, 12.06.

228

Preparation 119

Reaction of the product of Preparation 118 (2.0 g, 5.8 mmol) with 10% palladium on carbon (0.6 tetrahydrofuran (80 mL) under an atmosphere of hydrogen followed by coupling with the product of Preparation 1d (2.2 g, 5.8 mmol), 1-hydroxybenzotriazole (0.86 g, 6.4 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.3 g, 6.4 mmol) as described in Preparation 4 gave 1.82 g (47%) of the desired product as a tan foam: 1H-NMR is consistent with structure; MS (FD) 675.4 (M+); Anal. Calc'd for $C_{33}H_{40}F_3N_5O_7$: C, 58.66; H, 5.97; N, 10.36. Found: C, 58.67; H, 5.87; N, 10.51.

Preparation 120

Reaction of the product of Preparation 120 (1.67 g, 2.5 mmol) and lithium hydroxide (0.07 g, 2.8 mmol) in dioxane (40 mL) and water (20 mL) as described in Preparation 5 gave 1.60 g (99%) of the desired product as a yellow foam: H-NMR is consistent with structure; MS (FD) 648 (M+); Anal. Calc'd for $C_{31}H_{36}F_3N_5O_7$: C, 57.49; H, 5.60; N, 10.81. Found: C, 57.52; H, 5.62; N, 10.75.

229

Reaction of the product of Preparation 120 (0.6 g, 0.93 mmol), 4-methylpiperidine (0.11 mL, 0.93 mmol), hydroxybenzotriazole (0.13 g, 1.02 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.12 1.02 mmol)in dimethylformamide (40 mL) described as Preparation 6 gave 0.55 g (81%) of the desired product as a tan foam: 1H-NMR is consistent with structure; MS (FD) 728.9 (M+); Anal. Calc'd for $C_{37}H_{47}F_3N_6O_6$: C, 60.98; H, 6.50; N, 11.53. Found: C, 60.81; H, 6.57; N, 11.69.

Example 52

Reaction of the product of Preparation 121 (0.5 g, 0.68 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.33 g (69%) of the desired mixture of isomers as a yellow solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 628.8 (M+1); Anal. Calc'd for $C_{32}H_{39}F_{3}N_{6}O_{4}$ 2.3HCl: C, 53.94; H, 5.84; N, 11.79. Found: C, 53.89; H, 5.92; N, 11.65.

230

Reaction of 3-trifluoromethylphenylacetic acid (15.0 g, 73.4 mmol) and p-toluenesulfonic acid (2.8 g, 14.5 mmol)in absolute ethanol (200 mL) as described in Preparation 1 gave 16.11 g (94%) of the desired product as a colorless oil: 1H-NMR is consistent with structure; MS (FD) 232 (M+). Anal. Calc'd for $C_{11}H_{11}F_3O_2$: C, 56.90; H, 4.77. Found: C, 56.64; H, 4.90.

Preparation 123



Reaction of the product of Preparation 122 (15.0 g, 65 mmol), N-bromosuccinimide (11.9 g, 67.0 mmol) and 48% HBr (4 drops) in carbon tetrachloride (80 mL) as described in Preparation 2 gave 17.1 g (85%) of the desired product as a colorless oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 311, 313 (M+).

231

Preparation 124

Reaction of the product of Preparation 123 (15.0 g, 48.0 mmol), 4-nitroimidazole (6.0 g, 52.8 mmol) and sodium hydride (2.1 g, 52.8 mmol) in dimethylformamide (200 mL) as described in Preparation 3 gave 12.25 g (74%) of the desired product as a yellow oil: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 343 (M+); Anal. Calc'd for $C_{14}\text{H}_{12}\text{F}_3\text{N}_3\text{O}_4$: C, 48.99; H, 3.52; N, 12.24. Found: C, 49.10; H, 3.58; H, 12.22.

Preparation 125

Reaction of the product of Preparation 124 (2.0 g, 5.8 mmol) with 10% palladium on carbon (1.0 g)in tetrahydrofuran (60 mL) under a hydrogen atmosphere follwed by coupling with the product of Preparation 1d (2.2 g, 5.8 mmol), 1-hydroxybenzotriazole (0.86 g, 6.4 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.3 g, 6.4 mmol) as described in Preparation 4 gave 3.16 g (93%) of the desired product as a tan foam: 1 H-NMR is consistent with structure; MS (FD) 675.4 (M+); Anal. Calc'd for $C_{33}H_{40}F_{3}N_{5}O_{7}$: C, 58.66; H, 5.97; N, 10.36. Found: C, 58.81; H, 6.04; N, 10.12.

WO 99/08697

Reaction of the product of Preparation 125 (2.78 g, 4.1 mmol) with lithium hydroxide (0.12 g, 4.9 mmol) in dioxane (40 mL) and water (20 mL) as described in Preparation 5 gave 2.6 g (98%) of the desired product as a yellow foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 648.2 (M+); Anal. Calc'd for $C_{31}H_{36}F_{3}N_{6}O_{7}$: C, 57.49; H, 5.60; N, 10.81. Found: C, 58.06; H, 6.14; N, 10.27.

Preparation 127

Reaction of the product of Preparation 126 (0.7 g, 1.1 mmol), 4-methylpiperidine (0.13 mL, 1.1 mmol), 1.2 mmol) and 1-(3hydroxybenzotriazole (0.17 g, dimethylaminopropyl)-3-ethylcarbodiimide (0.25 1.2 g, mmol) in dimethylformamide (30 mL) as described Preparation 6 gave 0.32 g (40%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 728 (M+); Anal. Calc'd for C₃₇H₄₇F₃N₆O₆: C, 60.98; H, 6.50; N, 11.53. Found: C, 60.76; H, 6.59; N, 11.36.

Example 53

233

Reaction of the product of Preparation 127 (0.3 g, 0.41 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.28 g (97%) of the desired mixture of isomers as a white solid: 1H-NMR is consistent with structure; MS (FD) 628 (M+); Anal. Calc'd for $C_{32}H_{39}F_3N_6O_4$ 2.2HCl: C, 54.22; H, 5.86; N, Found: C, 54.33; H, 5.84; N, 11.56.

Preparation 128

Reaction of the product of Preparation 126 (0.5 g, 0.77 (0.07 mL, 0.77 mmol), pyrrolidine mmol), 1-(0.12 g, 0.85 mmol) and 1-(3hydroxybenzotriazole dimethylaminopropyl)-3-ethylcarbodiimide (0.18 g, mmol) in dimethylformamide (30 mL) as described in Preparation 6 gave 0.2 g (37%) of the desired product as a tan solid: 1H-NMR is consistent with structure; MS (FD) 700 (M+); Anal. Calc'd for $C_{35}H_{43}F_3N_6O_6$ 0.4 H_2 0: C, 59.38; H, 6.24; N, 11.87. Found: C, 59.17; H, 6.24; N, 11.87.

234

Example 54

Reaction of the product of Preparation 128 (0.2 g, 0.29 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.18 g (100%) of the desired mixture of isomers as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 600 (M+).

Preparation 129

Reaction of the product of preparation 126 (0.75 g, 1.2 mmol), dimethylamine hydrochloride (0.1 g, 1.2 mmol), triethylamine (0.19 g, 1.3 mmol), 1-hydroxybenzotriazole (0.18 g, 1.3 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.27 g, 1.3 mmol) in dimethylformamide (40 mL) as described in Preparation 6 gave 0.49 g (60%) of the desired product as a tan foam: 1 H-NMR is consistent with structure; MS (FD) 675 (M+); Anal. Calc'd for $C_{33}H_{41}F_{3}N_{6}O_{6}$: C, 58.75; H, 6.13; N, 12.46. Found: C, 58.69; H, 6.12; N, 12.28.

235

Example 55

Reaction of the product of Preparation 129 (0.42 g, 0.62 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.3 g (75%) of the desired mixture of isomers as a yellow solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 574 (M+); Anal. Calc'd for $C_{28}H_{33}F_{3}N_{6}O_{4}$ 2.8 ·HCl: C, 48.70; H, 5.33; N, 12.42. Found: C, 49.84; H, 5.27: N, 12.09.

Preparation 130

Reaction of the product of Preparation 126 (0.5 g, 0.77 mmol), 4-(4-fluorobenzoyl)piperidine hydrochloride (0.19 g, 0.77 mmol), triethylamine (0.12 mL, 0.85 mmol), 1-hydroxybenzotriazole (0.12 g, 0.85 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.18 g, 0.85 mmol) in dimethylformamide (40 mL) as described in Preparation 6 gave 0.45 g (69%) of the desired product as a yellow foam: 1 H-NMR is consistent with structure; MS (FD) 836.8 (M+); Anal. Calc'd for $C_{43}H_{48}F_{4}N_{6}O_{7}$ 0.4 $H_{2}O$: C, 61.19; H, 5.83; N, 9.96. Found: C, 60.92; H, 5.56; N, 10.09.

Example 46

236

Reaction of the prodcut of Preparation 130 (0.4 g, 0.48 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.26 g (67%) of the desired mixture of isomers as a white solid. $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 736.7 (M+); Anal. Calc'd for $C_{38}H_{40}F_{4}N_{6}O_{5}$ 2.1 HCl: C, 56.12; H, 5.22; N, 10.33. Found: C, 56.08; H, 5.46; N, 10.38.

237

Preparation 131

Reaction of alpha-bromocyclohexylacetic acid (5.0 g, 21.0 mmol), 4-nitroimidiazole (2.6 g, 23.1 mmol) and sodium hydride (0.93 g, 23.1 mmol) in dimethylformamide (45 mL) as described in Preparation 3 gave 1.9 g (34%) of the desired product as a clear oil: $^1\text{H-NMR}$ is consistent with structure; MS (ion spray) 268 (M+1); Anal. Calc'd for $C_{12}H_{17}N_3O_4$: C, 53.92; H, 6.41; N, 15.72. Found: C, 53.63; H, 6.33; N, 15.77.

Preparation 132

Reaction of the product of Preparation 131 (1.4 g, 5.2 mmol) with 10% palladium on carbon (0.8 g) in tetrahydrofuran (60 mL) under a hydrogen atmosphere followed by coupling with the product of Preparation 1d (2.0 g, 5.2 mmol), 1-hydroxybenzotriazole (0.8 g, 5.7 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.2 g, 5.7 mmol) as described in Preparation 4 gave 2.09 g (65%) of the desired product as a tan foam. 1 H-NMR is consistent with structure; MS (ion spray) 600.4 (M+1); Anal. Calc'd for $C_{31}H_{45}N_{5}O_{7}$: C, 62.08; H, 7.56; N, 11.68. Found: C, 62.04; H, 7.53; N, 11.74.

238

Reaction of the product of Preparation 132 (2.0 g, 3.3 mmol) with lithium hydroxide (0.1 g, 4.0 mmol) in dioxane (50 mL) and water (25 mL) as described in Preparation 5 gave 1.9 g (99%) of the desired product as a tan foam: 1 H-NMR is consistent with structure; MS (ion spray) 586.4 (M+1); Anal. Calc'd for $C_{30}H_{43}N_{5}O_{7}$: C, 61.52; H, 7.40; N, 11.96. Found: C, 61.41; H, 7.42; N, 11.82.

Preparation 134

Reaction of the product of Preparation 133 (0.8 g, 1.4 mmol), 4-methylpiperidine (0.17 mL, 1.4 mmol), 1-hydroxybenzotriazole (0.21 g, 1.54 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.32 g, 1.54 mmol) in dimethylformamide (30 mL) as described in Preparation 6 gave 0.92 g (99%) of the desired product as a tan foam: 1 H-NMR is consistent with structure; MS (ion spray) 667.5 (M+1); Anal. Calc'd for $C_{36}H_{54}N_{6}O_{6}$: C, 64.84; H, 8.16; N, 12.60. Found: C, 64.55; H, 7.73; N, 12.26.

Example 57

239

Reaction of the product of Preparation 134 (0.7 g, 1.0 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.43 g (64%) of the desired mixture of isomers as a tan solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 567.6 (M+1); Anal. Calc'd for $C_{31}H_{46}N_{6}O_{4}$ 2HCl: C, 58.21; H, 7.56; N, 13.14. Found: C, 58.36; H, 7.33; N, 13.19.

Preparation 135

Reaction of the product of Preparation 133 (0.8 g, 1.4 mmol), dimethylamine hydrochloride (0.12 g, 1.4 mmol), triethylamine (0.22 mL, 1.54 mmol), hydroxybenzotriazole (0.21 g, 1.54 mmol) and dimethylaminopropyl)-3-ethylcarbodiimide (0.32 mmol) in dimethylformamide (30 mL) as described in Preparation 6 gave 0.86 g (100%) of the desired product as a tan foam: 1H-NMR is consistent with structure; MS (ion spray) 613.4 (M+1); Anal. Calc'd for $C_{32}H_{48}N_6O_6$: C, 62.72; H, 7.90; N, 13.72. Found: C, 62.44; H, 7.64; N, 13.57.

240

Example 58

Reaction of the product of Preparation 135 (0.7 g, 1.0 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 1 gave 0.43 g (64%) of the desired mixture of isomers as a tan solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 567.6 (M+1); Anal. Calc'd for $C_{31}H_{46}N_{6}O_{4}$ 2HCl: C, 58.21; H, 7.56; N, 13.14. Found: C, 58.36; H, 7.33; N, 13.19.

Preparation 136

To a suspension of 2-naphthyl acetic acid (49.37 g, 265.0 mmol) in carbon tetrachloride (55 mL) was added and thionyl chloride (80 mL). The mixture was heated to reflux for 20 minutes then cooled to ambient temperature. Carbon tetrachloride (125 mL), N-bromosuccinimide (56.60 g, 318.0 mmol) and hydrobromic acid (48% aq., 0.5 mL) were added. The mixture was heated to reflux for 30 min, filtered. to ambient temperature, cooled concentrated. The resulting material was dissolved in dichloromethane (200 mL) and excess ethanol (100 mL) was added dropwise. After 1 h, the reaction was concentrated and the resulting crude material was purified by flash chromatography(silica gel, 30% ethyl acetate/hexane) to yield a tan solid. This crude material was dissolved

241

dimethylformamide (200 mL) and 4-nitroimidazole (29.78 g, 263.5 mmol) and potassium carbonate (72.70 g, 526.8 mmol) were added. After 16 h, the reaction was concentrated to 100 mL. Ethyl acetate and water were added and the mixture washed with sodium bicarbonate and brine. The 1 dried over sodium sulfate and organic laver was The crude material was was purified by concentrated . flash chromatography (silica, 30% ethyl acetate/hexane) to yield 40.2 g (47%) of the desired product as a brown ¹H NMR (300 MHz, CDCl₃) - consistent with foam: structure; Anal. calcd. for $C_{17}H_{15}N_3O_4$; 62.76 C, 4.65 H, 12.92 N; found 60.54 C, 4.35 H, 12.04 N; ISMS (M+) - 326.

Preparation 137


Reaction of the product of Preparation 136 (4.80 g, 14.77 with 5% palladium on carbon (2.5 mmol) q) in tetrahydrofuran (100 mL) under a hydrogen atmosphere followed by coupling with the product of Preparation 1d (5.61 g, 14.77 mmol), EDCI (2.79 g, 16.25 mmol), 1hydroxybenzotriazole (2.00 g, 14.77 mmol), methylmorpholine (1.6 mL, 14.77 mmol) as described in Preparation 4 gave (6.04 g, 62%) of the desired product as a light orange foam: ¹H NMR (300 MHz, CDCl₃) consistent with structure; Anal. calcd. for C36H43N5O7; 65.74 C, 6.59 H, 10.65 N; found 64.02 C, 6.09 H, 10.13 N; ISMS (M+) - 658.

242

A solution of lithium hydroxide (0.38 g, 9.16 mmol) in water (50 mL) was added to a solution of the product of Preparation 137 (6.04 g, 9.16 mmol) in tetrahydrofuran (100 mL). After 30 min, water was added and the mixture The aqueous layer was washed with diethyl ether. adjusted to pH = 3.0 with sodium bisulfate, saturated with sodium chloride, and washed with ethyl acetate. combined organic extracts were dried over sodium sulfate, To the resulting crude material and concentrated. stirring at room temperature in dimethylformamide (50 mL) added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (2.08 g, 10.01 mmol), 1-hydroxybenzotriazole (1.24 g, 9.16 mmol) and 4-methylpiperidine (1.1 mL, 9.16 mmol). After 18 h, the reaction was quenched with saturated bicarbonate, and washed with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, and concentrated. The crude material was purified by flash chromatography (silica qel, methanol/dichloromethane) to yield 4.9 g (75 %) of the desired product as a pale yellow foam: 1H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C40H50N6O6; 67.58 C, 7.09 H, 11.82 N; found 65.60 C, 7.09 H, 11.50 N; ISMS (M+) - 711.

Examples 59 and 60

243



To a solution of of the product of Preparation 138 (4.90 6.89 mmol) stirring at room temperature dichloromethane (40 mL) and anisole (1.0 mL) was added to triflouroacetic acid (10 mL). After 3 hours, the reaction was quenched with saturated sodium bicarbonate and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate and concentrated. The resulting crude material was purified by flash chromatography (silica gel, 5% methanol/dichloromethane) to give the product as a mixture of diastereomers. This material was resolved by HPLC (Kromsil CHI-DMP chiral stationary phase, alcohol/ dimethylethylamine/heptane eluant) to provide the free amine of the desired products. The individual diastereomers were dissolved in ethyl acetate and treated with a saturated solution of hydrochloric acid in diethyl The resulting precipiate was filtered to yield the desired products (426779 - 0.64 g, 14%) (426780 -0.43 g, 9%) as tan solids: **Example 59.** 1 H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{35}H_{44}N_6O_4Cl_2$; 61.49 C, 6.49 H, 12.29 N; found 60.28 C, 6.38 H, 11.74 N; ISMS (M+) - 611. Example 60. ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{35}H_{44}N_6O_4Cl_2$; 61.49 C, 6.49 H, 12.29 N; found 47.81 C, 5.29 H, 9.83 N; ISMS (M+) - 611.

244

Reaction of the product of Preparation 136 (1.31g, 4.02 with 10% palladium on carbon (0.5 g) in tetrahydrofuran (50 mL) under a hydrogen atmosphere followed by coupling with the product of Preparation 1j (1.52g, 4.02 mmol), 1-hydroxybenzotriazole (0.68g, 4.42 and 1-(3-dimethylaminopropyl)-3mmol), ethylcarbodiimide(0.91g, 4.42 mmol) as described in Preparation 4 to give 1.06 g (38 %) of the title compound as a tan solid: ${}^{1}H$ NMR (${d}^{6}$ -DMSO, d): 1.22(m, 18H), 1.50(m, 4H), 2.55(m, 2H), 4.26(q, J = 9.0Hz, 2H), 4.37(bs, 1H), 5.75(s, 1H), 6.60(s, 1H), 7.02(bs, 1H), 7.16(m, 1H)7.22(m, 3H), 7.43(m, 1H), 7.50(d, J = 9.3Hz, 2H), 7.60(m, 1H)2H), 7.97(m, 3H), 10.21(m, 1H). Ion spray MS (M⁺ +1): 656.

Preparation 140

Reaction of the product of Preparation 139 (1.06 g, 1.62 mmol) with lithium hydroxide 75 mg, 1.78 mmol) in dioxane (30 mL) and water (15 mL) as described in Preparation 5 gave 1.01 g (100 %) of the title compound as a golden yellow solid: 1 H NMR (d 6 -DMSO, d): 1.20(m, 15H), 1.50(m, 4H), 2.55(m, 2H), 4.38(bs, 1H), 6.58(s, 1H), 7.02(bs,

245

1H), 7.17(m, 3H), 7.25(m, 3H), 7.35(m, 1H), 7.50(m, 2H), 7.58(m, 2H), 7.98(m, 3H), 8.09(m, 1H), 10.36(bs, 1H). Ion spray MS (M* +1): 628.

246

Preparation 141

To a solution of the product of Preparation 140 (500 mg, 0.80 mmol) dimethylamine hydrochloric acid (72 mg, 0.88 (0.12 mL, 0.88 mmol), triethylamine mmol), 0.88 mmol) hydroxybenzotriazole (134 mg, and 1-(3dimethylaminopropyl) -3-ethylcarbodiimide (18 mg, mmol) in dimethylformamide (20 mL) as described in Preparation 6 gave 342 mg (66 %) of the title compound as a white solid: ^{1}H NMR ($d^{6}-DMSO$, d): 1.27(m, 15H), 1.57(m, 4H), 2.55(m, 2H), 2.90(s, 3H), 2.95(s, 3H), 4.38(bs, 1H), 6.80(s, 1H), 7.02(bs, 1H), 7.15(m, 3H), 7.22(m, 3H), 7.35(m, 1H), 7.47(m, 2H), 7.57(m, 1.35(m, 1.35)2H), 7.88(s, 7.98 (m, 3H), 10.15 (bs, 1H). Ion spray MS $(M^{+}+1)$: 655.

Anal. $(C_{37}H_{46}N_6O_5)$: H,N;C: calcd 67.87; found 66.19.

Example 61

Reaction of the product of Preparation 141 (333 mg, 0.51 mmol) with trifluoroacetic acid (5 mL) in dichloromethane (17 mL) as described in Example 1 gave 52 mg (65 %) of a tan solid which was purified by HPLC (Kromosil CHI-DMP chiral stationary phase, 3A alcohol/dimethylethylamine/heptane eluant) to give the free amine

247

which was acidied with hydrochloric acid to provide the desired product: 1H NMR (^6-DMSO , d): 1.21(m, 6H), 1.57(m, 4H), 2.54(m, 2H), 2.90(s, 3H), 2.95(s, 3H), 4.41(bs, 1H), 6.82(s, 1H), 7.02(bs, 1H), 7.14(m, 3H), 7.24(m, 3H), 7.48(m, 2H), 7.57(m, 2H), 7.87(s, 1H), 7.97(m, 3H), 8.12(bs, 1H), 10.40(s, 1H). FAB+ exact MS (6 +1): 555.3084 calcd, 555.3079 found Anal. (6 C₃₂H₄₁N₆O₃Cl₃):C,H,N.

248

Preparation 142

Reaction of the product of Preparation 136 (8.7 g, 27 mmol) with 10% palladium on carbon (4.0 g) under a hydrogen atmosphere follwed by coupling with the product Preparation 1d (10.14 g, 26.7 mmol), of hydroxybenzotriazole (4.49 g, 29.3 mmol), and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (6.05 g, mmol) as described in Preparation 4 gave 5.4 g (31 %) of the title compound as a tan solid: ^{1}H NMR ($d^{6}-DMSO$, d): 1.26(t, J = 7.4Hz, 3H), 1.40(s, 9H), 1.55(m, 6H), 3.55(m, 1H), 4.02(s, 1H), 4.25(m, 2H), 4.50(dd, J = 10.0Hz, 2H), 4.86(s, 1H), 5.92(s, 1H), 7.02(d, J = 7.0Hz, 1H), 7.22(m, IH)8H), 7.33(m, 3H), 7.41(s, 1H), 7.49(m, 1H), 7.80(m, 2H), 9.22(bs, 1H). Ion spray MS (M⁺ +1): 658. Anal. $(C_{36}H_{43}N_5O_7):C,H,N.$

Preparation 143

Reaction of the product of Preparation 142 (5.39 g, 8.19 mmol) with lithium hydroxide (361 mg, 8.60 mmol) in dioxane (120 mL) and water (75 mL) as described in Preparation 5 gave 4.92 g (95 %) of the title compound as

249

a golden yellow solid: ^{1}H NMR ($^{6}-DMSO$, d): 1.28 (m, 15H), 3.57 (m, 1H), 3.66 (m, 1H), 4.43 (s, 2H), 4.48 (d, J=5.3Hz, 1H), 4.56 (bs, 1H), 5.75 (bs, 1H), 7.13 (bs, 1H), 7.26 (m, 6H), 7.31 (d, J=6.0Hz, 2H), 7.40 (m, 1H), 7.45 (m, 2H), 7.65 (s, 1H), 7.83 (m, 3H), 10.10 (bs, 1H). Ion spray MS (4 +1): 630.

Preparation 144

Reaction of the product of Preparation 143 (4.88 g, 7.75 mmol), dimethylamine (4.2 mL, 8.53 mmol, 2.0M tetrahydrofuran), 1-hydroxy-7-azabenzotriazole(1.16 1-(3-dimethylaminopropyl)-3-8.53 mmol) and ethylcarbodiimide (1.76 g, 8.53 mmol) in tetrahydrofuran (120 mL) as described in Preparation 6 gave 2.06 g (40 %) of the title compound as a yellow foam: $4^{1}H$ NMR $(d^{6}-$ DMSO, d): 1.28(m, 15H), 2.92(s, 3H), 2.95(s, 3H), 3.60(m, 15H)1H), 4.43(d, J = 4.5Hz), 4.57(bs, 1H), 6.83(s, 1H)1H). 7.24(m, 8H), 7.39(m, 1H), 7.50(m, 1H), 7.56(m, 2H), 7.88(s, 1H), 7.96(m, 3H). Ion spray MS $(M^{+} + 1)$: 657 Anal. $(C_{36}H_{44}N_6O_4)$: H,N;C: calcd 65.84; found 63.70.

250

Example 62

To a solution of glacial acetic acid saturated with dry hydrochloric acid (50 mL, ~3N in hydrochloric acid) stirring at room temperature was added the product of Preparation 144 (1.87 g, 2.85 mmol). After 2h, solution was concentrated, washed with aqueous sodium bicarbonate solution, dried over magnesium sulfate and concentratd. The resulting crude material was purified by HPLC (Column) to give 0.5 g of the desired isomer which was dissolved in ethyl acetate and added dropwise to a stirred solution of anhydrous diethyl ether saturated with hydrochloric acid. The resulting white precipitate was collected by filtration and dried to give 474 mg (79 %) white solid: ^{1}H NMR (d^{6} -DMSO, d): 1.47(m, 6H), 2.90(s, 3H), 2.95(s, 3H), 3.65(dd, J = 9Hz, 2H), 4.49(d, J =7.9Hz, 2H), 4.73 (m, 1H), 6.93 (s, 1H), 7.18 (s, 1H), 7.26(m, 6H), 7.49(d, J = 8.7Hz, 1H), 7.60(m, 2H), 7.84(d, 7.84)J = 10.5Hz, 1H), 7.98(m, 3H), 8.14(d, J = 9.4Hz, 2H), 8.45(d, J = 6.8Hz, 1H), 10.74(bs, 1H). FAB+ exact MS (M⁺ 557.2876 calculated, 557.2873 found $(C_{31}H_{39}N_6O_4Cl_3):H,N;C: calcd, 56.01; found, 56.72.$

251

EXAMPLES PART 2B

Preparation 145



To a solution of m-anisaldehyde, 27.2 g (200 mmol) and benzyltriethylammonium chloride, 2.46 g (10.8 mmol) stirring in chloroform (32 mL) at 56°C was added sodium hydroxide (50 mL of 50% aqueous solution) dropwise over a period of 2 h keeping the temperature between 54°C and 10 58°C. After 1 h, the solution was cooled, then poured into an ice/water mixture. The resulting mixture was washed with ether. The aqueous layers were acidified with 6 N sulfuric acid, then extracted with ether. The organic extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated to yield 36.4 g of an oil which was used without purification.


Preparation 146

To a solution of the compound of Preparation 145 (36.4 g, 20 200 mmol) stirring in absolute ethanol (200 mL) was slowly added a solution of concentrated sulfuric acid (20 mL). The resulting mixture was heated to reflux for 5 h then cooled to ambient temperature, poured into an ice/water mixture, and extracted with ether. 25 combined organic extracts were washed with saturated sodium bicarbonate and brine then dried over sodium sulfate and concentrated. The resulting material was purificed by flash chromatography (silica gel, 15% ethyl acetate/hexanes) to give 13.6 g (33% over the two steps) 30 of the desired product an oil: 1H-NMR is consistent with structure.

PCT/US98/17201 WO 99/08697

252

Preparation 147



To a solution of the product of Preparation 146 (4.0 g, 19 mmol) stirring in chloroform (50 mL) at 0°C was added phosphorus tribromide (2.1 mL, 21 mmol). The reaction mixture was warmed to ambient temperature and stirred for 5 h, then poured into an ice/water mixture and extracted with chloroform. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered 10 and concentrated. The residue was purified by flash chromatography (silica gel, chloroform) to yield 3.9 g (75%) of the desired product as an oil: 1H-NMR is consistent with structure; MS (FD) 272, 274 (M+).

15

20

25

30

5

Preparation 148



To a slurry of sodium hydride (0.86 g, 20 mmol) stirring in dimethylformamide (30 mL) at room temperature was added 4-nitroimidazole (2.26 g, 20 mmol). The reaction was cooled to 0°C and 3.9 g (19 mmol) of the product of Preparation 147 was added. After 16 g, the mixture was slowly warming to ambient temperature. The reaction was poured into an ice/water mixture and extracted with ethyl acetate. The combined organic extracts were washed with water, brine, dried over sodium sulfate, and concentrated. The residue was purified by flash chromatography (silica gel, 40% ethyl acetate/hexanes) to yield 0.87 g (15%) of the desired product: 1H-NMR is consistent with structure; MS (ion spray) 306 (M+1);

253

Anal. Calc'd for $C_{14}H_{15}N_{3}O_{5}$: C, 55.08; H, 4.95; N, 13.76. Found: C, 55.63; H, 4.99' N, 12.98.

Preparation 149

5

To a slurry of 10% palladium on carbon (2.26 g, 20 mmol) in tetrahydrofuran was added a solution of the product of Preparation 148 (1.21 g, 3.96 mmol) in tetrahydrofuran (20 mL). The mixture was reacted under a hydrogen atmosphere (40 psi) on a Parr apparatus for 3 h and 10 subsequently filtered through celite. To this solution was added of the product of Preparation 1d (1.5 g, 3.96 mmol), 1-hydroxybenzotriazole (0.59 g, 4.35 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.9 g, 4.35 mmol). After 16 h, the reaction mixture was filtered and 15 concentrated. The resulting residue was purified by flash chromatography (silica gel, chloroform to 1% methanol/chloroform gradient) to yield 2.24 g (89%) of the desired product: 1H-NMR is consistent with structure; MS (ion spray) 638.4 (M+1); Anal. Calc'd for $C_{33}H_{43}N_5O_8$: C, 20 62.15; H, 6.80; N, 10.98. Found: C, 61.47; H, 6.41; N, 11.09.

Preparation 150

25

To a solution of the product of Preparation 149 (2.19 g, 3.4 mmol) stirring in dioxane (50 mL) at room temperature

254

was added a solution of lithium hydroxide (0.1 g, 4.08 mmol) in water (35 mL). After 15 min, the reaction was acidified to pH = 3.0 with 1 N hydrochloric acid and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated to yield 2.0 g (95%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (ion spray) 610 (M+1); Anal. Calc'd for C₃₁H₃₉N₅O₈·1.2dioxane: C, 60.10; H, 6.85; N, 9.79. Found: C, 59.78; H, 6.58; N, 10.14.

Preparation 151

To a solution of the product of Preparation 150 (0.6 g, 1.0 mmol) was added of 4-methylpiperidine (0.12 mL, 1.0 15 mmol), 1-hydroxybenzotriazole (0.15 g,1.1 mmol) and 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol). After 16 h, the reaction mixture was concentrated, slurried in ethyl acetate and filtered. Water was added and the mixture was extracted with ethyl 20 acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and The residue was purified by flash concentrated. chromatography (silica gel, 4% methanol/chloroform) to yield 0.46 g (67%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (ion spray) 691.3 (M+1); Anal. Calc'd for $C_{37}H_{50}N_6O_7$: C, 64.33; H, 7.29; N, 12.16. Found: C, 64.07; H, 7.29; N, 12.34.

WO 99/08697

255

PCT/US98/17201

To a solution of the product of Preparation 151 (0.37 g, 0.53 mmol) stirring in dichloromethane (6 mL) at room temperature was added trifluoroacetic acid (2 mL). After 1 h, the reaction mixture was poured into a solution of saturated sodium bicarbonate. The mixture was extracted with chloroform. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated. The residue was dissolved in ethyl acetate and this solution was acidified with a saturated 10 solution of hydrochloric acid in ether. The resulting slurry was concentrated to yield 0.3 g (86%) of the desired product: 1H-NMR is consistent with structure; MS (ion spray) 591.6 (M+1); Anal. Calc'd for $C_{32}H_{42}N_6O_5$ 2.2HCl: C, 57.29; H, 6.64; N, 12.53. Found: C, 57.18; H, 6.54; 15 N, 12.23. 0.14 g (0.2 mmol) of the free base was sent for chiral separation. Resolution of the diastereomers by HPLC gave two products:

Example 63. (Isomer 1). To the solution of the purified isomer in ethyl acetate was added a saturated solution of hydrochloric acid in ether. The resulting slurry was concentrated to yield 0.04 g (28%) of the desired isomer as a white solid: ¹H-NMR is consistent with structure; t_R = 6.92 min; MS (high res) Calc'd for C₃₂H₄₃N₆O₅: 591.3295.

25 Found: 591.3299. Anal. Calc'd for C₃₂H₄₃N₆O₅ 2.2HCl: C, 57.29; H, 6.64; N, 12.53. Found: C, 57.29; H, 6.25; N, 12.37.

Example 64. (Isomer 2). To the solution of the purified isomer in ethyl acetate was added a saturated solution of hydrochloric acid in ether. The resulting slurry was concentrated to yield 0.03 g (21%) of the desired isomer

256

as a tan foam: $^{1}\text{H-NMR}$ is consistent with structure; t_R = 9.64 min; MS (high res) Calc'd for $C_{32}H_{43}N_6O_5$: 591.3295. Found: 591.3288.

5

Preparation 152

To a solution of the product of Preparation 150 (0.6 g, 1.0 mmol), pyrrolidine (0.8 mL, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), in dimethylformamide (20 mL) as described in Preparation 150 gave 0.42 g (64%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (ion spray) 663.4 (M+1); Anal. Calc'd for C₃₅H₄₆N₆O₇: C, 63.43; H, 7.00; N, 12.68. Found: C, 63.39; H, 6.97; N, 12.58.

Example 65

To a solution of the product of Preparation 152 (0.35 g, 0.53 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as in described in Example 63 gave 0.23 g (68%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (ion spray) 563.5 (M+1); Anal. Calc'd for C₃₀H₃₈N₆O₅ 2.3HCl: C, 55.73; H, 6.28; N, 13.00. Found: C, 55.97; H, 6.18; N, 12.87.

257

Pr paration 153



To a solution of 2-anisaldehyde (13.5 g, 100 mmol), benzyltriethylammonium chloride (1.23 g, 5 mmol), chloroform (16 mL) and 50% sodium hydroxide (25 mL) as described Preparation 145 gave 15.0 g of the desired product as an oil which was carried without further purification.

10

Preparation 154



To a solution of the product of Preparation (15.04 g, 83 mmol), concentrated sulfuric acid (10 mL) and absolute ethanol (100 mL) as described in Preparation 146 gave 8.1 g (38% over the two steps) of the desired product as an oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 210 (M+).

Preparation 155



20

25

15

To a solution of the product of Preparation (4.0 g, 19 mmol), phosphorus tribromide (2.05 mL, 20 mmol) and chloroform (50 mL) as described in Preparation 147 gave 4.9 g (95%) of the desired product: ¹H-NMR is consistent with structure; MS (FD) 272, 274 (M+).

Preparation 156

258

Reaction of the product of Preparation 155 (5.18 g, 19.0 mmol), 4-nitroimidazole (2.37 g, 20.9 mmol) and sodium hydride (0.84 g, 20.9 mmol) in dimethylformamide (50 mL) as described in Preparation 148 gave 5.8 g (100%) of the desired product as a colorless oil which solidifies upon standing: ¹H-NMR is consistent with structure; MS (FD) 305 (M+); Anal. Calc'd for C₁₄H₁₅N₃O₅: C, 55.08; H, 4.95; N, 13.76. Found: C, 54.87; H, 4.96; N, 13.47.

10

15

20

5

Preparation 157

Reduction of the product of Preparation 156 (1.8 g, 5.8 mmol), 10% palladium on carbon (0.9 g) in tetrahydrofuran (80 mL) followed by coupling with the product of Preparation 1d (2.2 g, 5.8 mmol), 1-hydroxybenzotriazole (0.86 g, 6.4 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.3 g, 6.4 mmol), as described in Preparation 149 gave 2.0 g (54%) of the desired compound as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 637 (M+); Anal. Calc'd for C₃₃H₄₃N₅O₈·0.2H2O: C, 61.80; H, 6.82; N, 10.92. Found: C, 61.65; H, 6.93; N, 11.12.

Preparation 158

259

Reaction of the product of Preparation 157 (1.95 g, 3.0 mmol) and lithium hydroxide (0.09 g, 3.6 mmol) in dioxane (40 mL) and water (20 mL) as described in Preparation 150 gave 1.72 g (94%) of the desired product as a white foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 610 (M+).

Preparation 159

Reaction of the product of Preparation 158 (0.5 g, 0.82 mmol), 4-methylpiperidine (0.1 mL, 0.82 mmol), 1-hydroxybenzotriazole (0.12 g, 0.9 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.19 g, 0.9 mmol) in dimethylformamide (40 mL) as described in Preparation 151 gave 0.45 g (80%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 690 (M+); Anal. Calc'd for C₃₇H₅₀N₆O₇: C, 64.33; H, 7.30; N, 12.16. Found: C, 64.32; H, 7.21; N, 11.97.

20 Example 6

25

Reaction of the product of Preparation 159 (0.4 g, 0.58 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 63 gave 0.3 g (79%) of the desired product as a white solid: ¹H-NMR is consistent

260

with structure; MS (high res) Calc'd for $C_{32}H_{43}N_6O_5$: 591.3295. Found: 591.3298. Anal. Calc'd for $C_{32}H_{42}N_6O_5$ 2HCl: C, 57.92; H, 6.68; N, 12.66. Found: C, 57.27; H, 6.24; N, 11.82.

5

Preparation 160

Reaction of the product of Preparation 158 (0.5 g, 0.82 mmol), pyrrolidine (0.07 mL), 1-hydroxybenzotriazole


(0.12 g, 0.9 mmol) and 1-(3-dimethylaminopropyl)-3ethylcarbodiimide (0.19 g, 0.9 mmol) in dimethylformamide
(40 mL as described in Preparation 151 gave 0.35 g (65%)
of the desired product as a tan foam: ¹H-NMR is
consistent with structure; MS (FD) 662 (M+); Anal. Calc'd
for C₃₅H₄₆N₆O₇: C, 63.43; H, 7.00; N, 12.68. Found: C,
63.26; H, 6.94; N, 12.43.

Example 67

Reaction of the product of Preparation 160 (0.3 g, 0.4 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 63 gave 0.24 g (96%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (high res) Calc'd for C₃₀H₃₉N₆O₅: 563.2982. Found: 563.2989. Anal. Calc'd for C₃₀H₃₀N₆O₅ 2.4HCl: C, 55.42; H, 6.26; N, 12.93. Found: C, 55.51; H, 6.10; N, 12.30.

261


Preparation 161



Reaction of the product of Preparation 158 (0.5 g, 0.82 mmol), 4-(4-fluorobenzoyl)piperidine hydrochloride (0.2 g, 0.82 mmol), triethylamine (0.13 mL, 0.9 mmol), 1-hydroxybenzotriazole (0.12 g, 0.9 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.19 g, 0.9 mmol) in dimethylformamide (40 mL) as described in Preparation 151 gave 0.41 g (62%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (FD) 799 (M+); Anal. Calc'd for C₄₃H₅₁FN₆O₈: C, 64.65; H, 6.43; N, 10.52. Found: C, 64.44; H, 6.56; N, 10.53.

15

Example 68



Reaction of the product of Preparation 161 (0.36 g, 0.45 mmol) and trifluoroacetic acid (2 mL) in dichloromethane (6 mL) as described in Example 63 gave 0.26 g (74%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 699 (M+); Anal. Calc'd for C₃₈H₄₃FN₆O₆ 2HCl: C, 59.14; H, 5.88; N, 10.89. Found: C, 59.36; H, 5.99; N, 10.80.

25

20

Preparation 162

262

Reaction of 3-tolylbenzaldehyde (12 g, 100 mmol), benzyltriethylammonium chloride (1.23 g, 5 mmol), chloroform (16 mL) and 50% sodium hydroxide (25 mL) as described in Preparation 145 gave 16.2 g of an oil which was used without further purification.

Preparation 163

10 Reaction of the product of Preparation 162 (16.2 g, 98 mmol), conc. sulfuric acid (10 mL) and absolute ethanol (100 mL) as described in Preparation 146 gave 10.8 g (52% over the two steps) of the desired product as an oil: ¹H-NMR is consistent with structure; MS (FD) 194 (M+).

15

Preparation 164

Reaction of the product of Preparation 163 (4.85 g, 25 mmol), phosphorus tribromide (2.65 mL, 27.5 mmol) and chloroform (50 mL) as described in Preparation 147 gave 3.71 g (58%) of the desired product as an oil: 1H-NMR is consistent with structure; MS (FD) 256 (M+).

Preparation 164



25

Reaction of the product of Preparation 164 (3.7 g, 14.4 mmol), 4-nitroimidazole (1.8 g, 15.9 mmol) and sodium

263

hydride (0.64 g, 15.9 mmol) in dimethylformamide (30 mL) as described in Preparation 148 gave 3.2 g (77%) of the desired product as an oil: ¹H-NMR is consistent with structure; MS (FD) 289 (M+); Anal. Calc'd for C₁₄H₁₅N₃O₄: C, 58.13; H, 5.23; N, 14.52. Found: C, 58.18; H, 5.53; N, 14.89.

Preparation 165

10 Reduction of the product of Preparation 164 (1.15 g, 4.0 mmol), 5% palladium on carbon (0.57 g) in tetrahydrofuran (30 mL) followed by coupling with the product of Preparation 1d (1.5 g, 4.0 mmol), 1-hydroxybenzotriazole (0.6 g, 4.4 mmol) and 1-(3-dimethylaminopropyl)-3-

ethylcarbodiimide (0.9 g, 4.4 mmol) as described in Preparation 149 gave 2.24 g (90%) of the desired product: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 621 (M+); Anal. Calc'd for $C_{33}H_{43}N_{5}O_{7}$: C, 63.75: H, 6.97; N, 11.26. Found: C, 63.69; H, 7.06; N, 11.27.

20

Preparation 166

Reaction of the product of Preparation 165 (2.0 g, 3.3 mmol) with lithium hydroxide (0.088 g, 3.64 mmol) in

25 dioxane (20 mL) and water (10 mL) as described in


Preparation 150 gave 1.95 g (100%) of the desired

product: ¹H-NMR is consistent with structure; MS (FD) 594

264

(M+); Anal. Calc'd for $C_{31}H_{39}N_5O_7$ 1 H_2O : C, 61.59; H, 6.93; N, 10.38. Found: C, 61.26; H, 6.86; N, 10.77.

Preparation 167




5

Reaction of the product of Preparation 166 (0.6 g, 1.0 mmol), (0.7 g, 4-(4-fluorobenzoyl)piperidine hydrochloride (0.24 g, 1.0 mmol), triethylamine (0.1 g, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), in dimethylformamide (13 mL) as described in Preparation 151 gave 0.7 g (90%) of the desired product:

1H-NMR is consistent with structure; MS (FD) 782 (M+); Anal. Calc'd for C43H51FN6O7: C, 65.97; H, 6.57; N, 10.73.

15 Found: C, 66.01; H, 6.54; N, 10.52.

Examples 69 and 70



Reaction of the product of Preparation 167 (0.47 g, 0.6 mmol) with trifluoroacetic acid (4 mL) in dichloromethane (12 mL) as described Example 63 gave 0.44 g (98%) of the desired product as a foam: ¹H-NMR is consistent with structure; MS (ion spray) 682.2 (M+1); Anal. Calc'd for C₃₈H₄₃FN₆O₅·2.55HCl: C, 59.22; H, 5.94; N, 10.90. Found: C, 58.84; H, 5.94; N, 10.73. Resolution of the diastereomers (0.25 g, 0.37 mmol) by chiral HPLC gave the individual isomers:

265

Example 69. (Isom r 1) 0.069 g (27%) of the desired isomer as the free base. $^{1}\text{H-NMR}$ is consistent with structure; $t_R = 5.2$ min; MS (ion spray) 683.4 (M+1). Example 70. (Isomer 2) 0.065 g (26%) of the desired isomer as the free base. $^{1}\text{H-NMR}$ is consistent with structure; $t_R = 6.5$ min; MS (ion spray) 683.4 (M+1); Anal. Calc'd for $C_{38}H_{43}FN_{6}O_{5}$ 0.1 $H_{2}O$: C, 63.48; H, 6.17; N, 11.68. Found: C, 63.07; H, 6.07; N, 11.39.

10 Preparation 168

Reaction of the product of Preparation 166 (0.59 g, 1.0 mmol), 4-methylpiperidine (0.099 g, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol) in dimethylformamide (13 mL) as described in Preparation 151 gave 0.67 g (100%) of the desired product as a foam: ¹H-NMR is consistent with structure; MS (ion spray) 675 (M+1); Anal. Calc'd for C₂₇H₅₀N₆O₆: C, 65.85; H, 7.47; N, 12.45. Found: C, 66.09; H, 7.23; N, 12.53.

Examples 71 and 72

Reaction of the product of Preparation 168 (0.51 g, 0.76 mmol) with trifluoroacetic acid (4 mL) in dichloromethane (12 mL) as described in Example 63 gave 0.42 g (86%) of the desired product: ¹H-NMR is consistent with structure; MS (ion spray) 574.1 (M+1); Anal. Calc'd for

266

 $C_{32}H_{42}N_6O_4$ 2.25HCl: C, 58.52; H, 6.79; N, 12.80. Found: C, 58.54; H, 6.68; N, 12.80. Resolution of the diastereomers (0.173 g, 0.32 mmol) by chiral HPLC gave the individual isomers which were individually treated with a saturated solution of hydrochloric acid in diethyl ether to give the desired products:

Example 71. (Isomer 1) 1 H-NMR is consistent with structure; $t_{R} = 6.22$ min; MS (ion spray) 575.4 (M+1); Anal. Calc'd for $C_{32}H_{42}N_{6}O_{4}$ 2.3HCl: C, 58.36; H, 6.78; N, 12.76. Found: C, 58.23; H, 6.57; N, 12.53.

Example 72. (Isomer 2) 1 H-NMR is consistent with structure; $t_{R} = 8.53$ min; MS (ion spray) 575.4 (M+1); Anal. Calc'd for $C_{32}H_{42}N_{6}O_{4}$ 2.55HCl: C, 57.56; H, 6.73; N, 12.59. Found: C, 57.95; H, 6.91; N, 12.29.

15

10

Preparation 169

Reaction of the product of Preparation 166 (0.59 g, 1.0 mmol), pyrrolidine (0.071 g, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol) in dimethylformamide (13 mL) as described in Preparation 151 gave 0.57 g (88%) of the desired product:

1H-NMR is consistent with structure; MS (ion spray) 646 (M+1); Anal. Calc'd for C35H46N6O6: C, 65.00; H, 7.17; N, 12.99. Found: C, 64.95; H, 6.98; N, 13.19.

Example 73

267

Reaction of the product of Preparation 170 (0.55 g, 0.85 mmol) and trifluoroacetic acid (4 mL) in dichloromethane (12 mL) as described in Example 63 gave 0.37 g (70%) of the desired product: ¹H-NMR is consistent with structure; MS (ion spray) 546.1 (M+1); Anal. Calc'd for C₃₀H₃₈N₆O₄ 2.4HCl: C, 56.79; H, 6.42; N, 13.24. Found: C, 56.81; H, 6.35; N, 13.10.

10 Preparation 170



Reaction of 2-methylbenzaldehyde (12.0 g, 100 mmol), triethylbenzylammonium chloride (1.23 g, 5 mmol), chloroform (16 mL) and 50% sodium hydroxide (25 mL) as described in Preparation 145 gave 16.0 of an oil which was used without further purification.

Preparation 171

20 Reaction of the product of Preparation 170 (16.0 g, 96 mmol), concentrated sulfuric acid (10 mL) and absolute ethanol (100 mL) as described in Preparation 146 gave 15.2 g (78% over the two steps) of the desired product:

1H-NMR is consistent with structure; MS (ion spray) 194
25 (M+1).

Preparation 172

268


Reaction of the product of Preparation 171 (3.88 g, 20 mmol), phosphorus tribromide (2.2 mL, 22 mmol), chloroform (50 mL) as described in Preparation 147 gave 2.87 g (65%) of the desired product as a oil: ¹H-NMR is consistent with structure; MS (ion spray) 256, 258 (M+).

Preparation 173



Reaction of the product of Preparation 172 (3.3 g, 12.8 mmol), 4-nitroimidazole (1.53 g, 13.5 mmol) and sodium hydride (0.32 g, 13.5 mmol) in dimethylformamide (30 mL) as described in Preparation 148 gave 3.1 g (84%) of the desired product as an oil: ¹H-NMR is consistent with structure; MS (ion spray) 289 (M+1); Anal. Calc'd for C₁₄H₁₅N₃O₄: C, 58.13; H, 5.23; N, 14.52. Found: C, 58.41; H, 5.26; N, 14.47.

Preparation 174



20

Reduction of the product of Preparation 173 (1.15 g, 3.96 mmol) under a hydrogen atmosphere with 5% palladium on carbon (0.59 g) in tetrahydrofuran (30 mL) followed by coupling with the product of Preparation 1d (1.5 g, 3.96 mmol), 1-hydroxybenzotriazole (0.59 g, 4.35 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.9 g, 4.35

269

mmol) as described in Preparation 149 gave 2.43 g (99%) of the desired product as a foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 621.2 (M+1); Anal. Calc'd for $C_{33}H_{43}N_5O_7$: C, 63.75; H, 6.97; N, 11.26. Found: C, 63.59; H, 7.12; N, 11.38.

Preparation 175

Reaction of the product of Preparation 174 (2.2 g, 3.6 mmol) with lithium hydroxide (0.096 g, 4.0 mmol) in dioxane (20 mL) and water (10 mL) as described in Preparation 150 gave 2.1 g (100%) of the desired product:

1H-NMR is consistent with structure; MS (ion spray) 594 (M+1); Anal. Calc'd for C₃₁H₃₉N₅O₇·0.4H₂0: C, 61.27; H, 6.91; N, 10.57. Found: C, 60.93; H, 6.57; N, 10.92.

Preparation 176

Reaction of the product of Preparation 175 (0.59 g, 1.0 mmol), 4-methylpiperidine (0.099 g, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol) in dimethylformamide (10 mL) as described in Preparation 151 gave the desired product as an oil: ¹H-NMR is consistent with structure; MS (ion spray) 674.3 (M+1); Anal. Calc'd for C₃₇H₅₀N₆O₆: C, 65.85; H, 7.47; N, 12.45. Found: C, 65.56; H, 7.53; N, 12.69.

270

Example 74

Reaction of the product of Preparation 176 (0.49 g, 0.72 mmol) with trifluoroacetic acid (4 mL) in dichloromethane (12 mL) as described in Example 63 gave 0.31 g (75%) of the desired product: ¹H-NMR is consistent with structure; MS (ion spray) 574 (M+1); Anal. Calc'd for C₃₂H₄₂N₆O₄·2HCl: C, 59.35; H, 6.85; N, 12.98. Found: C, 59.27; H, 6.76; N, 13.02.

Preparation 177

271

Examples 75 and 76

Reaction of the product of Preparation 177 (0.40 g, 0.62 mmol) with trifluoroacetic acid (4 mL) in dichloromethane (12 mL) as described in Example 63 gave 0.36 g (95%) of the desired product: ¹H-NMR is consistent with structure; MS (ion spray) 546 (M+1); Anal. Calc'd for C₃₀H₃₈N₆O₄ 2HCl: C, 58.16; H, 6.51; N, 13.56. Found: C, 58.09; H, 6.43; N, 13.60. Resolution of the diastereomers (0.17 g, 0.31 mmol) by chiral HPLC gave the respective isomers which were individually treated with a saturated solution of hydrochloric acid in diethyl ether to give the desired products:

Example 75. (Isomer 1) 0.039 g (20%), ¹H-NMR is

consistent with structure; t_R = 6.50 min; MS (ion spray)
547.2 (M+1); Anal. Calc'd for C₃₀H₃₈N₆O₄ 2.3HCl: C, 57.15;
H, 6.44; N, 13.33. Found: C, 57.15; H, 6.17; N, 12.94.

Example 76. (Isomer 2) 0.054 g (28%) of the desired isomer as the hydrochloric acid salt. ¹H-NMR is

consistent with structure; t_R = 7.52 min; MS (ion spray)
547.2 (M+1); Anal. Calc'd for C₃₀H₃₈N₆O₄ 2.75HCl: C, 55.70;
H, 6.35; N, 12.99. Found: C, 55.68; H, 6.35; N, 12.66.

Preparation 178

25

Reaction of the product of Preparation 175 (0.59 g, 1.0 mmol), 4-(4-fluorobenzoyl)piperidine hydrochloride (0.24

272

g, 1.0 mmol), triethylamine (0.11 g, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol) and 1-(3- dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol) in dimethylformamide (13 mL) as described in Preparation 151 gave 0.58 g (82%) of the desired product as a product: $^1\text{H-NMR}$ is consistent with structure; MS (ion spray) 783 (M+1); Anal. Calc'd for $C_{43}\text{H}_{51}\text{FN}_5\text{O}_7$: C, 65.97; H, 6.57; N, 10.73. Found: C, 65.70; H, 6.69; N, 10.47.

10

Example 77

Reaction of the product of Preparation 178 (0.58 g, 0.74 mmol) with trifluoroacetic acid (4 mL) in dichloromethane (12 mL) as described in Example 63 gave 0.52 g (93%) of the desired product: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 682 (M+1); Anal. Calc'd for $C_{38}H_{43}FN_6O_5$ 2HCl: C, 60.40; H, 6.00; N, 11.12; Cl, 9.38. Found: C, 60.35; H, 5.96; N, 11.17; Cl, 9.23.

20

25

15

Preparation 179



To a solution of 4-methylphenylacetic acid, (7.5 g, 50 mmol) in carbon tetrachloride (10 mL) was added thionyl chloride (14.4 mL, 200 mmol). The reaction was heated to reflux. After 30 min, the mixture was cooled to 20°C and a solution of N-bromosuccinimde (8.9 g, 50 mmol) and HBr (8 drops of a 48% aqueous solution) in carbon tetrachloride (15 mL). The reaction was heated to reflux

273

and after 30 min, cooled to ambient temperature, filtered and concentrated. The resulting oil was added to absolute ethanol at 0°C and then concentrated. The residue was purified by flash chromatography (silica gel, 3% ethyl acetate/hexanes) to yield 5.1 g (40%) of the desired product as an oil: 1H-NMR is consistent with structure; MS (FD) 258 (M+); Anal. Calc'd for C₁₁H₁₃BrO₂·1.3CHCl₃: C, 44.44; H, 4.34. Found: C, 44.58; H, 4.51.

10

20

Preparation 180



Reaction of the product of Preparation 179 (5.0 g, 19.5 mmol), 4-nitroimidazole (2.2 g, 19.5 mmol) and sodium 15 hydride (0.47 g, 19.5 mmol) indimethylformamide (50 mL) as described in Preparation 148 gave 1.9 g (33%) of the desired product as a yellow oil: 1H-NMR is consistent with structure; MS (FD) 289 (M+); Anal. Calc'd for $C_{14}H_{15}N_3O_4$: C, 58.13; H, 5.23; N, 14.52. Found: C, 58.33; H, 5.17; N, 14.70.

Reduction of the product of Preparation 180 (4.9 g, 17.0 25 mmol) under a hydrogen atmosphere with 10% palladium on carbon (3.5 g) in tetrahydrofuran (120 mL) followed by coupling of the product of Preparation 1d (6.43 g, 17.0 mmol), 1-hydroxybenzotriazole (2.6 g, 18.7 mmol), 1-(3Found: C, 63.80; H, 7.09; N, 11.36.

WO 99/08697 PCT/US98/17201

274

dimethylaminopropyl)-3-ethylcarbodiimide (3.9 g, 18.7 mmol) as described in Preparation 149 gave 6.54 g (62%) of the desired compound as an orange foam: 1H -NMR is consistent with structure; MS (ion spray) 622.5 (M+1); Anal. Calc'd for $C_{33}H_{43}N_5O_7$: C, 63.75; H, 6.97; N, 11.26.

Preparation 182

Reaction of the product of Preparation 181 (6.49 g, 10.0 mmol) with lithium hydroxide (0.3 g, 12.0 mmol) in dioxane (90 mL) and water (50 mL) as described in Preparation 150 gave 5.93 g (100%) of the desired product: ¹H-NMR is consistent with structure; MS (ion spray) 594.6 (M+1);

Anal. Calc'd for C₃₁H₃₉N₅O₇ 0.4 dioxane: C, 62.26; H, 6.76; N, 11.14. Found: C, 62.33; H, 6.41; N, 11.19.

Preparation 183

(ion spray) 675.7 (M+1); Anal. Calc'd for $C_{37}H_{50}N_6O_6$ 0.3 H_2O :

20 Reaction of the product of Preparation 182 (5.0 g, 8.4 mmol), 4-methylpiperidine (1.0 mL, 8.4 mmol), 1-hydroxybenzotriazole (1.3 g, 9.24 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.9 g, 9.24 mmol) in dimethylformamide (80 mL) as described in
25 Preparation 151 gave 4.44 g (78%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS

275

C, 65.33; H, 7.50; N, 12.35. Found: C, 65.28; H, 7.37; N, 12.30.

Examples 78 and 79

5

Reaction of the product of Preparation 183 (4.17 g, 6.4 mmol) and trifluoroacetic acid (20 mL) in dichloromethane (50 mL) as described in Example 63 gave 2.59 g (65%) of the desired product: ¹H-NMR is consistent with structure;

10 MS (ion spray) 575.5 (M+1). Resolution of the diastereomers (0.22 g, 0.34 mmol) by chiral HPLC gave the respective isomers which were individually treated with a saturated solution of hydrochloric acid in diethyl ether to give the desired products:

Example 78. (Isomer 1): 0.091 g (36%), ¹H-NMR is consistent with structure; t_R = 4.40 min; MS (ion spray) 575.3 (M+1); Anal. Calc'd for C₃₂H₄₂N₆O₄: C, 66.88; H, 7.37; N, 14.62. Found: C, 66.30; H, 7.20; N, 14.40. Example 79. (Isomer 2): 0.059 g (24%); ¹H-NMR is consistent with structure; t_R = 5.3 min; MS (ion spray) 575.3 (M+1); Anal. Calc'd for C₃₂H₄₂N₆O₄: C, 66.88; H, 7.37; N, 14.62. Found: C, 67.02; H, 7.34; N, 14.40.

Preparation 184

276

Reaction of the product of Preparation 182 (0.7 g, 1.2 mmol), dimethylamine hydrochloride (0.1 g, 1.2 mmol), triethylamine (0.2 mL, 1.32 mmol), 1-hydroxybenzotriazole (0.18 g, 1.32 mmol) and 1-(3-dimethylaminopropyl)-3
5 ethylcarbodiimide (0.27 g, 1.32 mmol) in dimethylformamide (30 mL) as described in Preparation 151-gave 0.55 g (75%) of the desired product as a tan foam:

1H-NMR is consistent with structure; MS (ion spray) 621.7 (M+1); Anal. Calc'd for C33H44N6O6: C, 63.85; H, 7.15; N,

10 13.54. Found: C, 63.56; H, 7.37; N, 13.35.

Example 80

Reaction of the product of Preparation 184 (0.46 g, 0.74 mmol) with trifluoroacetic acid (2 mL)in dichloromethane (6 mL) as described in Example 63 gave 0.22 g (50%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (ion spray) 521.4 (M+1); Anal. Calc'd for C₂₈H₃₆N₆O₄ 2.5HCl: C, 54.97; H, 6.34; N, 13.74. Found: C, 54.85; H, 6.23; N, 13.58.

Preparation 185

Reaction of the product of Preparation 182 (0.59 g, 1 mmol), pyrrolidine (0.078 g, 1.1 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.27 g, 1.1

277

mmol) in dimethylformamide (12 mL) as described in Preparation 151 gave 0.52 g (80%) of the desired product: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 646 (M+1); Anal. Calc'd for $\text{C}_{35}\text{H}_{46}\text{N}_{6}\text{O}_{6}$: C, 65.00; H, 7.17; N, 12.99. Found: C, 65.00; H, 7.05; N, 12.82.

5

Example 81

Reaction of the product of Preparation 185 (0.39 g, 0.61 mmol) with trifluoroacetic acid (3.5 mL) indichloromethane (12 mL) as described in Example 63 gave 0.3 g (90%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (ion spray) 546 (M+1); Anal. Calc'd for C₃₀H₃₆N₆O₄ 1.4HCl: C, 60.99; H, 6.69; N, 14.22. Found: C, 61.08; H, 6.51; N, 13.89.

Preparation 186

Reaction of the product of Preparation 182 (0.7 g, 1,32 mmol), 4-(4-fluorobenzoyl)piperidine hydrochloride (0.3 g, 1.2 mmol), triethylamine (0.2 mL, 1.32 mmol), 1-hydroxybenzotriazole (0.18 g, 1.32 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.27 g, 1.32 mmol) in dimethylformamide (30 mL) as described in Preparation 151 gave 0.68 g (72%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (ion spray) 783.6 (M+1); Anal. Calc'd for C₄₃H₅₁FN₆O₇: C,

278

65.97; H, 6.57; N, 10.73. Found: C, 65.86; H, 6.62; N, 10.62.

Examples 82 and 83

5

Reaction of the product of Preparation 186 (0.7 g, 0.89 mmol) with trifluoroacetic acid (2 mL) in dichloromethane (6 mL), as in Example 63 from Examples Part 2A. (66%) of the desired product as a yellow solid: 1H-NMR is consistent with structure; MS (ion spray) 683.4 (M+1); 10 Anal. Calc'd for $C_{38}H_{43}N_6O_5$ 2.4HCl: C, 59.25; H, 5.94; N, 10.91. Found: C, 59.24; H, 5.66; N, 11.09. Resolution of the diastereomers (0.21 g, 0.32 mmol) by chiral HPLC gave the respective isomers which were individually treated with a saturated solution of hydrochloric acid in diethyl 15 ether to give the desired products: **Example 82.** (Isomer 1) $0.092 \text{ g } (38\%); ^{1}\text{H-NMR is}$ consistent with structure; $t_R = 5.83 \text{ min}$; MS (ion spray) 683.4 (M+1); Anal. Calc'd for C₃₈H₄₃FN₆O₅ 2HCl: C, 60.40; H, 20 6.00; N, 11.12. Found: C, 60.11; H, 6.12; N, 10.98. Example 83 (Isomer 2) 0.065 g (27%) of the desired isomer as a white solid: ¹H-NMR is consistent with structure; $t_R = 7.62 \text{ min}$; MS (ion spray) 683.4 (M+1); Anal. Calc'd for C38H43FN6O5 2HCl: C, 60.40; H, 6.00; N, 25 11.12. Found: C, 60.15; H, 5.82; N, 10.96.

Preparation 187

279


Reaction of 1-naphthylacetic acid (9.3 g, 50 mmol), thionyl chloride (14.4 mL, 200 mmol), carbon tetrachloride (35 mL), N-bromosuccinimide (8.9 g, 50 mmol), 48% HBr (8 drops) as described in Preparation 179 gave 12.6 g (86%) of the desired product as an oil: ¹H-NMR is consistent with structure.

Preparation 188



Reaction of the product of Preparation 187 (11.8 g, 40 mmol), 4-nitroimidazole (4.5 g, 40 mmol) and sodium hydride (1.6 g, 40 mmol) in dimethylformamide (50 mL) as described in Preparation 148 gave 6.03 g (50%) of the desired product as an oil: ¹H-NMR is consistent with structure; MS (ion spray) 325.1 (M+1); Anal. Calc'd for C₁₇H₁₅N₃O₄·0.37H₂O: C, 61.50; H, 4.78; N, 12.66. Found: C, 61.46; H, 4.60; N, 12.73.

Preparation 189



20

25

Reduction of the product of Preparation 188 (1.28 g, 4.0 mmol) under a hydrogen atmosphere with 5% palladium on carbon (0.6 g) followed by coupling with the product of Preparation 1d (1.5 g, 4.0 mmol), 1-hydroxybenzotriazole (0.59 g, 4.35 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.9 g, 4.35 mmol) in tetrahydrofuran (30 mL) as described in Preparation 149 gave 1.99 g (77%) of the desired product as an orange foam: ¹H-NMR is consistent with structure; MS (ion spray) 657 (M+1);

280

Anal. Calc'd for $C_{36}H_{43}N_{5}O_{7}$: C, 65.74; H, 6.59; N, 10.65. Found: C, 65.67; H, 6.53; N, 10.87.


Preparation 190

5

10

The product of Preparation 189 (1.97 g, 3.0 mmol), lithium hydroxide (0.08 g, 3.3 mmol), dioxane (20 mL), water (10 mL), as in Preparation 150. 1.8 g (95%) of the desired product. $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 630 (M+1); Anal. Calc'd for $C_{34}H_{39}N_{5}O_{7}$ ·1.05 $H_{2}O$: C, 62.96; H, 6.39; N, 10.80. Found: C, 63.09; H, 6.39; N, 10.40.

Preparation 191



15

Reaction of the product of Preparation 190 (0.63 g, 1.0 mmol), 4-methylpiperidine (0.099 g, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol) in dimethylformamide (12 mL) as described in Preparation 151 gave 0.60 g (85%) of the desired material as a white solid: ¹H-NMR is consistent with structure; MS (ion spray) 710 (M+1); Anal. Calc'd for C40H50N6O6: C, 67.58; H, 7.09; N, 11.82. Found: C, 67.33; H, 6.94; N, 11.58.

Example 84

281

Reaction of the product of Preparation 191 (0.60 g, 0.84 mmol) with trifluoroacetic acid (4 mL) in dichloromethane (12 mL) as described in Example 63 gave 0.47 g (92%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (ion spray) 610 (M+1); Anal. Calc'd for C₃₅H₄₂N₆O₄·2.7HCl: C, 59.28; H, 6.35; N, 11.85. Found: C, 59.34; H, 6.57; N, 11.75.

10 Preparation 19

5

Reaction of the product of Preparation 190 (0.63 g, 1.0 mmol), pyrrolidine (0.071 g, 1.0 mmol), 1-hydroxybenzotriazole(0.15 g, 1.1 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol) in dimethylformamide (12 mL) as described in Preparation 151 gave 0.54 g (78%) of the desired product as a solid: ¹H-NMR is consistent with structure; MS (ion spray) 682 (M+1); Anal. Calc'd for C₃₈H₄₆N₆O₆: C, 66.84; H, 6.79; N, 12.31. Found: C, 66.59; H, 6.78; N, 12.29.

Example 85

Reaction of the product of Preparation 192 (0.443 g, 0.65 mmol) with trifluoroacetic acid (4 mL) in dichloromethane

(12 mL) as described in Example 63 gave 0.27 g (65%) of the desired product as a white solid: ^{1}H -NMR is consistent with structure; MS (ion spray) 582 (M+1); Anal. Calc'd for $C_{33}H_{38}N_6O_4$ 2.8HCl: C, 57.88; H, 6.01; N, 12.27. Found: C, 57.83; H, 6.47; N, 12.11.

Preparation 193

To a solution of Preparation 8 from Examples Part 1 (1.0 10 g, 1.7 mmol), N,N-dimethylamine hydrochloride, 0.14 g (1.7 mmol), triethylamine, 0.26 mL (1.9 mmol) and 1hydroxybenzotriazole, 0.26 g (1.9 mmol) in 70 mL of dimethylformamide was added 0.4 g (1.9 mmol) of 1-(3dimethylaminopropyl)-3-ethylcarbodiimide. The reaction 15 mixture was stirred overnight then concentrated. residue was slurried in ethyl acetate, filtered and water was added. The mixture was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. 20 The residue was chromatographed on silica gel using 4% methanol/chloroform as eluant to yield 0.58 g (56%) of the desired compound as a white foam: 1H-NMR is consistent with structure; MS (FD) 606 (M+); Anal. Calc'd for $C_{32}H_{42}N_6O_6$: C, 63.35; H, 6.98; N, 13.85. Found: C, 25 63.18; H, 7.03; N, 13.84.

Example 86

283

To a solution of the product of Preparation 193, 0.5 g (0.82 mmol) in 12 mL of dichloromethane was added 4 mL of trifluoroacetic acid. After stirred for 1 h, water was The reaction was quenched with solid sodium added. bicarbonate and was extracted with chloroform. combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. residue was dissolved in ethyl acetate and hydrochloric acid-saturated ether was added. The resulting slurry was 10 concentrated to yield 0.4 g (85%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 506.4 (M+); Anal. Calc'd for $C_{27}H_{34}N_6O_4$ 2.9HCl: C, 53.85; H, 4.50; N, 13.95. Found: C, 53.91; H, 6.14; N, 15 13.76.

Preparation 194

Reaction of the product of Preparation 8 from Examples

20 Part 1 (1.0 g, 1.7 mmol), diethylamine (0.18 mL, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (80 mL) as described in Preparation 193 gave 0.53 g (49%) of the desired product

25 as a yellow foam: 1H-NMR is consistent with structure; MS (FD) 634.3 (M+).

284

Example 87

Reaction of the product of Preparation 194 (0.52 g, 0.82 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.47 g (100%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 534.1 (M+); Anal. Calc'd for C₂₉H₃₆N₆O₄ 2.4HCl: C, 55.99; H, 6.54; N, 13.51. Found: C, 55.88; H, 6.91; N, 13.32.

10

Preparation 195

Reaction of the product of Preparation 8 from Examples
Part 1 (1.0 g, 1.7 mmol), N,N-methylethylamine (0.15 mL,
1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (40 mL) as in Preparation 193 gave 0.56 g (56%) of the desired product as a tan foam:

1H-NMR is consistent with structure; MS (FD) 620 (M+);
20 Anal. Calc'd for C₃₃H₄₄N₆O₆: C, 63.85; H, 7.15; N, 13.54.
Found: C, 63.45; H, 7.19; N, 13.15.

Example 88

285

(0.4 g, 0.64 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as in Example 86 gave 0.32 g (84%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 520 (M+); Anal. Calc'd for C₂₈H₃₆N₆O₄ 2.2HCl: C, 55.97; H, 6.41; N, 13.99. Found: C, 56.11; H, 6.23; N, 13.60.

10 Preparation 19

Reaction of the product of Preparation 8 from Examples
Part 1 (0.6 g, 1.0 mmol), cyclopropylamine (0.07 mL, 1.0
mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3
dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1
mmol), dimethylformamide (30 mL) as described in
Preparation 193 gave 0.31 g (50%) of the desired product
as a white foam: ¹H-NMR is consistent with structure; MS
(ion spray) 619.6 (M+1); Anal. Calc'd for C₃₃H₄₂N₆O₆ 1.1H₂O:

C, 62.07; H, 6.98; N, 13.15. Found: C, 62.19; H, 6.43;
N, 12.82.

Example 89

286

Reaction of the product of Preparation 196 (0.31 g, 0.5 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL) as described in example 86 gave 0.27 g (90%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (ion spray) 518 (M+1); Anal. Calc'd for C₂₈H₃₄N₆O₄ 2.4HCl: C, 55.49; H, 6.05; N, 13.87. Found: C, 55.63; H, 5.27; N, 13.29.

Preparation 197

10

15

Reaction of the product of Preparation 8 from Examples Part 1 (1.0 g, 1.7 mmol), benzylamine (0.2 mL, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.7 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (80 mL) as in Preparation 193 gave 0.86 g (75%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (FD) 669 (M+).

Example 90

20

25

Reaction of the product 197 (0.76 g, 1.1 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.52 g (79%) of the desired product as a yellow solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 568 (M+); Anal. Calc'd for $C_{32}H_{35}N_6O_4$ ·2.5HCl: C, 58.25; H, 5.88; N, 12.74. Found: C, 57.95; H, 6.02; N, 13.18.

Preparation 198

Reaction of the product of Preparation 8 from Examples

Part 1 (1.0 g, 1.7 mmol), N,N-benzylmethylamine (0.22 mL,
1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (80 mL) as in Preparation 193 gave 0.65 g (56%) of the desired product as a white foam:

10 H-NMR is consistent with structure; MS (FD) 682.5 (M+).

Example 91

Reaction of the product of Preparation 198 (0.6 g, 0.88 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.53 g (96%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 582.2 (M+); Anal. Calc'd for C₃₃H₃₇N₆O₄ 2.5HCl: C, 58.82; H, 6.08; N, 12.47. Found: C, 58.85; H, 6.27; N, 12.39.

Preparation 199

288

Reaction of the product of Preparation 8 from Examples Part 1 (1.0 g, 1.7 mmol), methoxypropylamine (0.18 mL, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (40 mL) as in Preparation 193 gave 0.8 g (73%) of the desired product as a white foam: 1H-NMR is consistent with structure; MS (FD) 650 (M+).

Example 92

10

15

Reaction of the product of Preparation 199 (0.75 g (1.16 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.52 g (72%) of the desired product as a yellow solid: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 550 (M+); Anal. Calc'd for $C_{29}H_{38}N_6O_5$ 2.7HCl: C, 53.66; H, 6.32; N, 12.95. Found: C, 53.93; H, 6.27; N, 13.14.

Preparation 200

20

25

Reaction of the product of Preparation 8 from Examples Part 1 (0.6 g, 1.0 mmol), 2-(ethylthio)ethylamine hydrochloride (0.15 g, 1.0 mmol), triethylamine (0.16 mL, 1.1 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide (30 mL) as in Preparation 193 gave 0.42 g (63%) of the desired product as a white foam:

1H-NMR is consistent with structure; MS (FD) 663.3 (M+);

289

Anal. Calc'd for $C_{34}H_{46}N_6O_6S$: C, 61.24; H, 6.95; N, 12.60. Found: C, 61.00; H, 6.83; N, 12.48.

Example 93

5

10

Reaction of the product of Preparation 200 (0.36 g, 0.54 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 86 gave 0.28 g (80%) of the desired product as a white solid: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 566 (M+); Anal. Calc'd for $C_{29}H_{38}N_6O_4S.2.2HCl:$ C, 53.84; H, 6.26; N, 12.99. Found: C, 53.99; H, 6.03; N, 12.79.

Preparation 201

15

20

25

Reaction of the product of Preparation 8 from Examples Part 1 (0.6 g, 1.0 mmol), phenoxyethylamine (0.14 g, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide (20 mL) as described in Preparation 193 gave 0.53 g (76%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 698 (M+).

Example 94

290

Reaction of the product of Preparation 201 (0.46 g, 0.67 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL) as described in Example 86 gave 0.28 g (62%) of the desired product as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 598 (M+); Anal. Calc'd for $C_{33}H_{38}N_{6}O_{5}.2.0HCl$: C, 59.01; H, 6.00; N, 12.51. Found: C, 58.97; H, 6.09; N, 12.40.

10 Preparation 202

Reaction of the product of Preparation 8 from Examples
Part 1 (0.6 g, 1.0 mmol), 4-fluorophenethylamine (0.13
mL, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol),
1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g,
1.1 mmol), dimethylformamide (30 mL) as described in
Preparation 193 gave 0.5 g (71%) of the desired product
as a yellow foam: ¹H-NMR is consistent with structure; MS
(FD) 700.5 (M+); Anal. Calc'd for C₃₈H₄₅FN₆O₆ 0.5H₂O: C,
20 64.30; H, 6.53; N, 11.84. Found: C, 64.12; H, 6.38; N,
11.73.

Example 95

291

Reaction of the product of Preparation 202 (0.5 g, 0.71 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 86 gave 0.28 g (58%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 600 (M+); Anal. Calc'd for C₃₃H₃₇FN₆O₄ 2.2HCl: C, 58.21; H, 5.80; N, 12.34. Found: C, 58.32; H, 5.92; N, 12.07.

Preparation 203

10

Reaction of the product of Preparation 8 from Examples
Part 1 (1.0 g, 1.7 mmol), 3-pyrroline (0.13 mL, 1.7
mmol), 1-hydroxybenzotriazole (0.25 g, 1.9 mmol), 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9
mmol), dimethylformamide (80 mL) as described in
Preparation 193 gave 0.75 g (70%) of the desired product
as a white foam: ¹H-NMR is consistent with structure; MS
(FD) 630.2 (M+).


20

Example 96

Reaction of the product of Preparation 203 (0.7 g, 1.1 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.52 g (84%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 530 (M+). Anal. Calc'd for C₂₉H₃₃N₆O₄ 2.7HCl: C, 55.37; H, 5.88; N, 13.36. Found: C, 55.49; H, 5.95; N, 13.56.

292

Preparation 204



Reaction of the product of Preparation 8 from Examples

Part 1 (1.0 g, 1.7 mmol), R-2-methoxymethylpyrrolidine
(0.2 mL, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (80 mL), as described in Preparation 193 gave 0.82 g (71%) of the desired compound as a white foam: ¹H-NMR is consistent with structure; MS (FD) 676.4 (M+).

Example 97

Reaction of the product of Preparation 204 (0.7 g, 1.0 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86 gave 0.56 g (92%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 576 (M+); Anal. Calc'd for C₃₁H₃₉N₆O₅·2.5HCl: C, 55.75; H, 6.41; N, 12.58. Found: C, 55.45; H, 6.36; N, 13.17.

Preparation 205

293

Reaction of the product of Preparation 8 from Examples Part 1 (1.0 g, 1.7 mmol), D-proline methyl ester (0.22 g, 1.7 mmol), 1-hydroxybenzotriazole (0.23 g, 1.7 mmol), 1- (3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dichloromethane (50 mL) as described in Example 86 gave 0.2 g (17%) of the desired compound as a yellow foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 690 (M+).

10

15

Example 98

Reaction of the product of Preparation 205 (0.18 g, 0.26 mmol), trifluoroacetic acid (3.3 mL), dichloromethane (10 mL) as described in Example 86 gave 0.16 g (93%) of the desired product as a white foam: 1H-NMR is consistent with structure; MS (FD) 590 (M+).

Preparation 206

Reaction of the product of Preparation 8 from Examples
Part 1 (1.0 g, 1.7 mmol), (S)-2-methoxymethylpyrrolidine
(0.2 mL, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (80 mL) as described in
Preparation 193 gave 0.87 g (76%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 676 (M+).

294

Example 99

Reaction of the product of Preparation 206 (0.77 g, 1.1 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.54 g (77%) of the desired product as a tan solid: ¹H-NMR is consistent with structure; MS (FD) 576.1 (M+); Anal. Calc'd for C₃₁H₃₉N₆O₅ 2.3HCl: C, 56.37; H, 6.45; N, 12.72. Found: C, 56.28; H, 6.04; N, 13.36.

10

Preparation 208

Reaction of the product of Preparation 8 from Examples
Part 1 (1.0 g, 1.7 mmol), 3,5-dimethylpyrrolidine (0.2

mL, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol),
1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (80 mL), as described in
Preparation 193 gave 0.95 g (85%) of the desired product
as a tan foam: 1H-NMR is consistent with structure; MS

(FD) 661 (M+); Anal. Calc'd for C36H48N6O6: C, 65.43; H,
7.32; N, 12.72. Found: C, 65.22; H, 7.19; N, 12.87.

Example 100

295

Reaction of the product of Preparation 208 (0.9 g, 1.4 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.69 g (83%) of the desired product as a yellow solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 561 (M+); Anal. Calc'd for $C_{31}H_{39}N_6O_4$ 2.2HCl: C, 58.10; H, 6.64; N, 13.11. Found: C, 58.04; H, 6.75; N, 13.53.

Preparation 209

10

15

Reaction of the product of Preparation 8 from Examples Part 1 (1.0 g, 1.7 mmol), azetidine (0.11 mL, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (80 mL) as described in Preparation 193 gave 0.64 g (61%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 618 (M+).

20

Example 101

Reaction of the product of Preparation 209 (0.5 g, 0.81 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.42 g (93%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 518 (M+); Anal. Calc'd for C₂₈H₃₃N₆O₄·2.5HCl: C, 55.15; H, 6.03; N, 13.78. Found: C, 55.36; H, 5.87; N, 14.01.

296

Preparation 210

Part 1 (1.0 g, 1.7 mmol), heptamethyleneimine (0.22 mL, 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (80 mL) as described in Preparation 193 gave 0.89 g (77%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 674 (M+); Anal. Calc'd for C₃₇H₅₀N₆O₆ 0.6H₂O: C, 64.82; H, 7.53; N, 12.26. Found: C, 64.59; H, 7.39; N, 12.83.

15

25

Example 102

Reaction of the product of Preparation 210 (0.83 g, 1.2 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.67 g (92%) of the desired product as a tan solid: ¹H-NMR is consistent with structure; MS (high res) calc'd for C₃₂H₄₃N₆O₄: 575.3346. Found: 575.3352.

Preparation 211

297

Reaction of the product of Preparation 8 from Examples
Part 1 (1.0 g, 1.7 mmol), 3-azabicyclo(3.2.2)nonane (0.21 g, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol),

1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (80 mL) as described in
Preparation 193 gave 1.0 g (85%) of the desired compound as a yellow foam: ¹H-NMR is consistent with structure; MS (FD) 686 (M+); Anal. Calc'd for C38H50N6O6: C, 66.45; H,

7.34; N, 12.24. Found: C, 66.65; H, 7.42; N, 12.34.

Example 103

Reaction of the product of Preparation 211 (0.95 g, 1.4 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.75 g (86%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 586 (M+); Anal. Calc'd for C₃₃H₄₁N₆O₄·2.2HCl: C, 59.43; H, 6.68; n, 12.60. Found: C, 59.54; H, 6.86; N, 12.73.

Preparation 212

298

Reaction of the product of Preparation 8 from Examples Part 1 (1.0 g, 1.7 mmol), thiazolidine (0.134 mL, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (40 mL) as described in Preparation 193 gave 0.33 g (30%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (FD) 650 (M+).

10

Example 104

Reaction of the product of Preparation 212 (0.31 g, 0.48 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.28 g (93%) of the desired product as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 550 (M+); Anal. Calc'd for $C_{28}H_{34}N_{6}O_{4}S$ ·2.6HCl: C, 52.10; H, 5.72; N, 13.01. Found: C, 52.01; H, 5.78; N, 13.23.

20

25

15

Preparation 213

Reaction of the product of Preparation 8 from Examples Part 1 (0.6 g, 1.0 mmol), thiomorpholine (0.1 mL, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide (40 mL) as described in Preparation 193 gave 0.34 g (55%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS

299

(FD) 664 (M+); Anal. Calc'd for $C_{34}H_{44}N_6O_6S^{\circ}0.1H_2O$: C, 61.43; H, 6.67; N, 12.64. Found: 59.81; H, 6.79; N, 12.31.


5

Example 105

Reaction of the product of Preparation 213 (0.3 g, 0.45 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.26 g (100%) of the desired product as a yellow solid: 1 H-NMR is consistent with structure; MS (FD) 564.1 (M+); Anal. Calc'd for $C_{29}H_{36}N_{6}O_{5}$ 3HCl: C, 51.67; H, 5.83; N, 12.47. Found: C, 52.08; H, 6.24; N, 12.48.

15

Preparation 214



To a solution of the product of Preparation 213, 0.52 g (0.78 mmol) in 10 mL of dichloromethane was added 0.36 mL (3.12 mmol) of 30% hydrogen peroxide. The solution was 20 heated to reflux for 4 h, then quenched with sodium bisulfite and concentrated. The residue was chromatographed on silica gel using 10% methanol/chloroform as eluant to yield 0.16 g (30%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (FD) 681 (M+); Anal. Calc'd for C₃₄H₄₄N₆O₇S 1.5H₂O: C, 56.42; H, 6.54; N, 11.61. Found: C, 56.39; H, 6.15; N, 11.67.

300

Example 106

Reaction of the product of Preparation 214 (0.15 g, 0.21 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.13 g (100%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 580 (M+); Anal. Calc'd for C₂₉H₃₆N₆O₅S 2.5HCl: C, 51.84; H, 5.78; N, 12.51. Found: C, 51.81; H, 5.79; N, 11.84.

Preparation 215

Reaction of the product of Preparation 8 from Examples

Part 1 (0.5 g, 0.9 mmol), N-methylpiperazine (0.1 mL, 0.9 mmol), 1-hydroxybenzotriazole (0.13 g, 1.0 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.2 g, 1.0 mmol), dimethylformamide (30 mL) as described in Preparation 193 gave 0.12 g (20%) of the desired product as a clear oil: 1H-NMR is consistent with structure; MS (FD) 662 (M+).

Example 107

301

Reaction of the product of Preparation 215 (0.1 g, 0.15 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.085 g (94%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 561 (M+).

Preparation 216

Reaction of the product of Preparation 8 from Examples

10 Part 1 (0.6 g, 1.0 mmol), 1-acetylpiperazine (0.13 g, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide (40 mL) as described in Preparation 193 gave 0.52 g (75%) of the desired compound as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 689.4 (M+).

Example 108

20 Reaction of the product of Preparation 216 (0.47 g, 0.68 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.2 g (50%) of the desired product as a tan solid: ¹H-NMR is consistent with structure; MS (FD) 589.3 (M+); Anal. Calc'd for C₃₁H₃₉N₇O₅'4.5HCl: C, 49.40; H, 5.82; N, 13.01. Found: C,

49.84; H, 5.99; N, 12.57.

Preparation 217

302

Reaction of the product of Preparation 8 from Examples Part 1 (0.5 g, 0.9 mmol), N-phenylpiperazine (0.14 mL, 0.9 mmol), 1-hydroxybenzotriazole (0.13 g, 1.0 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.2 g, 1.0 mmol), dimethylformamide (30 mL) as described in Preparation 193 gave 0.42 g (65%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 723.7 (M+); Anal. Calc'd for C40H49N7O6: C, 66.37; H, 6.82; N, 13.55. Found: C, 62.92; H, 6.87; N, 13.24.

Example 109

Reaction of the product of Preparation 217 (0.38 g, 0.5 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.2 g (87%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 623 (M+).

Preparation 218

20

25

Reaction of the product of Preparation 8 from Examples
Part 1 (0.6 g, 1.0 mmol), 1-(4-fluorophenyl)piperazine
(0.18 g, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide

WO 99/08697

303

PCT/US98/17201

(0.23 g, 1.1 mmol), dimethylformamide (30 mL) as described in Preparation 193 gave 0.42 g (57%) of the desired product as a tan foam: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 742 (M+); Anal. Calc'd for $C_{40}H_{48}N_7O_6F^*0.4H_2O$: C, 64.14; H, 6.57; N, 13.09. Found: C, 64.06; H, 6.35; N, 12.75.

Example 110

10 Reaction of the product of Preparation 218 (0.37 g, 0.5 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL) as described in Example 86 gave 0.36 g (100%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 641 (M+); Anal. Calc'd for
15 C₃₅H₄₀N₇O₄F².8HCl: C, 56.52; H, 5.80; N, 13.18. Found: C, 56.92; H, 5.79; N, 12.86.

Preparation 219

Reaction of the product of Preparation 8 from Examples
Part 1 (0.6 g, 1.0 mmol), 4-piperazinoacetophenone (0.2 g, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol),
1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g,
1.1 mmol), dimethylformamide (30 mL) as described in
Preparation 193 gave 0.42 g (55%) of the desired compound
as a tan foam: ¹H-NMR is consistent with structure; MS
(ion spray) 766.2 (M+1); Anal. Calc'd for C₄₂H₅₂N₇O₇ 0.8H₂O:

304

C, 64.56; H, 6.91; N, 12.55. Found: C, 64.59; H, 6.59; N, 12.31.

Example 111

5

10

Reaction of the product of Preparation 219 (0.36 g, 0.47 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL) as described in Example 86 gave 0.33 g (94%) of the desired product as a yellow solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 666 (M+1); Anal. Calc'd for $C_{37}H_{43}N_{7}O_{5}$ 2.5HCl: C, 58.71; H, 6.06; N, 12.95. Found: C, 58.56; H, 6.44; N, 12.60.

Preparation 220

15

Reaction of the product of Preparation 8 from Examples
Part 1 (0.6 g, 1.0 mmol), 1-(2-pyridyl)piperazine (0.16
mL, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol),
1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g,
20 1.1 mmol), dimethylformamide (40 mL) as described in
Preparation 193 gave 0.48 g (66%) of the desired product
as a white foam: ¹H-NMR is consistent with structure; MS
(ion spray) 725 (M+1); Anal. Calc'd for C₃₉H₄₈N₈O₇.0.5H₂O:
C, 63.83; H, 6.73; N, 15.27. Found: C, 63.85; H, 6.76;
N, 15.09.

Exampl 112

305

Reaction of the product of Preparation 220 (0.42 g, 0.58 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL) as described in Example 86 gave 0.35 g (88%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (high res) calc'd for C₃₄H₄₁N₃O₄: 625.3251. Found: 625.3256.

Preparation 221

10

Reaction of the product of Preparation 8 from Examples Part 1 (0.6 g, 1.0 mmol), 1-benzylpiperazine (0.18 mL, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide (30 mL) as described in Preparation 193 gave 0.3 g (40%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 737.6 (M+); Anal. Calc'd for C₄₁H₅₁N₇O₆: C, 66.74; H, 6.97; N, 13.29. Found: C, 66.67; H, 7.08; N, 13.09.

20

Example 113

Reaction of the product of Preparation 221 (0.28 g, 0.38 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL) as described in Example 86 gave 0.19 g (70%) of the

306

desired product as a yellow solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 637 (M+); Anal. Calc'd for $C_{36}H_{43}N_{7}O_{4}$ 3.5HCl: C, 56.49; H, 6.12; N, 12.81. Found: C, 56.77; H, 6.44; N, 12.31.

5

Preparation 222

Reaction of the product of Preparation 8 from Examples
Part 1 (0.6 g, 1.0 mmol), 1-((3-trifluoromethyl)-210 pyridyl)piperazine (0.23 g, 1.0 mmol), 1hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1
mmol), dimethylformamide (40 mL) as described in
Preparation 193 gave 0.5 g (63%) of the product as a tan
15 foam: ¹H-NMR is consistent with structure; MS (ion spray)
793 (M+1).

Example 114

Reaction of the product of Preparation 222 (0.42 g, 0.53 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL) as described in Example 86 gave 0.34 g (85%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (ion spray) 693 (M+1); Anal. Calc'd for C₃₅H₃₉N₆O₄F₃ 2.3HCl: C, 54.13; H, 5.36; N, 14.43. Found: C, 54.00; H, 5.55; N, 14.07.

307

Preparation 223



To a solution of d-pipecolinic acid, 2.0 g (15.5 mmol) in 30 mL of 4N sodium hydroxide as 0°C was added 2.9 mL of benzyl chloroformate dropwise. The mixture was stirred overnight slowly warming to ambient temperature then was quenched into 5N hydrochloric acid. The mixture was extracted with chloroform. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated to yield 4.1 g (100%) of the desired product as a colorless oil: ¹H-NMR is consistent with structure; MS (FD) 264 (M+).

Preparation 224



15

20

25

A solution of the product of Preparation 223, 4.1 g (15.5 mmol) and p-toluenesulfonic acid, 0.6 g in 100 mL of absolute ethanol was refluxed for 7 h. The mixture was quenched with solid sodium bicarbonate and concentrated. The residue was partitioned between ethyl acetate and water and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was chromatographed on silica gel using chloroform as eluant to yield 0.91 g (20%) of the desired product as a colorless oil: ¹H-NMR is consistent with structure; MS (FD) 291 (M+); Anal. Calc'd for C₁₆H₂₁NO₄: C, 65.96; H, 7.27; N, 4.81. Found: C, 66.21; H, 7.23; N, 4.93.

308



A solution of the product of Preparation 224, 0.9 g (3.0 mmol) in 20 mL of ethyl acetate and 40 mL of absolute ethanol was added to a slurry of 10% palladium on carbon, 0.5 g in 20 mL of ethyl acetate. The slurry was hydrogenated at 40 psi for 40 min, then was filtered through celite and concentrated to yield 0.3 g (64%) of the desired product as a colorless oil: This material was carried on without purification.

10

15

20

Preparation 226

Reaction of the product of Preparation 8 from Examples Part 1 (1.1 g, 1.9 mmol), the product of Preparation 225 (0.3 g, 1.9 mmol), 1-hydroxybenzotriazole (0.28 g, 2.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.43 g, 2.1 mmol), dimethylformamide (40 mL), as described in Preparation 193 gave 0.81 g (59%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 719 (M+); Anal. Calc'd for C₃₈H₅₀N₆O₈: C, 63.49; H, 7.01; N, 11.69. Found: C, 62.98; H, 7.33; N, 11.51.

Example 115

25

309

Reaction of the product of Preparation 226 (0.75 g, 1.0 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL) as described in Example 86 gave 0.6 g (87%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 619 (M+); Anal. Calc'd for C₃₃H₄₂N₆O₆ 2HCl: C, 57.31; H, 6.41; N, 12.15. Found: C, 57.09; H, 6.50; N, 12.04.

Preparation 227

10

Reaction of the product of Preparation 8 from Examples
Part 1 (1.0 g, 1.7 mmol), ethyl nipecotate (0.27 mL, 1.7
mmol), 1-hydroxybenzotriazole (0.25 g, 1.9 mmol), 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9
mmol), dimethylformamide (80 mL) as described in
Preparation 193 gave 0.97 g (79%) of the desired product
as a tan foam: ¹H-NMR is consistent with structure; MS
(FD) 718 (M+).

20

25

Example 116

Reaction of the product of Preparation 227 (0.91 g, 1.26 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.7 g (84%) of the desired product as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 618 (M+); Anal. Calc'd for $C_{33}H_{42}N_{6}O_{6}$ 2.2HCl: C, 56.71; H, 6.37; N, 12.02. Found: C, 56.71; H, 6.44; N, 12.45.

PCT/US98/17201

Preparation 228

Reaction of the product of Preparation 8 from Examples

Part 1 (0.7 g, 1.2 mmol), 3,5-dimethylpiperidine (0.16 mL, 1.2 mmol), 1-hydroxybenzotriazole (0.18 g, 1.32 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.27 g, 1.32 mmol), dimethylformamide (80 mL) as described in Preparation 193 gave 0.8 g (100%) of the

desired product as a tan solid: 1H-NMR is consistent with structure; MS (FD) (674 (M+).

Example 117

- 15 Reaction of the product of Preparation 228 (0.47 g, 0.7 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as in Example 86 gave 0.4 g (93%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (high res) calc'd for C32H43N6O4: 575.3346.
- 20 Found: 575.3341.

Preparation 229

Reaction of the product of Preparation 8 from Examples
25 Part 1 (1.0 g, 1.7 mmol), 4-phenylpiperidine (0.28 g, 1.7

311

mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (80 mL) as described in Preparation 193 gave 0.92 g (75%) of the desired compound as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 722 (M+).

Example 118

Reaction of the product of Preparation 229 (0.88 g, 1.2 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.77 g (96%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 622.2 (M+); Anal. Calc'd for C₃₆H₄₂N₆O₄·2.3HCl: C, 61.19; H, 6.33; N, 11.89. Found: C,

61.02; H, 6.35; N, 11.97.

Preparation 229A

Reaction of the product of Preparation 8 from Examples
Part 1 (0.6 g, 1.0 mmol), 4-(3trifluoromethylphenyl)piperidine hydrochloride (DE
3500898, DE 2048589)(0.27 g, 1.0 mmol), triethylamine
(0.16 mL, 1.1 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1
mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide
(0.23 g, 1.1 mmol), dimethylformamide (40 mL) as
described in Preparation 193 gave 0.4 g (50%) of the
desired product as a tan foam: 1H-NMR is consistent with

312

structure; MS (FD) 791 (M+); Anal. Calc'd for $C_{42}H_{49}F_3N_6O_6$: C, 63.79; H, 6.35; N, 10.63. Found: C, 63.56; H, 6.53; N, 10.57.

5

Example 119

Reaction of the product of Preparation 229A (0.36 g, 0.45 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL) as described in Example 86 gave 0.34 g (100%) of the desired product. $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 690 (M+); Anal. Calc'd for $C_{37}H_{41}N_6O_4F_3$ 2.5HCl: C, 56.86; H, 5.61; N, 10.75. Found: C, 56.72; H, 5.88; N, 10.48.

15

10

Preparation 230

Reaction of the product of Preparation 8 from Examples
Part 1 (0.6 g, 1.0 mmol), 4-(4-methoxyphenyl)piperidine
hydrochloride (WO 9518118, EP 630887)(0.27 g, 1.0 mmol),

triethylamine (0.16 mL, 1.1 mmol), 1-hydroxybenzotriazole
(0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide
(40 mL), as described in Preparation 193 gave 0.45 g
(60%) of the desired product as a tan foam: 1H-NMR is

consistent with structure; MS (FD) 754 (M+); Anal. Calc'd
for C₄₂H₅₂N₆O₇: C, 67.00; H, 6.96; N, 11.16. Found: C,
66.66; H, 7.40; N, 10.95.

313

Exampl 120

Reaction of the product of Preparation 230 (0.4 g, 0.53 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 86 gave 0.32 g (84%) of the desired product as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 652 (M+); Anal. Calc'd for $C_{37}H_{44}N_{6}O_{5}$ 2HCl: C, 61.24; H, 6.39; N, 11.58. Found: C, 60.97; H, 6.38; N, 11.33.

10

15

20

Preparation 231

Reaction of the product of Preparation 8 from Examples Part 1 (0.6 g, 1.0 mmol), 4-(2-methoxyphenyl)piperidine (0.19 g, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide (30 mL) as described in Preparation 193 gave 0.63 g (84%) of the desired product as a white foam: 1 H-NMR is consistent with structure; MS (FD) 752 (M+); Anal. Calc'd for $C_{42}H_{52}N_6O_7$ 0.1H2O: C, 65.44; H, 7.06; N, 10.90. Found: C, 65.56; H, 7.13; N, 10.89.


Example 121

314

Reaction of the product of Preparation 231 (0.59 g, 0.78 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL) as described in Example 86 gave 0.42 g (74%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 652 (M+); Anal. Calc'd for C₃₇H₄₄N₆O₅ 2HCl: C, 61.24; H, 6.39; N, 11.58. Found: C, 60.94; H, 6.35; N, 11.33.

10

Preparation 232



Reaction of the product of Preparation 8 from Examples
Part 1 (1.0 g, 1.7 mmol), ethyl isonipecotate (0.27 mL,
1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9
mmol), dimethylformamide (60 mL) as described in
Preparation 193 gave 0.87 g (71%) of the desired product
as a yellow foam: ¹H-NMR is consistent with structure; MS
(FD) 718 (M+).

20

Example 122

Reaction of the product of Preparation 232 (0.85 g, 1.2 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as in Example 86 gave 0.72 g (91%) of the desired

315

product as a white solid: ^{1}H -NMR is consistent with structure; MS (FD) 618 (M+); Anal. Calc'd for $C_{33}H_{42}N_6O_6$: 2.6HCl: C, 55.56; H, 6.30; N, 11.78. Found: C, 55.89; H, 6.34; N, 12.27.

5

Preparation 233

Reaction of the product of Preparation 8 from Examples
Part 1 (0.84 g, 1.45 mmol), 4-methoxypiperidine

10 hydrochloride(Baker, et al.; J. Med. Chem. 1992, 35(10),
1722-34)(0.22 g, 1.45 mmol), triethylamine (0.22 mL, 1.45 mmol), 1-hydroxybenzotriazole (0.21 g, 1.6 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.32 g, 1.6 mmol), dimethylformamide (40 mL) as described in

15 Preparation 193 gave 0.74 g (76%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (FD) 676.6 (M+); Anal. Calc'd for C36H48N6O7: C, 63.89; H,
7.15; N, 12.42. Found: C, 63.84; H, 7.17; N, 12.12.

20

25

Example 123

Reaction of the product of Preparation 233 (0.66 g, 1.0 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL) as described in Example 86 gave 0.54 g (84%) of the desired product as a white solid: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 576.3 (M+); Anal. Calc'd for $C_{31}H_{40}N_6O_5$ 2.5HCl: C, 55.75; H, 6.41; N, 12.58. Found: C, 55.86; H, 6.52; N, 12.27.

316

Preparation 234



A slurry of 4-ethylpyridine, 10 g (93.3 mmol) and 5%

5 Rh/C, 5 g in 135 mL of absolute ethanol was hydrogenated at 60 psi and 50°C overnight. The reaction mixture was filtered through celite and distilled at atmospheric pressure to yield 3.6 g (34%) of the desired product as a colorless oil: ¹H-NMR is consistent with structure; MS

10 (FD) 114 (M+); b.p. = 153-155°C.

Preparation 235

Reaction of the product of Preparation 8 from Examples

Part 1 (0.6 g, 1.0 mmol), 4-ethylpiperidine (0.13 g, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide (40 mL) as described in Preparation 193 gave 0.57 g (85%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (ion spray) 675.4 (M+1). Anal. Calc'd for C₃₇H₅₀N₆O₆ 0.1CHCl₃: C, 64.88; H, 7.35; N, 12.24. Found: C, 65.01; H, 6.95; N, 12.16.

Example 124

25

317

Reaction of the product of Preparation 235 (0.57 g, 0.84 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL) as described in Example 86 gave 0.34 g (63%) of the desired product as a white solid: $^1\text{H-NMR}$ is consistent with structure; MS (ion spray) 575 (M+1); Anal. Calc'd for $C_{32}H_{42}N_6O_4$ 2HCl: C, 59.35; H, 6.85; N, 12.98. Found: C, 59.19; H, 6.65; N, 12.80.

Preparation 236


 \rightarrow

10

A slurry of 4-isopropylpyridine, 15.8 g (130 mmol) and 5% Rh/C, 16 g in 125 mL of absolute ethanol was hydrogenated at 60 psi and 50°C for 18 h. The reaction mixture was filtered through celite and distilled at atmospheric pressure to yield 2.4 g (14.4%) of the desired product as a colorless oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 127 (M+); b.p. = 178-180°C; Anal. Calc'd for C₈H₁₇N: C, 75.52; H, 13.47; N, 11.01. Found: C, 75.36; H, 13.40; N, 11.09.

20

Preparation 237



Reaction of the product of Preparation 8 from Examples
Part 1 (0.6 g, 1.0 mmol), 4-isopropylpiperidine (0.13 g,
1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1
mmol), dimethylformamide (20 mL) as described in
Preparation 193 gave 0.55 g (80%) of the desired product
as a tan foam: ¹H-NMR is consistent with structure; MS
(ion spray) 689.5 (M+1). Anal. Calc'd for C₃₈H₅₀N₆O₆ 2HCl:

318

C, 66.26; H, 7.61; N, 12.20. Found: C, 66.16; H, 7.46; N, 12.03.

Example 125

5

10

Reaction of the product of Preparation 237 (0.45 g, 0.65 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 86 gave 0.37 g (86%) of the desired product as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (high res) calc'd for $C_{33}H_{43}N_6O_4$: 589.3502. Found: 589.3509.

Preparation 238

A slurry of 4-t-butylpyridine, 17.4 g (128 mmol) and 5% Rh/Al₂O₃, 6.8 g in 130 mL of absolute ethanol was hydrogenated at 60 psi and 50°C overnight. The reaction mixture was filtered through celite and distilled at atmospheric pressure to yield 7.0 g (39%) of the desired product as a colorless oil: ¹H-NMR is consistent with structure; MS (FD) 142 (M+); b.p. = 155°C.

Preparation 239

Reaction of the product of Preparation 8 from Examples

25 Part 1 (0.6 g, 1.0 mmol), 4-t-butylpiperidine (0.14 g,

1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1
(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide (20 mL) as described in

319

Preparation 193 gave 0.59 g (84%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (ion spray) 703.6 (M+1). Anal. Calc'd for C₃₉H₅₂N₆O₆ 0.5H₂O: C, 65.80; H, 7.79; N, 11.80. Found: C, 65.63; H, 7.55; N, 11.88.

Example 126

Reaction of the product of Preparation 239 (0.48 g, 0.68 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL) as described in Example 86 gave 0.33 g (72%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (ion spray) 603.4 (M+1); Anal. Calc'd for C₃₄H₄₆N₆O₄ 2.2HCl: C, 59.80; H, 7.11; N, 12.30. Found: C, 59.63; H, 7.03; N, 12.37.

Preparation 240


Reaction of the product of Preparation 8 from Examples

20 Part 1 (1.0 g, 1.7 mmol), 4-propylpiperidine (0.22 g, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (60 mL) as described in Preparation 193 gave 0.93 g (80%) of the desired product

25 as a yellow foam: ¹H-NMR is consistent with structure; MS (FD) 688.5 (M+); Anal. Calc'd for C₃₈H₅₂N₆O₆: C, 66.20; H, 7.61; N, 12.20. Found: C, 66.19; H, 7.64; N, 12.47.

320

Example 127



Reaction of the product of Preparation 240 (0.93 g, 1.4 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86, gave 0.8 g (92%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 588.2 (M+); Anal. Calc'd for C₃₃H₄₃N₆O₄ 2.2HCl: C, 59.58; H, 6.98; N, 12.63. Found: C, 59.26; H, 7.19; N, 12.82.

10

15

20

25

Preparation 241

To a solution of 4-hydroxypiperidine, 5.0 g (50 mmol) in 80 mL of 4N sodium hydroxide at 0°C was added 9.3 mL (65 mmol) of benzyl chloroformate dropwise. The reaction mixture was stirred overnight while slowly warming to ambient temperature, then was poured into 100 mL of 5N hydrochloric acid. The mixture was extracted with chloroform. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was chromatographed on silica gel using a gradient of chloroform to 5% methanol/chloroform as eluant to yield 11.5 g (97%) of the desired product as a colorless oil: ¹H-NMR is consistent with structure; MS (FD) 235 (M+); Anal. Calc'd for C₁₃H₁₇NO₃: C, 66.36; H, 7.28; N, 5.95. Found: C, 66.39; H, 7.19; N, 5.95.

Preparation 242

PCT/US98/17201 WO 99/08697

321

To a solution of the product of Preparation 241, 10.6 g (45 mmol) in 100 mL of ether and 100 mL of dimethylformamide at 0°C was added 2.16 g (54 mmol) of sodium hydride. After stirring for 15 min at 0°C, 4.1 mL (54 mmol) of chloromethyl methyl ether was added dropwise. The reaction mixture was stirred overnight while slowly warming to ambient temperature and was then concentrated. The residue was partitioned between ethyl acetate and water and was extracted with ethyl acetate. 10 The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was chromatographed on silica gel using 3-5% methanol/chloroform as eluant to yield 6.64 g (65%) of the desired product as a colorless oil: 1H-NMR is 15 consistent with structure; MS (FD) 279.1 (M+); Anal. Calc'd for $C_{15}H_{21}NO_4$: C, 64.50; H, 7.58; N, 5.01. Found: C, 65.00; H, 7.20; N, 5.30.

Preparation 243 20

A solution of the product of Preparation 242, 2.0 g (7.1 mmol) in 20 mL of ethyl acetate and 40 mL of absolute ethanol was added to a slurry of 10% palladium on carbon, 1.0 g in 20 mL of ethyl acetate. The mixture was 25 hydrogenated at 40 psi for 4 h then filtered through celite and concentrated to yield 0.71 g (68%) of the desired product as a colorless oil: 1H-NMR is consistent with structure and was carried on without further characterization. 30

Preparation 244

322

Reaction of the product of Preparation 8 from Examples
Part 1 (1.0 g, 1.7 mmol), the product of Preparation 243
(0.25 g, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9

5 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4
g, 1.1 mmol), dimethylformamide (30 mL), as described in
Preparation 193, gave 0.63 g (53%) of the desired product
as a tan foam: ¹H-NMR is consistent with structure; MS
(FD) 706.5 (M+); Anal. Calc'd for C₃₇H₅₀N₆O₈ 0.5H₂O: C,
10 62.08; H, 7.18; N, 11.74. Found: C, 62.12; H, 7.32; N,
11.56.

Example 128

To a solution of the product of Preparation 244, 0.53 g (0.75 mmol) 12 mL of dichloromethane was added 6 mL of trifluoroacetic acid. The reaction mixture was stirred for 2 h, then concentrated. The residue was slurried in ether and filtered to yield 0.5 g (85%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 561.9 (M+); Anal. Calc'd for C₃₀H₃₈N₆O₅ 2 trifluoroacetic acid: C, 51.65; H, 5.10; N, 10.63. Found: C, 51.88; H, 5.36; N, 10.61.

25

323

Reaction of the product of Preparation 8 from Examples
Part 1 (0.6 g , 1.0 mmol), 4-phenoxypiperidine
hydrochloride(Boswell, et al.; J. Med. Chem. 1974, 17(9),

5 1000-8)(0.2 g, 1.0 mmol), triethylamine (0.16 mL, 1.1
mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1
mmol), dimethylformamide (40 mL) as described in
Preparation 193, gave 0.55 g (74%) of the desired product

10 as a tan foam: 1H-NMR is consistent with structure; MS
(FD) 738 (M+); Anal. Calc'd for C41H50N6O7: C, 66.65; H,
6.82; N, 11.37. Found: C, 66.35; H, 6.64; N, 11.28.

Example 129

15

20

Reaction of the product of Preparation 245 (0.46 g, 0.6 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 86, gave 0.3 g (70%) of the desired product as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 638.3 (M+); Anal. Calc'd for $C_{36}\text{H}_{42}\text{N}_{6}\text{O}_{5}$ 2HCl: C, 60.76; H, 6.23; N, 11.81. Found: C, 60.62; H, 6.32; N, 11.63.

PCT/US98/17201 WO 99/08697

324

Reaction of the product of Preparation 8 from Examples Part 1 (1.0 g, 1.7 mmol), 4-benzylpiperidine (0.3 mL, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (80 mL), as described in Preparation 193, gave 0.82 g (66%) of the desired product as a yellow foam: 1H-NMR is consistent with structure; MS (FD) 736.3 (M+); Anal. Calc'd for $C_{42}H_{52}N_6O_6$ ·1.1 H_2O : C, 66.66; H, 7.22; N, 11.11. Found: C, 66.47; H, 6.92; N, 10 11.67.

Example 130

Reaction of the product of Preparation 246 (0.8 g, 1.1 15 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86, gave 0.65 g (88%) of the desired product as a white solid: 1H-NMR is consistent with structure; MS (FD) 637 (M+); Anal. Calc'd for $C_{37}H_{44}N_6O_4\cdot 2.1HC1:$ C, 62.30; H, 6.51; N, 11.78. Found: C, 20 62.21; H, 6.59; N, 12.12.

325

(0.7 g, 1.2 mmol), 4-benzoylpiperidine hydrochloride (0.27 g, 1.2 mmol), triethylamine (0.18 mL, 1.32 mmol), 1-hydroxybenzotriazole (0.18 g, 1.32 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.27 g, 1.32 mmol), dimethylformamide (30 mL), as described in Preparation 193 gave 0.65 g (72%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 751 (M+).

10

Example 131

Reaction of the product of Preparation 247 (0.51 g, 0.68 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86, gave 0.47 g (96%) of the desired product as a yellow solid: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 651 (M+); Anal. Calc'd for $C_{37}H_{42}N_6O_5$ 2HCl: C, 60.49; H, 6.08; N, 11.44. Found: C, 60.44; H, 6.00; N, 11.41.

20

25

15


Preparation 248

Reaction of the product of Preparation 8 from Examples
Part 1 (1.0 g, 1.7 mmol), 4-(4-fluorobenzoyl)piperidine
hydrochloride (0.42 g, 1.7 mmol), triethylamine (0.26 mL,
1.9 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9
mmol), dimethylformamide (80 mL), as described in
Preparation 193, gave 0.97 g (75%) of the desired product

326

as a tan foam: ${}^{1}H-NMR$ is consistent with structure; MS (FD) 768.9 (M+).

Example 132




5

10

Reaction of the product of Preparation 248 (0.92 g, 1.2 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86, gave 0.75 g (88%) of the desired product as a yellow solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 669 (M+); Anal. Calc'd for $C_{37}\text{H}_{42}\text{FN}_{6}\text{O}_{5}$ 2.7HCl: C, 57.93; H, 5.74; N, 10.95. Found: C, 57.83; H, 5.81; N, 11.20.

Preparation 249



15

Reaction of the product of Preparation 8 from Examples
Part 1 (0.7 g, 1.2 mmol), 4-(4-chlorobenzoyl)piperidine
hydrochloride (0.32 g, 1.2 mmol), triethylamine (0.18 mL,
1.32 mmol), 1-hydroxybenzotriazole (0.18 g, 1.32 mmol),

1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.27 g,
1.32 mmol), dimethylformamide (30 mL), as described in
Preparation 193, gave 0.83 g (89%) of the desired product
as a tan foam: ¹H-NMR is consistent with structure; MS
(FD) 785 (M+).

25

Example 133

327

Reaction of the product of Preparation 249 (0.64 g, 0.82 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86, gave 0.51 g (82%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 685 (M+); Anal. Calc'd for C₃₇H₄₁N₆O₅Cl 2HCl: C, 58.61; H, 5.72; N, 11.08. Found: C, 58.34; H, 5.93; N, 11.00.

10 Preparation 250


Reaction of the product of Preparation 8 from Examples
Part 1 (0.6 g, 1.0 mmol), 4-(4-methylbenzoyl)piperidine
hydrochloride (0.24 g, 1.0 mmol), triethylamine (0.16 mL,
1.1 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1
mmol), dimethylformamide (30 mL), as described in
Preparation 193, gave 0.54 g (71%) of the desired product
as a tan foam: ¹H-NMR is consistent with structure; MS
(FD) 765.4 (M+); Anal. Calc'd for C₄₃H₅₂N₆O₇: C, 67.52; H,
6.85; N, 10.99. Found: C, 67.32; H, 6.65; N, 10.76.

Example 134

328

Reaction of the product of Preparation 250 (0.5 g, 0.65 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 86, gave 0.33 g (69%) of the desired product as a white solid: ¹H-NMR is consistent with structure; (FD) 655.2 (M+); Anal. Calc'd for C₃₈H₄₄N₆O₅Cl 2.3HCl: C, 60.96; H, 6.23; N, 11.23. Found: C, 60.92; H, 6.55; N, 11.10.

Preparation 251



10

15

20

Reaction of the product of Preparation 8 from Examples Part 1 (0.6 g, 1.0 mmol), 4-(4-methoxybenzoyl)piperidine hydrochloride (0.26 g, 1.0 mmol), triethylamine (0.16 mL, 1.1 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide (40 mL), as described in Preparation 193, gave 0.23 g (29%) of the desired product as a tan foam: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 780.8 (M+); Anal. Calc'd for C₄₃H₅₂N₆O₈: C, 66.14; H, 6.71; N, 10.76. Found: C, 66.14; H, 6.60; N, 10.65.

Example 135

Reaction of the product of Preparation 251 (0.2 g, 0.26 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 86, gave 0.19 g (100%) of the desired product. ¹H-NMR is consistent with structure; MS (FD) 681.3 (M+); Anal. Calc'd for C₃₈H₄₄N₆O₆·2HCl: C,

PCT/US98/17201 WO 99/08697

329

60.56; H, 6.15; N, 11.15. Found: C, 60.43; H, 6.29; N, 10.89.

Preparation 252

5

10

15

20

Reaction of the product of Preparation 8 from Examples Part 1 (0.6 g, 1.0 mmol), 4-(alpha-cyanobenzyl)piperidine hydrochloride (0.24 g, 1.0 mmol), triethylamine (0.16 mL, 1.1 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide (40 mL), as described in Preparation 193, gave 0.4 g (53%) of the desired product as a tan foam: 1H-NMR is consistent with structure; MS (FD) 761.7 (M+); Anal. Calc'd for $C_{43}H_{51}N_7O_6$: C, 67.79; H, 6.75; N, 12.87. Found: C, 67.54; H, 6.45; N, 12.67.

Example 136

Reaction of the product of Preparation 252 (0.31 g, 0.4 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 86, gave 0.22 g (76%) of the desired product as a tan foam: 1H-NMR is consistent with structure; MS (FD) 661.1 (M+); Anal. Calc'd for $C_{38}H_{43}N_{7}O_{4}$ 2HC1: C, 62.12; H, 6.17; N, 13.35. Found: C, 61.95; H, 8.38; H, 13.19. 25

330

Reaction of the product of Preparation 8 from Examples
Part 1 (0.6 g, 1.0 mmol), 4-(2-ethyl-2naphthyl))piperidine hydrochloride (Efange et al., J. Med

5 Chem. 1993 36(9), 1278-83) (0.28 g, 1.0 mmol),
triethylamine (0.16 mL, 1.1 mmol), 1-hydroxybenzotriazole
(0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide
(40 mL), as described in Preparation 193, gave 0.58 g

10 (74%) of the desired product as a tan foam: 1H-NMR is
consistent with structure; MS (FD) 801 (M+); Anal. Calc'd
for C47H56N6O6: C, 70.48; H, 7.05; N, 10.49. Found: C,
70.52; H, 6.87; N, 10.50.

15

20

Example 137

Reaction of the product of Preparation 253 (0.5 g, 0.62 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 86, gave 0.22 g (46%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 701 (M+).

331

Reaction of the product of Preparation 8 from Examples Part 1 (0.6 g, 1.0 mmol), 4-(phenpropyl)piperidine (0.2 g, 1.0 mmol), triethylamine (0.16 mL, 1.1 mmol), 1hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide (40 mL), as described in Preparation 193, gave 0.58 g (76%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 765.4 (M+); Anal. Calc'd for $C_{44}H_{56}N_6O_6$: C, 69.09; H, 7.38; N, 10.99. Found: C, 68.95; H, 7.29; N, 11.04. 10

Example 138

Reaction of the product of Preparation 254 (0.41 g, 0.5 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 15 mL), as described in Example 86, gave 0.32 g (87%) of the desired product as a white solid: 1H-NMR is consistent with structure; MS (FD)664 (M+); Anal. Calc'd for $C_{39}H_{48}N_6O_4$ 2HCl: C, 63.49; H, 6.83; N, 11.39. Found: C, 63.30; H, 6.77; N, 11.29. 20

Preparation 255

Reaction of the product of Preparation 8 from Examples Part 1 (1.0 g, 1.7 mmol), 4,4-dimethylpiperidine 25 hydrochloride (0.26 g, 1.7 mmol), triethylamine (0.26 mL, 1.9 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9

332

mmol), dimethylformamide (80 mL), as described in Preparation 193, gave 0.84 g (73%) of the desired product as a tan solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 674 (M+).

5

Example 139

Reaction of the product of Preparation 255 (0.82 g, 1.2 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86, gave 0.65 g (89%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 574.2 (M+).

Preparation 257

15

20

25

Reaction of the product of Preparation 8 from Examples Part 1 (0.6 g, 1.0 mmol), 3-azaspiro(5.5)undecane hydrochloride (Knoelker, et al; Synlett. 1992, Issue 5, 371-87)(0.19 g, 1.0 mmol), triethylamine (0.16 mL, 1.1 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.23 g, 1.1 mmol), dimethylformamide (40 mL), as described in Preparation 193, gave 0.42 g (60%) of the desired compound as a tan foam: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 715 (M+); Anal. Calc'd for C40H54N6O6: C, 67.21; H, 7.61; N, 11.76. Found: C, 67.13; H, 7.38; N, 11.71.

10

Example 140

Reaction of the product of Preparation 257 (0.37 g, 0.5 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 86, gave 0.32 g (94%) of the desired product as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 614 (M+); Anal. Calc'd for $C_{35}H_{46}N_{6}O_{4}$ 2.3HCl: C, 60.17; H, 6.97; N, 12.03. Found: C, 60.14; H, 7.13; N, 11.82.

Preparation 258

Reaction of the product of Preparation 8 from Examples

Part 1 (0.6 g, 1.0 mmol), 4-(10,11-dihydro-5H-dibenzo(A,D)cyclohepten-5-yl)piperidine (U.S. 4,626,542)

(0.28 g, 1.0 mmol), 1-hydroxybenzotriazole (0.15 g, 1.1 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide

(0.23 g, 1.1 mmol), dimethylformamide (40 mL), as

described in Preparation 193, gave 0.69 g (82%) of the desired compound as a tan foam: 1H-NMR is consistent with structure; MS (FD) 839 (M+).

Example 141

PCT/US98/17201 WO 99/08697

334

Reaction of the product of Preparation 258 (0.63 g, 0.75 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 86, gave 0.44 g (72%) of the 5 desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 739 (M+); Anal. Calc'd for $C_{45}H_{50}N_6O_4$ 2HCl: C, 66.58; H, 6.46; N, 10.35. Found: C, 66.41; H, 6.62; N, 10.25.

10

20

Preparation 259

Reaction of the product of Preparation 8 from Examples Part 1 (1.0 g, 1.0 mmol), 4-cyano, 4-phenylpiperidine hydrochloride (0.38 g, 1.7 mmol), triethylamine (0.26 mL, 1.9 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-15 (3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (40 mL), as described in Preparation 193, gave 1.0 g (79%) of the desired product as a white foam: 1H-NMR is consistent with structure; MS (FD) 747.8 (M+).

Example 142

Reaction of the product of Preparation 259 (1.0 g, 1.3 25 mmol), trifluoroacetic acid (4 mL), dichloromethane (12

335

mL), as described in Example 86, gave 0.71 g (76%) of the desired product as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 647.2 (M+); Anal. Calc'd for $C_{37}\text{H}_{41}\text{N}_{7}\text{O}_{4}$ 2.6HCl: C, 59.85; H, 5.92; N, 13.20. Found: C, 59.76; H, 5.88; N, 13.10.

Preparation 260

Reaction of the product of Preparation 8 from Examples

Part 1 (1.0 g, 1.0 mmol), 4-acetyl-4-phenylpiperidine
hydrochloride (0.4 g, 1.7 mmol), triethylamine (0.26 mL,
1.9 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9
mmol), dimethylformamide (80 mL), as described in

Preparation 193, gave 0.52 g (40%) of the desired product
as a white foam: ¹H-NMR is consistent with structure; MS
(FD) 765 (M+); Anal. Calc'd for C₄₃H₅₂N₆O₇ 0.1H₂O: C, 65.97;
H, 6.95; N, 10.73. Found: C, 65.68; H, 6.93; N, 11.17.

20 Example 14

Reaction of the product of Preparation 260 (0.5 g, 0.65 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86, gave 0.44 g (96%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 664.4 (M+); Anal. Calc'd for C₃₈H₄₃N₆O₅ 2.7HCl: C, 59.80; H, 6.17; N, 11.01. Found: C, 59.61; H, 6.18; N, 11.24.

336

Preparation 261

Reaction of the product of Preparation 8 from Examples

Part 1 (0.7 g, 1.2 mmol), 4-phenyl-4-propionylpiperidine hdyrochloride (0.3 g, 1.2 mmol), triethylamine (0.18 mL, 1.32 mmol), 1-hydroxybenzotriazole (0.18 g, 1.32 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.27 g, 1.32 mmol), dimethylformamide (40 mL), as described in Preparation 193, gave 0.71 g (76%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 779 (M+); Anal. Calc'd for C44H54N6O7: C, 67.85; H, 6.99; N, 10.79. Found: C, 67.56; H, 7.10; N, 10.95.

15

20

25

Example 144

Reaction of the product of Preparation 261 (0.58 g, 0.74 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86, gave 0.54 g (100%) of the desired product. ¹H-NMR is consistent with structure; MS (FD) 678 (M+); Anal. Calc'd for C₃₉H₄₆N₆O₅·2.6HCl: C, 60.55; H, 6.33; N, 10.86. Found: C, 60.40; H, 6.27; N, 10.79.

337

Reaction of the product of Preparation 8 from Examples Part 1 (0.7 g, 1.2 mmol), 4-butyryl-4-phenylpiperidine hdyrochloride (0.3 g, 1.2 mmol), triethylamine (0.18 mL, 1.32 mmol), 1-hydroxybenzotriazole (0.18 g, 1.32 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.27 g, 1.32 mmol), dimethylformamide (40 mL), as described in Preparation 193, gave 0.76 g (80%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (FD) 793 (M+).

10

Example 145

Reaction of the product of Preparation 262 (0.65 g, 0.82 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86, gave 0.5 g (80%) of the desired product as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 692 (M+); Anal. Calc'd for $C_{40}H_{48}N_{6}O_{5}$ 2.3HCl: C, 61.86; H, 6.53; N, 10.82. Found: C, 61.96; H, 6.63; N, 10.60.

Preparation 263

Reaction of the product of Preparation 8 from Examples
25 Part 1 (1.0 g, 1.7 mmol), 1,2,3,6-tetrahydropyridine

(0.16 mL, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (80 mL), as described in Preparation 193, gave 0.63 g (57%) of the desired product as a tan solid: ¹H-NMR is consistent with structure; MS (FD) 644 (M+); Anal. Calc'd for C₃₅H₄₄H₆O₆: C, 65.20; H, 6.88; N, 13.03. Found: C, 65.30; H, 7.04; N, 13.14.

Example 146

10

Reaction of the product of Preparation 263 (0.62 g, 0.96 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86. gave 0.53 g (95%) of the desired product as a white solid: $^1\text{H-NMR}$ is consistent with structure; MS (high res) calc'd for $C_{50}H_{37}N_6O_4$: 545.2876. Found: 545.2883.

Preparation 264

Reaction of the product of Preparation 8 from Examples
Part 1 (0.7 g, 1.2 mmol), 4-phenyl-1,2,3,6tetrahydropyridine (0.16 g, 1.2 mmol), 1hydroxybenzotriazole (0.18 g, 1.32 mmol), 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.27 g, 1.32
mmol), dimethylformamide (40 mL), as described in
Preparation 193, gave 0.36 g (50%) of the desired product
as a tan foam: ¹H-NMR is consistent with structure; MS
(FD) 721.2 (M+); Anal. Calc'd for C41H48N6O6.0.5H2O: C,

339

67.47; H, 6.77; N, 11.51. Found: C, 67.56; 6.81; N, 11.20.

Example 147

5

10

Reaction of the product of Preparation 264 (0.34 g, 0.47 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 86, gave 0.28 g (85%) of the desired product as a white solid: $^1\text{H-NMR}$ is consistent with structure; MS (ion spray) 621.4 (M+1).

Preparation 265

Reaction of the product of Preparation 8 from Examples

Part 1 (1.0 g, 1.7 mmol), 3,5-dimethylpiperidine (0.23 mL, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol),

1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (80 mL), as described in

Preparation 193, gave 0.93 g (81%) of the desired product

as a tan solid: 1H-NMR is consistent with structure; MS


(FD) 674 (M+).

Example 148

340

Reaction of the product of Preparation 265 (0.78 g, 1.16 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86, gave 0.63 g (89%) of the desired product as a yellow solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 574 (M+); Anal. Calc'd for $C_{32}H_{41}N_{6}O_{4}$ 2.8HCl: C, 56.79; H, 6.67; N, 12.42. Found: C, 56.75; H, 6.70; N, 12.12.

Preparation 266



10

Reaction of the product of Preparation 8 from Examples
Part 1 (1.0 g, 1.7 mmol), decahydroquinoline (0.26 mL,
1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9
mmol), dimethylformamide (80 mL), as described in
Preparation 193, gave 1.0 g (83%) of the desired product
as a tan solid: ¹H-NMR is consistent with structure; MS
(FD) 700.5 (M+); Anal. Calc'd for C₃₉H₅₂N₆O₆: C, 66.84; H,
7.48; N, 11.99. Found: C, 66.69; H, 7.48; N, 12.15.

20

Example 149

Reaction of the product of Preparation 266 (0.93 g, 1.3 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86, gave 0.77 g (93%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 600 (M+); Anal. Calc'd for

341

 $C_{34}H_{43}N_6O_4$ 2.6HCl: C, 58.71; H, 6.75; N, 12.08. Found: C, 58.54; H, 6.98; N, 11.93.

Preparation 267

5

Reaction of the product of Preparation 8 from Examples Part 1 (1.0 g, 1.7 mmol), 1,2,3,4-tetrahydroisoquinoline (0.22 mL, 1.7 mmol), 1-hydroxybenzotriazole (0.26 g, 1.9 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.4 g, 1.9 mmol), dimethylformamide (40 mL), as described in Preparation 193, gave 0.91 g (77%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (FD) 694.8 (M+).

15

25

10

Example 150

Reaction of the product of Preparation 267 (0.88 g, 1.3 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 86, gave 0.87 g (100%) of the desired product as a white solid: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 594.3 (M+); Anal. Calc'd for $C_{34}H_{38}N_6O_4$ 2.5HCl: C, 59.54; H, 5.95; N, 12.25. Found: C, 59.31; H, 6.02; N, 12.06.

To a solution of tert-butylcarbonyl-alpha-aminoisobutic acid(10.0 g, 50 mmol) and methyl iodide, 3.6 mL (55 mmol) in 200 mL of tetrahydrofuran stirring at 0°C was added 4.4 g (55 mmol) of sodium hydride. After 1 h, the 5 mixture was warmed to ambient temperature and stirred overnight. The reaction mixture was quenched with water, concentrated, and dissolved in water. The aqueous solution was acidified to pH = 3.4 with solid citric acid. Ethyl acetate was added and the mixture was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated to yield 10.8 g (100%) of the desired product as a white solid: 1H-NMR is consistent with structure; MS (ion spray) 218 (M+1).

15

10

Preparation 269

To a solution of the product of Preparation 268, 9.75 g (45.0 mmol), tert-butyloxycarbonyl-O-benzyl-D-serin, 11.0 g (45.0 mmol), Hunig's base, 28 mL (157.5 mmol) and 1-20 hydroxybenzotriazole, 6.8 g (49.5 mmol) in 100 mL of dimethylformamide at 0°C, was added 9.5 g (49.5 mmol) of 1-(3-(dimethylamino)propyl)-3-ethylcarbodiimide hydrochloride. The reaction mixture was slowly warmed to ambient temperature, stirred overnight, then 25 concentrated. The residue was partitioned between ethyl acetate and water and extracted with ethyl acetate. combined organic extracts were washed with 0.1 N hydrochloric acid, saturated sodium bicarbonate and brine, dried over sodium sulfate, filtered and 30 concentrated. The residue was chromatographed on silica gel using 3% methanol/chloroform as eluant to yield 13.8 g (75%) of the desired product as a tan oil: $^{1}\text{H-NMR}$ is

343

consistent with structure; MS (ion spray) 409.4 (M+1); Anal. Calc'd for $C_{21}H_{32}N_2O_6$: C, 61.60; H, 8.12; N, 6.84. Found: C, 61.45; H, 8.10; N, 6.87.

5

Preparation 270

To a solution of the product of Preparation 269, 0.5 g (1.22 mmol) in 10 mL of tetrahydrofuran was added a solution of lithium hydroxide, 0.09 g (3.7 mmol) in 5 mL of water. The reaction mixture was stirred for 2 h and acidified to pH = 2.4 with 1N hydrochloric acid. Water was added and the mixture was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated to yield 0.48 g (100%) of the desired product as a clear oil: ¹H-NMR is consistent with structure; MS (ion spray) 395.2 (M+1); Anal. Calc'd for C20H30N2O6: C, 60.90; H, 7.67; N, 7.10. Found: C, 61.04; H, 7.70; N, 7.34.

20


Preparation 271

Reaction of the product of Preparation 165 (4.5 g, 14.7 mmol), 10% palladium on carbon (5.8 g), tetrahydrofuran (120 mL), the product of Preparation 270 (5.8 g, 14.7 mmol), 1-hydroxybenzotriazole (2.2 g, 16.2 mmol), dicyclohexylcarbodiimide (3.3 g, 16.2 mmol), as in Example 63, gave 6.55 g (68%) of the desired compound as a tan foam: ¹H-NMR is consistent with structure; MS (ion spray) 652.4 (M+1).

WO 99/08697

344

Preparation 272



Reaction of the product of Preparation 271 (6.5 g, 10.0 mmol), lithium hydroxide (0.3 g, 12.0 mmol), dioxane (100 mL), water (50 mL), as in Example 63, gave 6.24 g (100%) of the desired compound as a tan foam: ¹H-NMR is consistent with structure; MS (ion spray) 624.4 (M+1); Anal. Calc'd for C₄₃H₄₁N₅O₈: C, 61.62; H, 6.63; N, 11.23.

10 Found: C, 61.34; H, 6.68; N, 11.33.

Preparation 273

Reaction of the product of Preparation 272 (6.1 g, 1.0 mmol), 4-methylpiperidine (1.2 mL, 1.0 mmol), 1-hydroxybenzotriazole (1.5 g, 1.1 mmol), dicyclohexylcarbodiimide (2.3 g, 1.1 mmol), dimethylformamide (90 mL), as in Example 63, gave 6.18 g (88%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (ion spray) 705.6 (M+1); Anal. Calc'd for C₃₀H₅₂N₆O₇: C, 64.75; H, 7.44; N, 11.92. Found: C, 64.63; H, 7.39; N, 11.99.

Examples 151 and 152

Reaction of the product of Preparation 273 (5.65 g, 8.0 mmol), trifluoroacetic acid (16 mL), dichloromethane (40 mL), as in Example 63, gave 4.08 g (85%) of the desired product as the free base: 1H-NMR is consistent with structure; MS (ion spray) 605.4 (M+1); Anal. Calc'd for $C_{33}H_{44}N_6O_5$: C, 65.54; H, 7.33; N, 13.90. Found: C, 65.30; H, 7.54; N, 13.93. Resolution of the diastereomers (3.5 g, 5.8 mmol) by chiral HPLC gave the respective isomers which were individually treated with a saturated solution of hydrochloric acid in diethyl ether to give the desired products:

10

Example 151. (Isomer 1): 1.4 g (36%); ¹H-NMR is consistent with structure; $t_R = 6.50 \text{ min}$; MS (ion spray) 605.1 (M+1); Anal. Calc'd for $C_{33}H_{44}N_6O_5$ 2.4HCl: C, 57.26; 15 H, 6.76; N, 12.14. Found: C, 57.30; H, 6.67; N, 12.00. **Example 152.** (Isomer 2): 1.43 g (36%); ¹H-NMR is consistent with structure; $t_R = 7.91 \text{ min; MS (ion spray)}$ 605.1 (M+1); Anal. Calc'd for $C_{33}H_{44}N_6O_5$ 2.2HCl: C, 57.87; H, 6.80; N, 12.27. Found: C, 57.71; H, 6.84; N, 12.09. 20

To a solution of 5-phenylvaleric acid, 19.55 g (110 mmol) in 100 mL of absolute ethanol was added 2.0 g (10.4 mmol) of p-toluenesulfonic acid. The mixture was heated to reflux overnight, cooled to ambient temperature and concentrated. The residue was chromatographed on silica gel using 20% ethyl acetate/hexanes as eluant to yield 21.3 g (94%) of the desired product as a colorless oil: 30

PCT/US98/17201 WO 99/08697


346

¹H-NMR is consistent with structure; MS (FD) 206 (M+); Anal. Calc'd for $C_{12}H_{18}O_2$: C, 75.69; H, 8.81. Found: C, 75.69; H, 8.91.

5

10

15



To a solution of diisopropylamine, 11.5 mL (80 mmL) in 250 mL of tetrahydrofuran at -78°C was added 50.0 mL of 1.0 M n-butyl lithium dropwise. After stirring at -78°C for 20 minutes, 15.0 mL (116 mmol) of trimethylsilyl chloride was added followed by a solution of the product of Preparation 274 (15.0 g, 72.5 mmol) in 50 mL of tetrahydrofuran. The reaction mixture was stirred 1 h at -78°C and was then concentrated. To a slurry of the resulting white solid in 500 mL of ethylene glycol dimethylether at -78°C was added 13.5 g (76.1 mmol) of Nbromosuccinimide. The reaction mixture was stirred 3 h at -78°C, then was quenched with water and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was chromatographed on silica gel using 2.5% ethyl acetate/hexanes to yield 14.8 g (72%) of the desired product as a yellow oil: 1H-NMR is consistent with structure; MS (FD) 284, 286 (M+).

25

20

Preparation 276



To a slurry of sodium hydride, 2.4 g (60 mmol) in 300 mL of dimethylformamide at 0°C was added 6.8 g (60 mmol) of 4-nitroimidazole. After stirring for 20 min at 0°C, a 30 solution of 14.1 g (50 mmol) of the product of

347

Preparation 275 in 30 mL of dimethylformamide was added. The reaction mixture was stirred overnight, slowly warming to ambient temperature and was concentrated. The residue was partitioned between ethyl acetate and water and was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The resulting residue was chromatographed on silica gel using a chloroform as eluant to yield 12.7 g (71%) of the desired product as an orange oil: ¹H-NMR is consistent with structure; MS (FD) 317 (M+); Anal. Calc'd for C₁₆H₁₉N₃O₄: C, 60.56; H, 6.04; N, 13.24. Found: C, 60.78; H, 6.09; N, 12.98.

10

15

20

25

Preparation 277

To a solution of the product of Preparation 276, 12.2 g (38.4 mmol) in 60 mL of tetrahydrofuran was added 40 mL of 5N sodium hydroxide. The mixture was stirred at ambient temperature for 1 h, then acidified to pH = 3 with 5N hydrochloric acid and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated to yield 9.3 g (84%) of the desired product as an orange oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 290 (M+).



To a solution of the product of Preparation 277, 9.0 g (31 mmol), pyrrolidine, 2.6 mL (31 mmol) and 1-hydroxybenzotriazole, 4.6 g (34.1 mmol) in 200 mL of dimethylformamide was added 7.0 g (34.1 mmol) of dicyclohexylcarbodiimide. The reaction mixture was stirred overnight at ambient temperature and was then concentrated in vacuo. The residue was partitioned between ethyl acetate and water and was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was chromatographed on silica gel using 5% methanol/chloroform as eluant to yield 10.3 g (97%) of the desired product as a tan oil: ¹H-NMR is consistent with structure; MS (FD) 312 (M+).

Preparation 278A

To a slurry of 0.8 g of 10% palladium on carbon in 20 mL of dioxane was added a solution of 0.8 g (3.5 mmol) of the product of Preparation 278 in 20 mL of dioxane. The mixture was hydrogenated at 40 psi of hydrogen for 3 h and filtered through celite. To this solution was added 1.2 g (5.8 mmol) of the product of Preparation 1d, 0.52 g (3.85 mmol) of 1-hydroxybenzotriazole and 0.8 g (3.85 mmol) of dicyclohexylcarbodiimide. The reaction was stirred overnight at ambient temperature, filtered and concentrated. The residue was partitioned between ethyl acetate and water and was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The resulting residue was chromatographed on silica gel

349

using 2.5% methanol/chloroform as eluant to yield 0.65 g (16%) of the desired product as a red foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 674 (M+).

5

10

Example 153

To a solution of the product of Preparation 278A, 0.55 g (0.8 mmol) in 24 mL of dichloromethane was added 8 mL of trifluoroacetic acid. The reaction mixture was stirred for 2 h, quenched with aqueous sodium carbonate and extracted with chloroform. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. To a solution of the residue in ethyl acetate was added ether/hydrochloric acid. The resultant slurry was concentrated to yield 0.47 g (94%) of the desired product as a tan solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 573.9 (M+); Anal. Calc'd for $C_{32}H_{42}N_6O_4$ 0.2HCl: C, 59.35; H, 6.85; N, 12.98. Found: C, 59.04; H, 6.96; N, 12.52.

20

Preparation 279

To a solution of hydrocinnamic acid, 12.35 g (82.2 mmol) in 300 mL of dichloromethane was added 4.2 mL (82.2 mmol) of bromine dropwise. The reaction mixture was illuminated for 15 min and then stirred an additional 15 min and concentrated. The residue was trituated with ether/hexanes to yield 14.7 g (78%) of the desired product as a white solid: ¹H-NMR is consistent with

350

structure; MS (FD) 228, 230 (M+); Anal. Calc'd for C₉H₉O₂Br: C, 47.19; H, 3.96. Found: C, 47.28; H, 4.02.

Preparation 280



5

15

20

To a slurry of sodium hydride, 1.1 g (27.3 mmol) in 50 mL of dimethylformamide at 0°C was added 3.1 g (27.3 mmol) of 4-nitroimidazole. The resulting mixture was stirred 20 min, then 3.0 g (13.0 mmol) of product of Preparation 279 was added. The reaction mixture was stirred overnight slowly warming to ambient temperature, then concentrated. The residue was partitioned between ethyl acetate and 1N hydrochloric acid and was basified to pH = 3.0 with 1N sodium hydroxide. The aqueous layer was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was absorbed onto a silica pad and chromatographed on silica gel using a gradient of 10-40% methanol/chloroform to yield 0.55 g (14.4%) of the desired product as a white solid: 1H-NMR is consistent with structure; MS (FD) 261 (M+).

Preparation 281



To a solution of the product of Preparation 280, 2.4 g (9.3 mmol), pyrrolidine, 0.77 mL (9.3 mmol) and 1-hydroxybenzotriazole, 1.6 g (10.2 mmol) in 200 mL of dimethylformamide was added 2.1 g (10.2 mmol) of dicyclohexylcarbodiimide. The reaction mixture was stirred overnight at ambient temperature and was then

351

concentrated in vacuo. The residue was partitioned between ethyl acetate and water and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was chromatographed on silica gel using 2.5% methanol/chloroform as eluant to yield 0.84 g (29%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (FD) 314 (M+).

5

10

Preparation 282

O'STYYY

To a slurry of 0.7 g of 10% palladium on carbon in 30 mL of ethyl acetate was added a solution of 0.8 g (2.5 mmol) of the product of Preparation 281 in 30 mL of ethyl acetate, 60 mL of tetrahydrofuran and 60 mL of ethanol. 15 The mixture was hydrogenated at 40 psi of hydrogen for 1 h, filtered through celite and concentrated. The residue was dissolved in 60 mL of dimethylformamide. solution was added 0.95 g (2.5 mmol) of the product of Preparation 1d, 0.37 g (2.75 mmol) of 1-20 hydroxybenzotriazole and 0.57 g (2.75 mmol) of dicyclohexylcarbodiimide. The reaction was stirred overnight at ambient temperature, filtered and concentrated. The residue was partitioned between ethyl acetate and water and was extracted with ethyl acetate. 25 The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The resulting residue was chromatographed on silica gel using 2.5% methanol/chloroform as eluant to yield 0.32 g (20%) of the desired product as a tan foam: 1H-NMR is 30 consistent with structure; MS (FD) 646.3 (M+).

352

To a solution of the product of Preparation 282, 0.25 g (0.38 mmol) in 12 mL of dichloromethane was added 4 mL of trifluoroacetic acid. The reaction mixture was stirred for 2 h, quenched with aqueous sodium carbonate and extracted with chloroform. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. To a solution of the residue 10 in 20 mL of ethyl acetate was added 40 mL of ether hydrochloric acid. The resultant slurry was concentrated to yield 0.22 g (96%) of the desired product as a tan solid: 1H-NMR is consistent with structure; MS (high res) calc'd for $C_{30}H_{39}N_6O_4$: 547.3033. Found: 15 547.3037.

Preparation 283



To a slurry of sodium hydride, 6.72 g (168 mmol) in 400 20 mL of dimethylformamide at 0°C was added 19.0 g (168 mmol) of 4-nitroimidazole. The resulting slurry was stirred 20 min, then a solution of 25 g (140 mmol) of ethyl 3bromopropionate in 10 mL of dimethylformamide was added dropwise. The reaction mixture was stirred overnight 25 slowly warming to ambient temperature, then concentrated. The residue was partitioned between ethyl acetate and was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium

353

sulfate, filtered and concentrated. The residue was absorbed onto a silica pad and chromatographed on silica gel using a gradient of 20--70% ethyl acetate/hexanes to yield 22.71 g (76%) of the desired product as a yellow oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 213 (M+); Anal. Calc'd for $C_8H_{11}N_3O_4$: C, 45.07; H, 5.20; N, 19.71. Found: C, 45.08; H, 5.18; N, 19.42.

Preparation 284



10

15

20

To a solution of the product of Preparation 283, 22.46 g (100 mmol) in 100 mL of tetrahydrofuran and 100 mL of ethanol was added 100 mL of 5N sodium hydroxide. The mixture was stirred at ambient temperature for 1 h, then acidified to pH = 2.5 with 5N hydrochloric acid and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated to yield 7.8 g (42%) of the desired product as an orange oil: 1H -NMR is consistent with structure; MS (FD) 185 (M+).

Preparation 285



To a solution of the product of Preparation 284, 5.3 g

(29 mmol), pyrrolidine, 2.4 mL (29 mmol) and 1hydroxybenzotriazole, 4.3 g (32 mmol) in 50 mL of
dimethylformamide was added 6.6 g (32 mmol) of
dicyclohexylcarbodiimide. The reaction mixture was
stirred overnight at ambient temperature and was then

concentrated in vacuo. The residue was partitioned

5

10

PCT/US98/17201

between ethyl acetate and water and was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was absorbed onto a silica pad and chromatographed on silica gel using a gradient of 2.5-5% methanol/chloroform as eluant to yield 0.69 g (10%) of the desired product as a yellow solid: ¹H-NMR is consistent with structure; MS (FD) 238 (M+); Anal. Calc'd for $C_{10}H_{14}N_4O_3$: C, 50.41; H, 5.92; N, 23.52. Found: C, 50.59; H, 5.75; N, 23.47.

Preparation 286

To a slurry of 0.66 g of 10% palladium on carbon in 15 mL of ethyl acetate was added a solution of 0.66 g (2.8 15 mmol) of the product of Preparation 285 in 6 mL of dichloromethane and 15 mL of ethanol. The mixture was hydrogenated at 40 psi of hydrogen for 40 min and filtered through celite and concentrated. The residue was dissolved in 30 mL of dimethylformamide. To this 20 solution was added 1.1 g (2.8 mmol) of the product of Preparation 1d, 0.42 g (3.1 mmol) of 1hydroxybenzotriazole and 0.63 g (3.1 mmol) of dicyclohexylcarbodiimide. The reaction was stirred overnight at ambient temperature, filtered and 25 concentrated. The residue was partitioned between ethyl acetate and water and was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The resulting residue was chromatographed on silica gel 30 using a gradient of chloroform to 3% methanol/chloroform as eluant to yield 0.67 g (42%) of the desired product as

355

a tan foam: $^{1}H-NMR$ is consistent with structure; MS (FD) 570.2 (M+); Anal. Calc'd for $C_{29}H_{42}N_6O_6$: C, 61.03; H, 7.42; N, 14.73. Found: C, 60.94; H, 7.26; N, 14.55.

5

Example 155

To a solution of the product of Preparation 286, 0.6 g (1.1 mmol) in 12 mL of dichloromethane was added 4 mL of trifluoroacetic acid. The reaction mixture was stirred for 2 h, quenched with solid sodium carbonate and 10 extracted with chloroform. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. To a solution of the residue in 10 mL of ethyl acetate was added 10 mL of 15 ether hydrochloric acid. The resultant slurry was concentrated to yield 0.52 g (94%) of the desired product as a tan foam: 1H-NMR is consistent with structure; MS (FD) 471 (M+1); Anal. Calc'd for $C_{24}H_{34}N_6O_4$ 2.25HCl: C, 52.16; H, 6.61; N, 15.21. Found: C, 52.48; 6.87; N, 14.81. 20

Preparation 287



Ethyl-4-bromobutyrate (18.4 mL, 156 mmol), 4
25 nitroimidazole (17.6 g, 130 mmol), sodium hydride (6.24 g, 156 mmol), dimethylformamide (400 mL), as described Preparation 283 gave 23.4 g (80%) of the desired product as a yellow oil: ¹H-NMR is consistent with structure; MS

356

(FD) 227 (M+); Anal. Calc'd for $C_9H_{13}N_3O_4$: C, 47.58; H, 5.77; N, 18.49. Found: C, 47.48; H, 5.50; N, 18.30.

Preparation 288



5

Reaction of the product of Preparation 287 (22.84 g, 100 mmol), 5N sodium hydroxide (100 mL), tetrahydrofuran (100 mL), ethanol (100 mL), as described in Preparation 284 gave 9.8 g (50%) of the desired product as an orange solid: H-NMR is consistent with structure; MS (FD) 200.1 (M+); Anal. Calc'd for C₇H₉N₃O₄: C, 42.22; H, 4.56; N, 21.10. Found: C, 41.97; H, 4.63; N, 21.04.

Preparation 289



15

20

Reaction of the product of Preparation 288 (4.87 g, 25 mmol), pyrrolidine (2.4 mL, 25 mmol), 1-hydroxybenzotriazole (3.7 g, 27.5 mmol), dicyclohexylcarbodiimide (5.6 g, 27.5 mmol), dimethylformamide (50 mL), as described in Preparation 285, gave 0.65 g (11%) of the desired product as a tan oil: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 252 (M+); Anal. Calc'd for $C_{11}H_{16}N_4O_3$: C, 52.37; H, 6.39; N, 22.21. Found: C, 52.59; H, 6.50; N, 22.44.

25

357

Reaction of the product of Preparation 289 (0.6 g, 2.4 mmol), 10% palladium on carbon (0.3 g), dichloromethane (6 mL), ethyl acetate (20 mL), absolute ethanol (20 mL), 386979 (0.91 g, 2.4 mmol), 1-hydroxybenzotriazole (0.36 g, 2.64 mmol), dicyclohexylcarbodiimide (0.54 g, 2.64 mmol), dimethylformamide (40 mL), as described in Preparation 286 gave 0.69 g (50%) of the desired product as a tan oil: ¹H-NMR is consistent with structure; MS (FD) 585.4 (M+); Anal. Calc'd for C30H44N6O6: C, 61.63; H, 7.58; N, 14.37. Found: C, 61.35; H, 7.50; N, 14.30.

Example 156

15 Reaction of the product of Preparation 290 (0.65 g, 1.1 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 155 gave 0.58 g (98%) of the desired hydrochloric acid salt as an oil: ¹H-NMR is consistent with structure; MS (FD) 484 (M+); Anal. Calc'd for C₂₅H₃₆N₆O₄·2.3HCl: C, 52.82; H, 6.79; N, 14.78. Found: C, 53.01; H, 6.88; N, 14.40.

Preparation 291



Reaction of ethyl-5-bromovalerate (25 g, 120 mmol), 4-nitroimidazole (16.3 g, 144 mmol), sodium hydride (5.8 g, 144 mol), dimethylformamide (400 mL), as described

358

Preparation 283 gave 21.9 g (75%) of the desired product as a colorless oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 241.1 (M+); Anal. Calc'd for $C_{10}H_{15}N_{3}O_{4}$: C, 49.79; H, 6.27; N, 17.42. Found: C, 49.63; H, 6.16; N, 17.22.

5

Preparation 292



Reaction of the product of Preparation 291 (21.08 g, 87 mmol), 5N sodium hydroxide (100 mL), tetrahydrofuran (100 mL), ethanol (100 mL), as described in Preparation 284 gave 11.9 g (64%) of the desired product as a tan solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 214 (M+); Anal. Calc'd for $C_{8}\text{H}_{11}\text{N}_{3}\text{O}_{4}$: C, 45.07; H, 5.20; N, 19.71. Found: C, 44.89; H, 4.92; N, 19.44.

15

10

Preparation 293



Reaction of the product of Preparation 292 (9.9 g, 46.4 mmol), pyrrolidine (4 mL, 46.4 mmol), 1
20 hydroxybenzotriazole (7.0 g, 51.0 mmol), dicyclohexylcarbodiimide (10.6 g, 51.0 mmol)), triethylamine (3.23 mL, 46.4 mmol), as described in Preparation 285 gave 8.4 g (69%) of the desired product as a colorless oil: ¹H-NMR is consistent with structure;

25 MS (FD) 266 (M+); Anal. Calc'd for C₁₂H₁₈N₄O₃: C, 54.12; H, 6.81; N, 21.04. Found: C, 54.35; H, 6.91; N, 20.91.

359

Preparation 294

Reaction of the product of Preparation 293 (1.0 g, 3.8 mmol), 10% palladium on carbon (0.5 g), dichloromethane (5 mL), ethyl acetate (10 mL), ethanol (20 mL), the product of Preparation 1d (1.45 g, 3.8 mmol), 1-hydroxybenzotriazole (0.56 g, 4.2 mmol), dicyclohexylcarbodiimide (0.86 g, 1.1 mmol), dimethylformamide (40 mL), as described in Preparation 286 gave 0.8 g (35%) of the desired product as a yellow oil: ¹H-NMR is consistent with structure; MS (FD) 598.2 (M+).

Example 157

15

20

Reaction of the product of Preparation 294 (0.75 g, 1.2 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 155 gave 0.57 g (100%) of the desired product as the hydrochloric acid salt as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 499 (M+).

Preparation 295

Reaction of N-(2-chloroethyl)morpholine hydrochloride (6.9 g, 37.0 mmol), 4-nitroimidazole (5.0 g, 44.0 mmol), sodium hydride (3.27 g, 80.7 mmol), dimethylformamide

360

(150 mL), as described in Preparation 283 gave 1.1 g (13%) of the desired product as a yellow solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 226.1 (M+1); Anal. Calc'd for $C_{9}H_{14}N_{4}O_{3}$: C, 47.78; H, 6.24; N, 24.77. Found: C, 48.01; H, 6.13; N, 24.56.

Preparation 296

Reaction of the product of Preparation 295 (0.6 g, 2.6 mmol), 10% palladium on carbon (0.6 g), ethyl acetate (20 mL), ethanol (10 mL), the product of Preparation 1d (1.0 g, 2.6 mmol), 1-hydroxybenzotriazole (0.4 g, 2.9 mmol), dicyclohexylcarbodiimide (0.6 g, 2.9 mmol), dimethylformamide (40 mL), as described in Preparation 286 gave 0.34 g (24%) of the desired product as a yellow foam: ¹H-NMR is consistent with structure; MS (FD) 558 (M+1).

Example 158

20

25

5

Reaction of the product of Preparation 296 (0.14 g, 0.25 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 155 gave 0.13 g (100%) of the desired product as the hydrochloric acid salt as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 458.2 (M+).

361

Preparation 297



To a solution of 4-nitroimidazole, 4.6 g (40 mmol) and bromodiphenylmethane, 10.0 g (40 mmol) in 150 mL of dimethylformamide was added 16.6 g (120 mmol) of 5 potassium carbonate. The reaction mixture was stirred overnight at ambient temperature, filtered and concentrated. The residue was partitioned between ethyl acetate and water and was extracted with ethyl acetate. The combined organic extracts were washed with brine, 10 dried over sodium sulfate, filtered and concentrated. The resulting crude material was absorbed onto a silica pad and chromatographed on silica gel using 25-60% ethyl acetates/hexanes as eluant to yield 3.2 g (27%) of the desired product as a clear oil: ¹H-NMR is consistent with 15 structure; MS (ion spray) 280 (M+1).

Preparation 298

20 To a slurry of 0.8 g of 10% palladium on carbon in 30 mL of tetrahydrofuran was added a solution of 1.2 g (4.3 mmol) of the product of Preparation 297 in 40 mL of tetrahydrofuran. The mixture was hydrogenated at 40 psi of hydrogen for 1 h and filtered through celite. To this solution was added 1.64 g (4.3 mmol) of Preparation 4 from Examples Part 1, 0.7 g (4.7 mmol) of 1-hydroxybenzotriazole and 1.04 g (4.7 mmol) of dicyclohexylcarbodiimide. The reaction was stirred overnight at ambient temperature, filtered and

362

concentrated. The resulting residue was chromatographed on silica gel using 4% methanol/chloroform as eluant to yield 1.57 g (60%) of the desired product. $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 612.4 (M+1); Anal. Calc'd for $C_{35}H_{41}N_{5}O_{5}$: C, 68.72; H, 6.76; N, 11.45. Found: C, 68.44; H, 6.72; N, 11.15.

Example 159

To a solution of the product of Preparation 298, 0.56 g 10 (0.91 mmol) in 6 mL of dichloromethane was added 2 mL of trifluoroacetic acid. The reaction mixture was stirred for 2.5 h, then poured into a solution of saturated sodium carbonate and extracted with chloroform. combined organic extracts were washed with brine, dried 15 over sodium sulfate, filtered and concentrated. desired product was trituated with ethyl acetate/hexanes to yield 0.45 g (96%) of the desired product as a tan solid: 1H-NMR is consistent with structure; MS (ion spray) 512.6 (M+1); Anal. Calc'd for $C_{30}H_{33}N_5O_3$.0.2 H_2O : C, 20 69.94; H, 6.53; N, 13.59. Found: C, 69.88; H, 6.36; N, 13.25.

Preparation 299



25

To a solution of 4,4'-dimethoxybenzhydrol, 5.0 g (20 mmol), 4-nitroimidazole, 2.31 g (20 mmol) and triphenylphosphine, 5.3 g (20 mmol) in 200 mL of tetrahydrofuran at 0°C was added 5.0 mL (32 mmol) of

diethyl azodicarboxylate dropwise. The resulting mixture was slowly warmed to ambient temperature, stirred overnight and concentrated. The residue was partitioned between ethyl acetate and water and extracted with ethyl acetate. The combined organic extracts were washed with 5 brine, dried over sodium sulfate, filtered and concentrated. The residue was absorbed onto a silica pad and chromatographed on silica gel using 20-40% ethyl acetates/hexane as eluant. The resulting residue was chromatographed on silica gel using 3% 10 methanol/chloroform as eluant to yield 4.92 g (72%) of the desired product as a yellow oil: 1H-NMR is consistent with structure; MS (ion spray) 340 (M+1); Anal. Calc'd for $C_{18}H_{17}N_3O_4 \cdot 0.3H_2O$: C, 62.71; H, 5.15; N, 12.19. Found: 15 C, 62.48; H, 4.83; N, 11.84.

Preparation 300

Reaction of the product of Preparation 299 (1.8 g, 5.3 mmol), 10% palladium on carbon (1.0 g), tetrahydrofuran (60 mL), 386979 (2.0 g, 5.3 mmol), 1-hydroxybenzotriazole (0.8 g, 5.83 mmol), dicyclohexylcarbodiimide (1.2 g, 5.83 mmol), as described in Preparation 159 gave 1.3 g (36%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (ions spray) 672.7 (M+1); Anal. Calc'd for C₃₇H₄₅N₅O₇: C, 66.15; H, 6.75; N, 10.43. Found: C, 66.29; H, 6.82; N, 10.63.

Example 160

364

Reaction of the product of Preparation 300 (1.24 g, 1.8 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Example 159, Chromatographed over silica gel (5% methanol/chloroform) to yield 0.73 g (69%) of the desired product as a white foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 572.4 (M+1); Anal. Calc'd for $C_{32}H_{37}N_{5}O_{5}$: C, 67.23; H, 6.52; N, 12.25. Found: C, 67.05; H, 6.59; N, 11.97.

10

15

20

25

Preparation 301



To a slurry of sodium hydride, 2.64 g (66.0 mmol) in 300 mL of dimethylformamide at 0°C was added 7.4 g (66.0 mmol) of 4-nitroimidazole. After stirring for 20 min at 0°C, 15.0 g (55.0 mmol) of desyl bromide was added. The reaction mixture was stirred 65 h, as it warmed to ambient temperature and was concentrated. The residue was partitioned between ethyl acetate and water and was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The resulting residue was chromatographed on silica gel using a gradient of chloroform to 5% methanol/chloroform as eluant to yield 10.1 g (60%) of the desired product as a yellow oil which solidifies upon standing. 1H-NMR is consistent with structure; MS (ion spray) 307.1 (M+1); Anal. Calc'd for $C_{17}H_{13}N_3O_3$: C, 66.44; H, 4.26; N, 13.67. Found: C, 66.31; H, 4.22; N, 13.39.

365

Preparation 302

To a slurry of 0.3 g of 10% palladium on carbon in 20 mL of tetrahydrofuran was added a solution of 0.8 g (2.6 mmol) of the product of Preparation 301 in 20 mL of tetrahydrofuran. The mixture was hydrogenated at 40 psi of hydrogen for 1 h and filtered through celite. To this solution was added 1.0 g (2.6 mmol) of the product of Preparation 1d, 0.4 g (2.9 mmol) of 1-10 hydroxybenzotriazole and 0.6 g (2.9 mmol) of dicyclohexylcarbodiimide. The reaction was stirred overnight at ambient temperature, filtered and concentrated. The resulting residue was chromatographed on silica gel using a gradient from chloroform to 5% 15 methanol/chloroform as eluant to yield 1.3 g (76%) of the desired product. 1H-NMR is consistent with structure; MS (FD) 639.4 (M+); Anal. Calc'd for $C_{36}H_{41}N_5O_6$: C, 65.74; H, 6.59; N, 10.65. Found: C, 65.91; H, 6.21; N, 10.67.

20

Example 161

To a solution of the product of Preparation 302, 1.0 g (1.56 mmol) in 12 mL of dichloromethane was added 4 mL of trifluoroacetic acid. The reaction mixture was stirred for 1.5 h, then quenched with solid sodium carbonate and extracted with chloroform. The combined organic extracts were washed with brine, dried over sodium sulfate,

366

filtered and concentrated. The residue was dissolved in ethyl acetate and ether hydrochloric acid was added. The mixture was concentrated to yield 0.68 g (71%) of the desired product as a tan solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 539 (M+); Anal. Calc'd for $C_{31}H_{33}N_{5}O_{4}$ 2HCl: C, 60.78; H, 5.76; N, 11.43. Found: C, 60.51; H, 5.84; N, 11.12.

Preparation 303

10

To a slurry of desoxyanisoin, 20 g (78 mmol) in 200 mL of carbon tetrachloride at 0°C was added 4 mL (78 mmol) of bromine, dropwise over 30 min. The reaction mixture was stirred overnight while slowly warming to ambient

15 temperature, then concentrated. The residue was absorbed onto a silica pad and chromatographed on silica gel using a gradient of 10-30% ethyl acetate/hexanes as eluant to yield 18 g (69%) of the desired product as a white solid:

1 h-NMR is consistent with structure; MS (ion spray) 335,

20 337 (M+1).

Preparation 304



Reaction of the product of Preparation 303 (7.0 g, 20.0 mmol), 4-nitroimidazole (2.34 g, 20.0 mmol), sodium hydride (1.0 g, 24.0 mmol), dimethylformamide (100 mL), as described in Preparation 301. The crude material was absorbed onto a silica pad and chromatographed over silica gel using a gradient of 30-80% ethyl

30 acetate/hexanes to yield 2.8 g (38%) of the desired product as a tan foam: ¹H-NMR is consistent with

367

structure; MS (ion spray) 366.1 (M+1); Anal. Calc'd for $C_{19}H_{17}N_3O_5$: C, 62.12; H, 4.66; N, 11.44. Found: C, 62.05; H, 4.55; N, 11.57.

Preparation 305

5

20

25

Reaction of the product of Preparation 304 (1.5 g, 4.1 mmol), 10% palladium on carbon (1.0 g), tetrahydrofuran (40 mL), the product of Preparation 1d (1.56 g, 4.1 mmol), 1-hydroxybenzotriazole (0.61 g, 4.51 mmol), dicyclohexylcarbodiimide (0.93 g, 4.51 mmol), as in described in Preparation 302. The residue was chromatographed on silica gel using 3% methanol/chloroform as eluant to yield 1.95 g (68%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (ion spray) 700.8 (M+1); Anal. Calc'd for C38H45N5O8: C, 65.22; H, 6.48; N, 10.01. Found: C, 65.05; H, 6.20; N, 10.25.

Examples 162 and 163

SHOOL SHOOL

Reaction of the product of Preparation 305 (1.85 g, 2.6 mmol), trifluoroacetic acid (10 mL), dichloromethane (24 mL), as in described in Example 161, gave 1.56 g (88%) of the desired product as a product as the hydrochloric acid salt. $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 599.2 (M+1); Anal. Calc'd for $C_{33}H_{37}N_{5}O_{6}$ '2HCl: C,

368

58.93; H, 5.84; N, 10.41. Found: C, 59.05; H, 5.87; N, 10.43. Resolution of the diastereomers (0.21 g, 0.38 mmol) by chiral HPLC gave the respective isomers which were individually treated with a saturated solution of hydrochloric acid in diethyl ether to give the desired products:

Example 162. (Isomer 1) 0.104 g (40%); 1 H-NMR is consistent with structure; $t_{R} = 9.45$ min; MS (ion spray) 600.3 (M+1); Anal. Calc'd for $C_{33}H_{37}N_{5}O_{6}$ 2HCl: C, 58.93; H, 5.84; N, 10.41. Found: C, 58.66; H, 5.80; N, 10.20. Example 163. (Isomer 2) 0.066 g (25%); 1 H-NMR is consistent with structure; $t_{R} = 12.93$ min; MS (ion spray) 600.3 (M+1); Anal. Calc'd for $C_{33}H_{37}N_{5}O_{6}$ 2.3HCl: C, 57.99; H, 5.80; N, 10.25. Found: C, 57.94; H, 5.80; N, 10.12.

Preparation 306

5

10

15

20

25



To a slurry of 4-nitroimidazole, 10.0 g (89 mmol) in 30 mL of glacial acetic acid was slowly added 10 mL of fuming nitric acid. To this mixture was added 20 mL of acetic anhydride at a rate such that the temperature remained below 50°C. The reaction mixture was stirred for one hour then poured into 300 g of ice/water. A white solid precipitated out over a period of 30 min. The solid was filtered and dried to yield 6.8 g of the desired product. The filtrate was extracted with dichloromethane. The combined organic extracts were washed with saturated sodium bicarbonate and water, then dried over sodium sulfate, filtered and concentrated to yield an additional 5.1 g of the desired white solid, affording a total of 11.9 g (85%) of the desired product.

1H-NMR is consistent with structure; MS (FD) 159 (M+);

369

Anal. Calc'd for $C_3H_2N_4O_4^{\circ}0.3H_20$: C, 22.04; H, 1.60; N, 34.27. Found: C, 22.35; H, 1.52; N, 33.87.

Preparation 307



5


10

15

20

To a slurry of 9-aminofluorene hydrochloride, 2.1 g (10.0 mmol) in 20 mL of methanol and 20 mL of water at 0°C was added 0.8 g (10.0 mmol) of sodium bicarbonate. After stirring for 10 min at 0°C, 1.5 g (10.0 mmol) of the product of Preparation 306 was added. The reaction mixture was stirred overnight, slowly warming to ambient temperature, then concentrated. The residue was partitioned between ethyl acetate and water and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate and concentrated. The residue was absorbed onto a silica pad and chromatographed on silica gel using a gradient of 25-80% ethyl acetate/hexanes as eluant to yield 2.39 g (88%) of the desired product as a tan solid: 1H-NMR is consistent with structure; MS (ion spray) 276.2 (M+1); Anal. Calc'd for $C_{16}H_{11}N_3O_2$: C, 69.31; H, 4.00; N, 15.15. Found: C, 69.53; H, 4.08; N, 14.93.

Preparation 308



25

To a slurry of 0.95 g of 10% palladium on carbon in 30 mL of tetrahydrofuran was added a solution of 1.1 g (4.0 mmol) of the product of Preparation 307 in 30 mL of

370

tetrahydrofuran. The mixture was hydrogenated at 40 psi of hydrogen for 1 h and filtered through celite. To this solution was added 1.5 g (4.0 mmol) of the product of Preparation 1d, 0.6 g (4.4 mmol) of 1-

Preparation 1d, 0.6 g (4.4 mmol) of 1
hydroxybenzotriazole and 0.9 g (4.4 mmol) of
dicyclohexylcarbodiimide. The reaction was stirred
overnight at ambient temperature, filtered and
concentrated. The residue was slurried in ethyl acetate,
filtered and water was added. The mixture was extracted
with ethyl acetate. The combined organic extracts were
washed with brine, dried over sodium sulfate, filtered
and concentrated. The resulting residue was
chromatographed on silica gel using 5%
methanol/chloroform as eluant to yield 1.16 g (48%) of
the desired product. H-NMR is consistent with structure;
MS (ion spray) 610 (M+1); Anal. Calc'd for C35H39N5O5: C,
68.95; H, 6.59; N, 11.49. Found: C, 68.83; H, 6.46; N,

20

25

30

11.27.

Example 164

To a solution of the product of Preparation 308, 1.1 g (1.8 mmol) in 6 mL of dichloromethane was added 2 mL of trifluoroacetic acid. The reaction mixture was stirred for 3 h, then poured into a solution of saturated sodium carbonate. The mixture was extracted with chloroform. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was dissolved in ethyl acetate and ether/hydrochloric acid was added. The mixture was concentrated to yield 0.9 g (90%) of the desired product

371

as a pink solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 510.4 (M+1); Anal. Calc'd for $C_{30}H_{31}N_{5}O_{3}$ 2HCl: C, 61.86; H, 5.71; N, 12.02. Found: C, 61.70; H, 5.78; N, 11.86.

5

Preparation 309




Reaction of 2,2-diphenylethylamine (2.5 g, 12.7 mmol), sodium bicarbonate (1.07 g, 12.7 mmol), the product of Preparation 306 (2.0 g, 12.7 mmol), methanol (20 mL), water (10 mL), as described in Preparation 307 gave 2.75 g (74%) of the desired product as a tan solid: ¹H-NMR is consistent with structure; MS (ion spray) 294.2 (M+1).

15

10

Preparation 310



Reaction of the product of Preparation 309 (1.0 g, 3.4 mmol), 10% palladium on carbon (1.2 g), tetrahydrofuran (40 mL), the product of Preparation 1d (1.3 g, 3.4 mmol), 1-hydroxybenzotriazole (0.51 g, 3.74 mmol), dicyclohexylcarbodiimide (0.8 g, 3.74 mmol), as in Example Preparation 308. 1.42 g (67%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (ion spray) 626.3 (M+1); Anal. Calc'd for C₃₆H₄₃N₅O₅: C, 69.10; H, 6.93; N, 11.19. Found: C, 68.92; H, 6.70; N, 11.21.

PCT/US98/17201

372

Example 165

Reaction of the product of Preparation 310 (1.37 g, 2.2 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as described in Example 164 -426737 gave 1.14 g (87%) of the desired product as a tan solid: 1H-NMR is consistent with structure; MS (ion spray) 526 (M+1); Anal. Calc'd for $C_{31}H_{35}N_5O_3$ 2.2HCl: C, 61.46; H, 6.19; N, 11.56. Found: C, 61.72; H, 5.95; N, 11.11.

10

20



To a slurry of 2-aminoindan hydrochloride, 2.0 g (12.7 mmol) in 20 mL of methanol and 10 mL of water at 0°C was added 1.4 g (16.4 mmol) of sodium bicarbonate. After 15 stirring for 10 min at 0°C, 2.0 g (12.7 mmol) of the product of Preparation 306 was added. The reaction mixture was stirred overnight, slowly warming to ambient temperature, then concentrated. The residue was partitioned between ethyl acetate and water and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate and concentrated. The residue was absorbed onto a silica pad and chromatographed on silica gel using a gradient of 50-80% ethyl acetate/hexanes as eluant to yield 1.91 g (66%) 25 of the desired product as a tan solid: 1H-NMR is consistent with structure; MS (ion spray) 230.2 (M+1);

373

Anal. Calc'd for $C_{12}H_{11}N_3O_2$: C, 62.88; H, 4.84; N, 18.33. Found: C, 63.28; H, 4.90; N, 17.89.

Preparation 312

5

To a slurry of 1.3 g of 10% palladium on carbon in 30 mL of tetrahydrofuran was added a solution of 1.2 g (5.3 mmol) of the product of Preparation 311 in 30 mL of tetrahydrofuran. The mixture was hydrogenated at 40 psi of hydrogen for 40 min and filtered through celite. To 10 this solution was added 2.0 g (5.3 mmol) of the product of Preparation 1d, 0.8 g (5.8 mmol) of 1hydroxybenzotriazole and 1.2 g (5.8 mmol) of dicyclohexylcarbodiimide. The reaction was stirred overnight at ambient temperature, filtered and 15 concentrated. The residue was slurried in ethyl acetate, filtered and water was added. The mixture was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The resulting residue was 20 chromatographed on silica gel using 3% methanol/chloroform as eluant to yield 1.38 g (486) of the desired product: 1H-NMR is consistent with structure; MS (ion spray) 562.4 (M+1); Anal. Calc'd for $C_{31}H_{39}N_5O_5$ 0.1CHCl3: C, 65.12; H, 6.87; N, 12.21. Found: C, 25

Example 166

65.39; H, 7.18; N, 11.97.

374

To a solution of the product of Preparation 312, 1.4 g (2.5 mmol) in 12 mL of dichloromethane was added 4 mL of trifluoroacetic acid. The reaction mixture was stirred for 2 h, then concentrated. The residue was partitioned between ethyl acetate and saturated sodium carbonate and was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was dissolved in ethyl acetate and ether hydrochloric acid was 10 added. The mixture was concentrated to yield 1.2 g (90%) of the desired product as a yellow solid: 1H-NMR is consistent with structure; MS (ion spray) 462.4 (M+1); Anal. Calc'd for $C_{26}H_{31}N_5O_3$ 2HCl: C, 57.99; H, 6.93; N, 12.01. Found: C, 57.87; H, 6.82; N, 12.73. 15

Preparation 313



20

To a slurry of 1-aminoindan, 1.25 mL (10.0 mmol) in 20 mL of methanol and 20 mL of water at 0°C, was added 1.5 g (10.0 mmol) of the product of Preparation 306. The reaction mixture was stirred overnight, slowly warming to ambient temperature and concentrated. The residue was partitioned between ethyl acetate and water and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was absorbed onto a silica pad and chromatographed on silica gel using 30-80% ethyl

375

acetate/hexanes as eluant to yield 1.13 g (49%) of the desired product as a tan oil: ¹H-NMR is consistent with structure; MS (ion spray) 230.2 (M+1).

376

Preparation 314

To a slurry of 1.0 g of 10% palladium on carbon in 20 mL of tetrahydrofuran was added a solution of 1.07 g (4.7 mmol) of the product of Preparation 313 in 20 mL of tetrahydrofuran. The mixture was hydrogenated at 40 psi of hydrogen for 30 min and filtered through celite. this solution was added 1.8 g (4.7 mmol) of the product of Preparation 1d,, 0.7 g (5.2 mmol) of 1hydroxybenzotriazole and 1.06 g (5.2 mmol) of 10 dicyclohexylcarbodiimide. The reaction was stirred overnight at ambient temperature, filtered and concentrated. The residue was dissolved in ethyl acetate, filtered, then water was added. The mixture was extracted with ethyl acetate. The combined organic 15 extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The resulting residue was chromatographed on silica gel using 3% methanol/chloroform as eluant to yield 0.99 g (38%) of the desired product: 1H-NMR is consistent with structure; 20 MS (ion spray) 562 (M+1); Anal. Calc'd for $C_{31}H_{39}N_5O_5$: C, 66.29; H, 7.00; N, 12.47. Found: C, 66.05; H, 7.12; N,

25

12.58.

Example 167

To a solution of the product of Preparation 314, 0.97 g (1.7 mmol) in 6 mL of dichloromethane was added 2 mL of

377

trifluoroacetic acid. The reaction mixture was stirred for 2 h, then poured into a solution of saturated sodium carbonate. The mixture was extracted with chloroform. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was dissolved in ethyl acetate and a saturated solution of hydrochloric acid in diethyl ether was added. The resulting mixture was concentrated to yield 0.73 g (76%) of the desired mixture of isomers as a white solid: ¹H-NMR is consistent with structure; MS (ion spray) 462.4 (M+1); Anal. Calc'd for C₂₆H₃₁N₅O₃·2.3HCl: C, 57.26; H, 6.15; N, 12.84. Found: C, 57.53; H, 6.04; N, 12.57.

15

10

5

Preparation 315



Reaction of (1S, 2R)-(-)-cis-1-Amino-2-indanol (0.9 g, 6.3 mmol), sodium bicarbonate (0.53 g, 6.3 mmol), the product of Preparation 306, (1.0 g, 6.3 mmol) in methanol (10 mL) and water (5 mL), as described in Preparation 311 gave 1.02 g (66%) of the desired product as a tan foam:

¹H-NMR is consistent with structure; MS (ion spray) 246.3 (M+1).

25

30

20

Preparation 316



To a solution of the product of Preparation 315, 2.33 g (9.5 mmol) and imidazole, 2.0 g (36.9 mmol) in 100 mL of dimethylformamide was added 1.86 g (12.3 mmol) of t-butyldimethylsilyl chloride. The reaction mixture was

378

stirred overnight at ambient temperature then concentrated. The residue was partitioned between ethyl acetate and water and was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was chromatographed over silica gel using 50% ethyl acetate/hexanes as eluent to yield 3.07 g (90%) of the desired product as a tan oil: 1H-NMR is consistent with structure; MS (ion spray) 360.2 (M+1); Anal. Calc'd for C18H25N3O3Si: C, 60.14; H, 7.01; N, 11.69. Found: C, 60.37; H, 6.92; N, 11.43.

Preparation 317

Reaction of the product of Preparation 316 (1.5 g, 4.2 mmol), 10% palladium on carbon (1.0 g), tetrahydrofuran (80 mL), the product of Preparation 1d, (1.6 g, 4.2 mmol), 1-hydroxybenzotriazole (0.63 g, 4.6 mmol), dicyclohexylcarbodiimide (0.95 g, 4.6 mmol), as described in Preparation 314 gave 0.72 g (25%) of the desired product as a reddish foam: ¹H-NMR is consistent with structure; MS (ion spray) 692.1 (M+1); Anal. Calc'd for C₃₇H₅₃N₅O₆Si: C, 64.34; H, 7.72; N, 10.12. Found: C, 64.14; H, 7.65; N, 10.05.

25

Example 168

PCT/US98/17201 WO 99/08697

379

A solution of the product of Preparation 317, 0.65 g (0.94 mmol) in 10 mL of a saturation solution of hydrochloric acid in acetic acid was stirred at ambient temperature. After 1.5 h, the solution was concentrated. The residue was partitioned between ethyl acetate and saturated sodium bicarbonate and was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate and concentrated. residue was chromatographed on silica gel using a gradient of 5-10% methanol/chloroform/ammonia as eluant to yield the desired product as the free base. resulting foam was dissolved in ethyl acetate and a saturated solution of hydrochloric acid in diethyl ether was added. The resulting slurry was concentrated to yield 0.18 g (35%) of the desired product as a white 15 solid: 1H-NMR is consistent with structure; MS (ion spray) 478.3 (M+1); Anal. Calc'd for C₂₆H₃₁N₅O₄·2.5HCl: C, 47.85; H, 5.53; N, 10.73. Found: C, 48.08; H, 5.17; N, 10.40.

20

25

30

5

10

Preparation 318



To a solution of the product of Preparation 315, 1.71 g (7.0 mmol) and triethylamine, 2.2 mL (15.6 mmol) in 80 mL of dichloromethane at 0°C was added 0.9 mL (12.6 mL) of acetyl chloride. The reaction mixture was stirred 2.5 h at 0°C, then was quenched with saturated sodium bicarbonate. The mixture was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was absorbed onto a silica pad and chromatographed on silica gel using a gradient of 30-

380

80% ethyl acetate/hexanes as eluant to yield 1.06 g (53%) of the desired product as a yellow oil: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 288.0 (M+1); Anal. Calc'd for $C_{14}H_{13}N_{3}O_{4}$: C, 58.53; H, 4.56; N, 14.63. Found: C, 58.78; H, 4.58; N, 14.36.

Preparation 319

Reaction of the product of Preparation 318 (1.0 g, 3.5 mmol), 10% palladium on carbon (2.0 g), tetrahydrofuran (60 mL), the product of Preparation 1d, (1.33 g, 3.5 mmol), 1-hydroxybenzotriazole (0.52 g, 3.9 mmol), dicyclohexylcarbodiimide (0.8 g, 3.9 mmol), described in Preparation 314 gave 0.53 g (25%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (ion spray) 620.4 (M+1); Anal. Calc'd for C₃₃H₄₁N₅O₇ 0.15CHCl₃: C, 62.44; H, 6.50; N, 10.98. Found: C, 62.42; H, 6.51; N, 11.16.

20

25


Example 169

A solution of the product of Preparation 319, 0.43 g (0.7 mmol) in 8 mL of a saturated solution of hydrochloric acid in glacial acetic acid was stirred for 1.5 h at ambient temperature then concentrated. The residue was dissolved in toluene and concentrated to yield 0.26 g (61%) of the desired product as a tan solid: ¹H-NMR is consistent with structure; MS (ion spray) 520.3 (M+1);

381

Anal. Calc'd for $C_{28}H_{33}N_5O_5$ 2HCl: C, 56.76; H, 5.95; N, 11.82. Found: C, 56.87; H, 6.07; N, 11.52.

Preparation 320




5

To a solution of the product of Preparation 315, 1.36 g (5.6 mmol) in 50 mL of dichloromethane was added 1.09 g (6.72 mmol) of 1,1'-carbonyldiimidazole. The reaction mixture was stirred overnight at ambient temperature then 3.3 mL (28 mmol) of 4-methylpiperidine was added. 10 reaction mixture was stirred overnight, washed with a 0.1N aqueous solution of hydrochloric acid, brine, dried over sodium sulfate, filtered and concentrated. residue was chromatographed on silica gel using a gradient of chloroform to 1% methanol/chloroform as 15 eluant to yield 1.4 g (67%) of the desired product as a tan foam: 1H-NMR is consistent with structure; MS (ion spray) 371.1 (M+1); Anal. Calc'd for $C_{19}H_{22}N_4O_4$: C, 61.61; H, 5.99; N, 15.13. Found: C, 61.95; H, 6.01; N, 14.95.

20

Preparation 321



Reaction of the product of Preparation 320 (0.87 g, 2.3 mmol), 10% palladium on carbon (2.0 g), tetrahydrofuran (80 mL), the product of Preparation 1d, (0.9 g, 2.3 mmol), 1-hydroxybenzotriazole (0.35 g, 2.53 mmol), dicyclohexylcarbodiimide (0.52 g, 2.53 mmol), as described in Preparation 314, gave 0.23 g (14%) of the

382

desired product as a tan foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 703.6 (M+1); Anal. Calc'd for $C_{38}H_{50}N_{6}O_{7}$ 0.15CHCl₃: C, 63.57; H 7.01; N, 11.66. Found: C, 63.60; H, 7.32; N, 11.31.

5

Example 170

To a solution of the product of Preparation 321, 0.21 g (0.3 mmol) in 6 mL of dichloromethane was added 2 mL of trifluoroacetic acid. The mixture was stirred 40 min at 10 ambient temperature then concentrated. The resulting residue was partitioned between ethyl acetate and saturated sodium bicarbonate and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and 15 concentrated. The crude product was chromatographed on silica gel using 10% methanol/chloroform as eluant. solution of the product in chloroform was added a saturates solution of hydrochloric acid in diethyl ether. 20 The slurry was concentrated to yield the desired product as a white solid: 1H-NMR is consistent with structure; MS (ion spray) 603.4 (M+1); Anal. Calc'd for $C_{33}H_{42}N_6O_5$ 2.3HCl: C, 57.73; H, 6.50; N, 12.24. Found: C, 57.80; H, 6.34; N, 12.15.

25

Preparation 322




383

Reaction of (1R, 2S)-(+)-1-amino-2-indanol (3.8 g, 26.0 mmol), sodium bicarbonate (2.2 g, 26.0 mmol), the product of Preparation 306. (4.0 g, 26.0 mmol), methanol (28 mL), water (12 mL), as in Preparation 313 gave 4.0 g (63%) of the desired product as a tan foam: $^1\text{H-NMR}$ is consistent with structure; MS (ion spray) 246.3 (M+1); Anal. Calc'd for $C_{12}H_{11}N_3O_3$: C, 58.77; H, 4.52; N, 17.13. Found: c, 58.68; H, 4.48; H, 17.00.

10


Preparation 323



Reaction of the product of Preparation 322 (2.2 g, 9.0 mmol), 1,1'-carbonyldiimidazole (1.73 g, 10.8 mmol), 4-methylpiperidine (5.3 mL, 45.0 mmol), dichloromethane (80 mL), as described in Preparation 320, gave 2.3 g (70%) of the desired product as an orange foam: ¹H-NMR is consistent with structure; MS (ion spray) 371.1 (M+1); Anal. Calc'd for C₁₉H₂₂N₄O₄'0.15CHCl₃: C, 59.23; H, 5.75; N, 14.43. Found: C, 59.38; H, 5.87; N, 14.55.

20

Preparation 324



Reaction of the product of Preparation 323 (2.22 g, 6.0 mmol), 10% palladium on carbon (3.0 g), tetrahydrofuran (60 mL), the product of Preparation 1d, (2.3 g, 6.0 mmol), 1-hydroxybenzotriazole (0.9 g, 6.6 mmol), dicyclohexylcarbodiimide (1.4 g, 6.6 mmol), as described in Preparation 314 gave 1.0 g (24%) of the desired

384

product as a tan foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 703.6 (M+1); Anal. Calc'd for $C_{38}H_{50}N_{6}O_{7}$: C, 64.94; H, 7.17; N, 11.96. Found: C, 64.81; H, 7.09; N, 11.83.

5

Example 171

Reaction of the product of Preparation 324 (0.97 g, 1.4 mmol), trifluoroacetic acid (4 mL), dichloromethane (12 mL), as in described in Example 170 gave 0.95 g (100%) of the desired product as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 603.3 (M+1); Anal. Calc'd for $C_{33}H_{42}N_6O_5$ 2.3HCl: C, 57.73; H, 6.50; N, 12.24. Found: C, 57.46; H, 6.53; N, 11.90

15

10

Preparation 325



Prepared as shown in Preparation 1 in Examples Part 4. ¹H-NMR consistent with structure.

20

Preparation 326

Reaction of the product of Preparation 4 from Examples
Part 4 (0.4 g, 1.4 mmol), 10% palladium on carbon (0.8
g), tetrahydrofuran (40 mL), the product of Preparation
1d, (0.53 g, 1.4 mmol), 1-hydroxybenzotriazole (0.21 g,

385

1.54 mmol), dicyclohexylcarbodiimide (0.32 g, 1.54 mmol), as described in Preparation 314 gave 0.25 g (29%) of the desired product as a tan foam: $^{1}\text{H-NMR}$ is consistent with structure; MS (ion spray) 620.7 (M+1); Anal. Calc'd for $C_{33}H_{41}N_{5}O_{7}$: C, 63.96; H, 6.67; N, 11.30. Found: C, 63.91; H, 6.72; N, 11.04.

Example 172

10 A solution of the product Preparation 326, 0.22 g (0.35 mmol) in 10 mL of a saturated solution of hydrochloric acid in glacial acetic acid was stirred for 2 h at ambient temperature and then concentrated. The residue was dissolved in toluene and concentrated, then trituated with hexanes to yield 0.187 g (89%) of the desired product as a tan solid: ¹H-NMR is consistent with structure; MS (ion spray) 520.3 (M+1); Anal. Calc'd for C28H33N5O5 2HCl: C, 56.76; H, 5.95; N, 11.82. Found: C, 56.96; H, 6.06; N, 11.83.

20

Preparation 327



To a solution of R-(-)-2-phenylglycinol, 2.75 g (20 mmol) in 20 mL of methanol and 10 mL of water at 0°C, was added 1.7 g (20 mmol) of sodium bicarbonate. The reaction mixture was stirred for 20 min at 0°C, then 3.16 g (20 mmol) of the product of Preparation 306 was added. The residue was absorbed onto silica and chromatographed on

386

silica gel using 80% ethyl acetate/hexanes as eluant to yield 3.5 g (74%) of the desired product as a colorless oil: 1H-NMR is consistent with structure; MS (ion spray) 234 (M+1).

5

10

Preparation 328



To a solution of the product of Preparation 327, 3.08 g (13.0 mmol), triethylamine, 2.4 mL (16.9 mmol) and 4dimethylaminopyridine, 0.3 g (2.6 mmol) in 150 mL of dichloromethane at 0°C was added 1.5 mL (15.6 mmol) of acetic anhydride dropwise. The reaction mixture was stirred overnight slowly warming to ambient temperature, then concentrated. The residue was partitioned between ethyl acetate and saturated sodium bicarbonate and 15 extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate and concentrated. The residue was chromatographed on silica gel using 80% ethyl acetate/hexanes as eluant to yield 3.4 g (95%) of the desired product as a tan oil: 1H-NMR is consistent with structure; MS (ion spray) 276.2 (95%); Anal. Calc'd for $C_{13}H_{13}N_3O_4$: C, 56.73; H, 4.76; N, 15.27. Found: C, 56.46; H, 4.89; N, 15.17.

25

20

To a slurry of 4.0 g of 10% palladium on carbon in 60 mL of tetrahydrofuran was added a solution of 3.1 g (11.3

mmol) of the product of Preparation 328 in 60 mL of tetrahydrofuran. The mixture was hydrogenated at 40 psi of hydrogen for 1.5 h and filtered through celite. To this solution was added 4.3 g (11.3 mmol) of the product of Preparation 1d, 1.7 g (12.4 mmol) of 1hydroxybenzotriazole and 2.6 g (12.4 mmol) of dicyclohexylcarbodiimide. The reaction was stirred overnight at ambient temperature, filtered and concentrated. The reaction mixture was stirred overnight, slowly warming to ambient temperature, then 10 was concentrated. The residue was dissolved in ethyl acetate, filtered, then water was added. The mixture was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium 15 sulfate, filtered and concentrated. The resulting residue was chromatographed on silica gel using 3% methanol/chloroform as eluant to yield 2.41 g (35%) of the desired product: 1H-NMR is consistent with structure; MS (ion spray) 608.3 (M+1); Anal. Calc'd for $C_{32}H_{41}N_5O_7$: C, 63.25; H, 6.80; N, 11.52. Found: C, 63.51; H, 6.79; N, 20 11.74.

Example 173

To a solution of the product of Preparation 329, 2.2 g
(3.6 mmol) in 25 mL of dichloromethane was added 8 mL of
trifluoroacetic acid. The reaction mixture was stirred
for 1 h, then concentrated. The residue was partitioned
between chloroform and saturated sodium carbonate and was
extracted with chloroform. The combined organic extracts
were washed with brine, dried over sodium sulfate,

388

filtered and concentrated. The residue was dissolved in ethyl acetate and a saturated solution of hydrochloric acid in diethyl ether was added. The mixture was concentrated to yield 1.7 g (81%) of the desired isomer as a white solid: ¹H-NMR is consistent with structure; MS (ion spray) 508.3 (M+1); Anal. Calc'd for C₂₇H₃₁N₅O₅ 2.3HCl: C, 55.00; H, 5.69; N, 11.89. Found: C, 55.06; H, 6.01; N, 11.41.

10

15

20

25

Preparation 330



To a solution of the product of Preparation 327, 3.45 g (15.0 mmol) in 100 mL of dichloromethane was added 3.0 g (18.0 mmol) of 1,1'-carbonyldiimidazole. The reaction mixture was stirred overnight at ambient temperature then 8.9 mL (75 mmol) of 4-methylpiperidine was added. reaction mixture was stirred overnight at ambient temperature then concentrated. The residue was partitioned between ethyl acetate and 0.1N aqueous hydrochloric acid and washed with 0.1N aqueous hydrochloric acid. The combined aqueous layers were extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. The residue was chromatographed on silica gel using 80% ethyl acetate/hexanes as eluant to yield 4.4 g (85%) of the desired product as a yellow oil: 1H-NMR is consistent with structure; MS (ion spray) 359.2 (M+1); Anal. Calc'd for $C_{18}H_{22}N_4O_4^{\circ}0.3CHCl_3$: C, 56.46; H, 5.78; N, 14.43.

30 Found: C, 56.81; H, 5.98; N, 14.42.

389

Pr paration 331

Reaction of the product of Preparation 330 (4.2 g, mmol), 10% palladium on carbon (5.2 g), tetrahydrofuran (80 mL), the product of Preparation 1d, (4.5 g, 11.7 mmol), 1-hydroxybenzotriazole (1.8 g, 12.9 mmol), dicyclohexylcarbodiimide (2.7 g, 12.9 mmol), as in described in Preparation 329 gave 2.2 g (27%) of the desired product as a tan foam: ¹H-NMR is consistent with structure; MS (ion spray) 691.3 (M+1); Anal. Calc'd for C₃₇H₅₀N₆O₇: C, 64.33; H, 7.30; N, 12.17. Found: C, 64.04; H, 7.41; N, 11.88.

Example 174

15

Reaction of the product of Preparation 331 (1.95 g, 2.8 mmol), trifluoroacetic acid (8 mL), dichloromethane (25 mL), as described in Example 173 gave 1.7 g (90%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (ion spray) 591.6 (M+1); Anal. Calc'd for C₃₂H₄₂N₆O₅ 2.4HCl: C, 56.67; H, 6.60; N, 12.39. Found: C, 56.94; H, 6.63; N, 12.04.

390

To a solution of the product of Preparation 8, Examples, Part 1, 2.5 g (4.1 mmol) in 12 mL of dichloromethane was added 4 mL of trifluoroacetic acid. The solution was stirred for 2 h and was then concentrated. The residue was partitioned between ethyl acetate and saturated sodium bicarbonate and was then extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated to yield 1.88 g (90%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (ion spray) 508.3 (M+1); Anal. Calc'd for C₂₇H₃₃N₅O₅·1H₂O: C, 61.70; H, 6.71; N, 13.32. Found: C, 61.79; H, 6.25; N, 13.07.

Example 176

To a solution of the product of Example 175 -438377, 1.66

20 g (3.3 mmol) in 50 mL of methanol was added 0.6 g of
lithium borohydride. The reaction mixture was stirred 10
min. then concentrated. The residue was partitioned
between ethyl acetate and water and was extracted with
ethyl acetate. The combined organic extracts were washed

25 with brine, dried over sodium sulfate, filtered and
concentrated. The desired product was recrystallized
from ethyl acetate/hexanes to yield 0.9 g (58%) of a
white solid: ¹H-NMR is consistent with structure; MS (ion

391

spray) 466.3 (M+1); Anal. Calc'd for $C_{25}H_{31}N_5O_4^{-1}0.3H_2O$: C, 63.76; H, 6.76; N, 14.87. Found: C, 63.87; H, 6.90; N, 14.77.

5

10

15

20

Preparation 332

To a solution of the product of Preparation 8 from Examples Part 1, 3.0 g (5.2 mmol), N-methylmorpholine, 0.63 mL (5.7 mmol) and 4-dimethylaminopyridine, 20 mg in 30 mL of dichloromethane at 0°C was added 0.55 mL (5.7 mmol) of ethyl chloroformate. After stirring at 0°C for 1 h, the reaction mixture was concentrated to dryness. The residue was partitioned between ethyl acetate and 1N HCl and was extracted with ethyl acetate. The combined organics were washed with 1N HCl, brine, dried over sodium sulfate, filtered and concentrated to dryness. The crude material was chromatographed on silica gel using 3% methanol/chloroform as eluant to yield 2.7 g (85%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (ion spray) 608.1 (M+1); Anal. Calc'd for C₃₂H₄₁N₅O₇: C, 63.

Preparation 333

Reaction of the product of Preparation 8 (0.4 g, 0.69 mmol), N-methylmorpholine (0.08 mL, 0.69 mmol), isobutyl chloroformate (0.1 mL, 0.76 mmol) in dichloromethane (6 mL), as described in Preparation 332, gave 0.37 g (84%)

392

of the desired product as a tan foam: ^{1}H -NMR is consistent with structure; MS (FD) 635 (M+); Anal. Calc'd for $C_{34}H_{45}N_{5}O_{7}$: C, 64.23; H, 7.13; N, 11.02. Found: C, 64.15; H, 7.02; N, 10.94.

5

Example 177

Reaction of the product of Preparation 333 (0.31 g, 0.5 mmol), trifluoroacetic acid (2 mL), dichloromethane (6 mL), as described in Preparation 175, gave 0.25 g (83%) of the desired product as a tan solid: ¹H-NMR is consistent with structure; MS (FD) 535 (M+).

Preparation 334

15

20

Reaction of the product of Preparation 8 from Examples Part 1 (0.6 g, 1.0 mmol), N-methylmorpholine (0.1 mL, 1.0 mmol), phenyl chloroformate (0.14 mL, 1.1 mmol), dichloromethane (6 mL), as described in Preparation 332 gave 0.4 g (61%) of the desired mixture of isomers as a tan foam: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 655.3 (M+); Anal. Calc'd for $C_{36}H_{41}N_5O_7$: C, 65.94; H, 6.30; N, 10.68. Found: C, 65.64; H, 6.42; N, 10.43.

393

Example 178

Reaction of the product of Preparation 334 (0.37 g, 0.56 mmol), triflouroacetic acid (2 mL), dichloromethane (6 mL), as in described in Example 175 gave 0.32 g (91%) of the desired product as a white solid: $^{1}\text{H-NMR}$ is consistent with structure; MS (FD) 555 (M+); Anal. Calc'd for $C_{31}H_{33}N_{5}O_{5}$ 2HCl: C, 59.24; H, 5.61; N, 11.14. Found: C, 59.30; H, 5.88; N, 10.97.

10

Preparation 335

Reaction of the product of Preparation 8 from Examples

Part 1 (0.4 g, 0.69 mmol), N-methylmorpholine (0.08 mL, 0.69 mmol), benzyl chloroformate (0.11 mL, 0.76 mmol), dichloromethane (6 mL), as described in Preparation 332 gave 0.32 g (70%) of the desired product as a white foam: ¹H-NMR is consistent with structure; MS (FD) 669

(M+).

Example 179

Reaction of the product of Preparation 335 (0.27 g, 0.4 mmol), trifluoroacetic acid (2 mL), dichloromethane (6

394

mL), as described in Preparation 175, gave 0.24 g (92%) of the desired product as a tan solid: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 569 (M+); Anal. Calc'd for $\text{C}_{32}\text{H}_{35}\text{N}_{5}\text{O}_{5}$ 2HCl: C, 59.81; H, 5.80; N, 10.90. Found: C, 60.05; H, 6.07; N, 10.66.

Preparation 336




To a suspension of 60% sodium hydride in mineral oil

(26.0 g , 0.6 mol) stirring in dimethylformamide (500 mL)

at -5 °C was slowly added 4-nitroimidazole (73.5 g, 0.650 mol). After 30 min, ethyl bromoacetate (100 g, 0.599 mol) was added neat over 15 minutes via addition funnel. After 24 h, the solution was concentrated, diluted with ethyl acetate washed with water. The organic extract was dried over sodium sulfate and concentrated then triturated in ether to collect 67.1 g (56%) of the desired product as pale yellow needles: ¹H NMR consistent with structure; MS FD+ = 199; Anal. Calcd. for C7H9N3O4:

C, 42.21; H, 4.55; N, 21.20. (Found) C, 43.90; H, 4.58; N, 22.02.

Preparation 337



25

5

The product of Preparation 336 was reduced and coupled to give the ester which was then dissolved in methanol (400ml) at room temp. 1 M lithium hydroxide (45.2 ml, 45.3 mol) was added drop-wise over 10 minutes. After 20

395

min, the mixture was concentrated then diluted with 150 mL 10% acetic acid/ethyl acetate and concentrated and extracted with ethyl acetate. The combined organic extracts were dried over sodium sulfate, and concentrated to give 9.26 g (81%) of the desired material as a white amorphous powder: ^{1}H NMR consistent with structure; MS FD+ = 504; Anal. Calcd. For $C_{24}H_{32}N_{5}O_{7}$: C, 57.25; H, 6.61; N, 13.91. (Found) C, 57.04; H, 6.83; N, 14.07.

10 Preparation 338

To a solution of the product of Preparation 337 (250 mgs, 0.49 mmol) and 1-hydroxybenzotriazole (67 mgs, 0.5 mmol) stirring in dry dioxane (5 mL) was added

15 dicyclohexylcarbodiimide (107 mg, 0.521). After 15 min, piperidine(49 mg, 0.5 mmol) was added neat. After 18 h, the reaction mixture was filtered and concentrated. Purification by via radial chromatography (4 mm silica gel plate, 6% methanol/methylene chloride) gave 105 mg

20 (36%) of a white amorphous foam: used in the next reaction without further purification.

Example 180


25

To a solution of the product of Preparation 338 (105 mg, 0.18 mmol)stirring in dichloromethane (5 mL) at room temperature was added anisole 0.2 mL) followed by

396

trifluoroacetic acid (2 mL). After 2 h, the reaction mixture was concentrated and triturated with diethyl ether to give 86 mg (67%) of desired product as a white amorphous solid: ^{1}H NMR consistent with structure; MS FD+ = 484; Anal. Calcd. for $C_{25}H_{36}N_{6}O_{4}$ X 2 trifluoroacetic acid: C, 48.87; H, 5.38; N, 11.79. (Found) C, 48.71; H, 5.30; N, 11.59.

Example 181




10

15

Reaction of the product of Preparation 337 (250 mg, 0.5 mmol) and 3-methyl piperidine (49 mg, 0.5 mmol) as described in Preparation 338 gave the corresponding amide which deprotected according to Example 180 to provide the desired product as a white amorphous solid: ^{1}H NMR consistent with structure; MS FD+ = 485; Anal. Calcd. for $C_{25}H_{36}N_6O_4$ X 2 trifluoroacetic acid: C, 48.87; H, 5.38; N, 11.79. (Found) C, 48.61; H, 5.25; N, 11.61.

20

Example 182




Reaction of the product of Preparation 337 (250 mg, 0.5 mmol) and 2-methyl piperidine (49 mg, 0.5 mmol) as

described in Preparation 338 gave the corresponding amide which deprotected according to Example 180 to provide the desired product as a white amorphous solid: ¹H NMR consistent with structure; MS FD+ = 485; Anal. Calcd.

397

for $C_{25}H_{36}N_6O_4$ X 2 trifluoroacetic acid: C, 48.87; H, 5.38; N, 11.79. (Found) C, 48.65; H, 5.39; N, 11.64.

Example 183




5

Reaction of the product of Preparation 337 (250 mg, 0.5 mmol) and 2-methyl pyrrolidine (42 mg, 0.5 mmol) as described in Preparation 338 gave the corresponding amide which deprotected according to Example 180 to provide the desired product as a white amorphous solid: ^{1}H NMR consistent with structure; MS FD+ = 470; Anal. Calcd. for $C_{24}H_{34}N_{6}O_{4}$ X 2 trifluoroacetic acid: C, 48.14; H, 5.19; N, 12.03. (Found) C, 47.21; H, 5.37; N, 11.73.

15

10

Example 184




Reaction of the product of Preparation 337 (250 mg, 0.5 mmol) and 3-methyl pyrrolidine (42 mg, 0.5 mmol) as described in Preparation 338 gave the corresponding amide which deprotected according to Example 180 to provide the desired product as a white amorphous solid: ^{1}H NMR consistent with structure; MS IS+ = 471; Anal. Calcd. for $C_{24}H_{34}N_{6}O_{4}$ X 2 trifluoroacetic acid: C, 48.14; H, 5.19; N, 12.03. (Found) C, 48.02; H, 5.11; N, 11.83.

25

20

Example 185

398




Reaction of the product of Preparation 337 (250 mg, 0.5 mmol) and 4-propylpiperidine (63 mg, 0.5 mmol) as described in Preparation 338 gave the corresponding amide which deprotected according to Example 180 to provide the desired product as an off-white solid: 1H NMR consistent with structure; MS IS+ = 513; Anal. Calcd. for $C_{27}H_{40}N_6O_4$ X 2 trifluoroacetic acid: C, 50.27; H, 5.72; N, 11.35. (Found) C, 52.30; H, 6.37; N, 12.29.

10

15

20

Example 186



Reaction of the product of Preparation 337 (250 mg, 0.5 mmol) and piperidine (49 mg, 0.5 mmol) as described in Preparation 338 gave the corresponding amide which deprotected according to Example 180 to provide the desired product as an off-white amorphous solid: ^{1}H NMR consistent with structure; MS FD+ = 546; Anal. Calcd. for $C_{30}H_{38}N_{6}O_{4}$ X 2 trifluoroacetic acid: C,48.14; H,5.19; N, 12.03. (Found) C, 48.02; H, 5.11; N, 11.83.

Example 187

Reaction of the product of Preparation 337 (1.75 g, 3.48 mmol), dicyclohexylcarbodiimide (717 mg, 3.48 mmol), 1-

399

hydroxybenzotriazole (470 mg, 3.48 mmol), and pyrrolidine (0.29 mL, 3.48 mmol) as described in Preparation 338 gave 870 mg (45%) of the corresponding amide which was deprotected according to Example 180 using anisole (1 mL) and trifluoroacetic acid (10 mL) to yield 580 mg (54%) of final product as a tan amorphous solid: 1 H NMR consistent with structure; MS IS+ = 457; Anal. Calcd. for $C_{23}H_{32}N_6O_4$ X 2 trifluoroacetic acid: C, 47.37; H, 5.01; N, 12.28. (Found) C, 51.40; H, 5.81; N, 14.34.


10

15

20

25

Example 188



Reaction of the product of Preparation 337 (1.7 g, 3.38 mmol), dicyclohexylcarbodiimide (697 mg, 3.38 mmol), 1-hydroxybenzotriazole (456 mg, 3.38 mmol), 4-(p-fluorobenzoyl) piperidine hydrochloride (823 mg, 3.38 mmol), and triethylamine (0.47 mL, 3.38 mmol) as described in Preparation 338 gave 1.96 g (84%) of the corresponding amide which was deprotected according to Example 180 using anisole (1 mL) and trifluoroacetic acid (10 mL) to yield 1.28 g (55%) of final product as an off-white powder: ¹H NMR consistent with structure; MS IS+ = 593; Anal. Calcd. for C₃₁H₃₇FN₆O₅ X 2 trifluoroacetic acid: C, 51.22; H, 4.79; N, 10.24. (Found) C, 52.71; H, 5.07; N, 10.92.

Example 189

400

Reaction of the product of Preparation 337(400 mg, 0.79 mmol), dicyclohexylcarbodiimide (164 mg, 0.79 mmol), 1-hydroxybenzotriazole (107 mg, 0.79 mmol), and cyclohexylamine (78 mg, 0.79 mmol) as described in

5 Preparation 338 gave 234 mg (50%) of the corresponding amide which was deprotected according to Example 180 using anisole (0.05 mL) and trifluoroacetic acid (4 mL) to yield 220 mg (77%) of final product as a white amorphous solid: ¹H NMR consistent with structure; MS

10 IS+ = 485; Anal. Calcd. for C₂₅H₃₆N₆O₄ X 2 trifluoroacetic acid: C, 48.88; H, 5.38; N, 11.79. (Found) C, 48.70; H, 5.52; N, 11.81.

Example 190

15

30

Reaction of the product of Preparation 337(400 mg, 0.79 mmol), dicyclohexylcarbodiimide (164 mg, 0.79 mmol), 1-hydroxybenzotriazole (107 mg, 0.79 mmol), and 4
20 benzylpiperidine (139 mg, 0.79 mmol) as described in Preparation 338 gave 196 mg (68%) of the corresponding amide which was deprotected according to Example 180 using anisole (0.05 mL) and trifluoroacetic acid (4 mL) to yield 196 mg(68%) of final product as a white

25 amorphous solid: ¹H NMR consistent with structure; MS IS+ = 561; Anal. Calcd. for C₃₁H₄₀N₆O₄ X 2 trifluoroacetic acid: C, 52.37; H, 5.02; N, 10.47. (Found) C, 52.55; H, 5.33; N, 10.56.

Example 191

401

Reaction of the product of Preparation 337 (400 mg, 0.79 mmol), dicyclohexylcarbodiimide (164 mg, 0.79 mmol), 1- hydroxybenzotriazole (107 mg, 0.79 mmol), and 4- ethylpiperidine (89 mg, 0.79 mmol) as described in Preparation 338 gave 236 mg (50%) of the corresponding amide which was deprotected according to Example 180 using anisole (0.05 mL) and trifluoroacetic acid (4 mL) to yield 204 mg (71%) of final product as a white

10 amorphous solid: ¹H NMR consistent with structure; MS FD+ = 498; Anal. Calcd. for C₂₆H₃₈N₆O₄ X 2 trifluoroacetic acid: C, 49.59; H, 5.55; N, 11.57. (Found) C, 48.77; H, 5.29; N, 11.37.

15

20

25

Example 192


Reaction of the product of Preparation 337 (400 mg, 0.79 mmol), dicyclohexylcarbodiimide (164 mg, 0.79 mmol), 1-hydroxybenzotriazole (107 mg, 0.79 mmol), 4-benzoylpiperidine hydrochloride (179 mg, 0.79 mmol), and triethylamine (0.11 mL, 0.79 mmol) as described in Preparation 338 gave 175 mg (33%) of the corresponding amide which was deprotected according to Example 180 using anisole (0.05 mL) and trifluoroacetic acid (4 mL) to yield 170 mg (82%) of final product as a white amorphous solid: 1 H NMR consistent with structure; MS IS+ = 575; Anal. Calcd. for $C_{31}H_{38}N_{6}O_{5}$ X 2 trifluoroacetic

402

acid: C, 52.37; H, 4.99; N, 10.47. (Found) C, 52.13; H, 5.03; N, 10.62.

403

Example 193




Reaction of the product of Preparation 337(400 mg, 0.79 mmol), dicyclohexylcarbodiimide (164 mg, 0.79 mmol), 1
5 hydroxybenzotriazole (107 mg, 0.79 mmol), and 4-tbutylpiperidine (112 mg, 0.79 mmol) as described in
Preparation 338 gave 342 mg (69%) of the corresponding
amide which was deprotected according to Example 180
using anisole (0.05 mL) and trifluoroacetic acid (4 mL)

10 to yield 287 mg (70%) of final product as a white
amorphous solid: ¹H NMR consistent with structure; MS
IS+ = 527; Anal. Calcd. for C28H42N6O4 X 2 trifluoroacetic
acid: C, 50.93; H, 5.84; N, 11.14. (Found) C, 50.65;
H, 5.99; N, 11.28.

15

Example 194




Reaction of the product of Preparation 337 (400 mg, 0.79 mmol), dicyclohexylcarbodiimide (164 mg, 0.79 mmol), 1
20 hydroxybenzotriazole (107 mg, 0.79 mmol), and 4isopropylpiperidine (101 mg, 0.79 mmol)) as described in Preparation 338 gave 208 mg (43%) of the corresponding amide which was deprotected according to Example 180 using anisole (0.05 mL) and trifluoroacetic acid (4 mL)

25 to yield 197 mg (78%) of final product as a white amorphous solid: ¹H NMR consistent with structure; MS IS+ = 513; Anal. Calcd. for C₂₇H₄₀N₆O₄ X 2 trifluoroacetic acid: C, 50.27; H, 5.68; N, 11.35. (Found) C, 50.21; H, 5.78; N, 11.64.

404


Preparation 339



To a solution of the product of Preparation 337 (400 mg, 0.79 mmol) and N-methyl morpholine (0.1 mL) stirring in dichloromethane (20 mL) at 0°C was added isobutyl chloroformate (0.13 mL, 1 mmol). After 15 min, benzylamine (0.11 mL, 1 mmol) was added drop-wise and after 30 min the reaction mixture was quenched with water and washed with saturated sodium bicarbonate solution, 10 water, and 1N HCl. The organic extract was then dried over sodium sulfate, concentrated and then purified by flash chromatography (silica gel, 5% methanol/dichloromethane) to yield 199 (42%) mg of the 15 desired product as a yellow foam: ¹H NMR consistent with structure; MS FD+ = 592; Anal. Calcd. for $C_{31}H_{40}N_6O_6$: C, 62.82; H, 6.80; N, 14.18. (Found) C, 61.82; H, 6.72; N, 13.75.

20

Example 195



To a solution of the product of Preparation 339 (183 mg, 0.31) stirring in dichloromethane (5 mL) at room temperature was added anisole (0.2 mL) followed by 25 trifluoroacetic acid (2 mL). After 2 h, the reaction mixture was concentrated and subsequently triturated with diethyl ether to provide 93 mg (42%) of desired product as a white amorphous solid: ¹H NMR consistent with structure; MS FD+ = 492; Anal. Calcd. for C₂₆H₃₂N₆O₄

405

x 2trifluoroacetic acid: C, 50.00; H, 4.76; N, 11.66. (Found) C, 50.28; H, 4.93; N, 11.59.

Preparation 340



5

10


15

To a mixture of potassium t-butoxide (16.2 g, 145 mmol) stirring in dimethylformamide (100 mL) was added 4-nitroimidazole (13.6 g, 120.7 mmol) portion-wise over 30 min. After 1 h, ethyl-1-bromocyclobutanecarboxylate (25 g, 120.7 mmol) was slowly added. The mixture was heated to 100 0°C, and after 36 h, the mixture was cooled to room temperature and concentrated. The resulting solid was triturated in diethyl ether and filtered. The filtrate was concentrated and purified by flash chromatography (silica gel; 60% ethyl acetate/hexanes) to yield 3.86 g (13%) of the desired product as a clear yellow oil: 1 H NMR consistent with structure; MS FD+ = 239; Anal. Calcd. for $C_{10}H_{13}N_{3}O_{4}$: C_{7} 50.21; H_{7} 5.48; N_{7} 17.56; (Found) C_{7} 50.36; H_{7} 5.60; N_{7} 17.54.

20

25

Preparation 341




To a solution of the product of Preparation 340 (3.6 g, mmol) in dioxane (50 mL) was added 5% palladium on carbon (0.9 g). The resulting mixture was hydrogenated at room temperature at 40 p.s.i on a Parr apparatus for 30 min. This mixture was then filtered through celite and added to solution of the product of Preparation 1d (5.73 g, 15.05 mmol), 1-hydroxybenzotriazole (2.03 g, 15.05 mmol),

406

and dicyclohexylcarbodiimide (3.11 g, 15.05 mmol) in dioxane (75 mL). After 18 h, the reaction was filtered and the filtrate concentrated. The resulting orange foam was purified by flash chromatography (silica gel; methanol /dichloromethane) to yield 2.63 g of the desired product as a yellow foam: ¹H NMR consistent with structure; MS IS+ = 572.

Preparation 342




10

To a solution of the product of Preparation 341 (2.6 g, 4.55 mmol) stirring in methanol (80 mL) at room temperature was slowly added 1M LiOH (9.1 mL, 9.1 mmol) drop-wise. After 90 min, the reaction mixture was concentrated then diluted with 30 mL 10% acetic acid/ethyl acetate and concentrated. The resulting concentrate was diluted with 50 mL 10% acetic acid/ethyl acetate and washed with water. The organic extract was dried over sodium sulfate and then concentrated.

20 Triturate the solid in diethyl ether gave 620 mg (25%) of desired product as a white amorphous powder: ¹H NMR

Preparation 343

consistent with structure; MS IS+ = 544.



25

To a solution of the product of Preparation 342 (600 mg, 1.10 mmol) and 1-hydroxybenzotriazole (149 mg, 1.10 mmol)stirring in dioxane (10 mL) at room temperature was added dicyclohexylcarbodiimide (228 mg, 1.10 mmol).

407


After 15 min, 4-methylpiperidine (109 mg, 1.10 mmol) was added. After 18 h, the mixture was filtered and the filtrate concentrated. The resulting material was purified by radial chromatography to yield 200 mg (29%)of the desired product as a white amorphous foam: ¹H NMR consistent with structure; MS IS+ = 625.

Example 196

To a solution of the product of Preparation 343 (200 mg, 0.32 mmol) stirring in dichloromethane (5 mL) at room temperature was added anisole (0.05 mL) followed by trifluoroacetic acid (2 mL). After 2 h, the reaction mixture was concentrated then triturate in diethyl ether to provide 118 mg (49%) of desired product as a white amorphous solid: ¹H NMR consistent with structure; MS IS+ = 525; Anal. Calcd. for C28H40N6O4 x 2trifluoroacetic acid: C, 51.06; H, 5.62; N, 11.16. (Found) C, 53.40; H, 6.06; N, 11.99.

20

Preparation 344



To a solution of the product of Preparation 5 (10.0 g, 16.4 mmol) stirring in tetrahydrofuran (150 mL) at room temperature was added 1-hydroxybenzotriazole (2.22 g, 16.4 mmol) and 1,3- dicyclohexylcarbodiimide (3.38 g, 16.4 mmol). After 15 min, pyrrolidine (1.37 mL, 16.4 mmol) was added. After 16 h, the reaction mixture was

408

filtered and concentrated. The resulting crude material was purified by flash chromatography (silica gel, 5% methanol/dichloromethane) to yield 7.05 g (65%) of the desired product as a yellow foam: $^{1}\text{H-NMR}$ consistent with product; MS (ion spray) 663 (M+1); Anal. Calcd for $C_{35}H_{46}N_{6}O_{7}$: C, 63.43; H, 7.00; N, 12.68. (Found) C, 62.69; H, 6.87; N, 12.91.

Examples 197 and 198

10

To the product of Preparation 344 (7.0 g, 10.6 mmol) was added a saturated solution of HCl(g)/acetic acid (100 mL). After 4 h, the reaction mixture was concentrated then partitioned between ethyl acetate and saturated 15 sodium bicarbonate. The organic layer was removed, dried over sodium sulfate and concentrated to yield 5.59 g (94%) of the free base as a light yellow foam. diastereomeric material (3.45 g) was chromatographed on an 8 x 15 cm Prochrom column packed with Kromsil CHI-DIMETHYLFORMAMIDE chiral phase using an eluent mixture of 20 3A alcohol and dimethylethylamine in heptane to provide the individual diastereomers in pure form: 1H NMR consistent with product; MS (ion spray) 563 (M+1); Anal. Calcd. for $C_{30}H_{38}N_6O_5$: C, 64.04; H, 6.81; N, 25 14.94. (Found) C, 63.98; H, 6.82; N, 14.87. Example 197. (Isomer 1) To a solution of the purified isomer in ethyl acetate was added a saturated solution of hydrochloric acid in ethyl acetate. The resulting slurry was concentrated to dryness to yield 1.50 g (39%) of the desired product as an off-white solid: 1H NMR consistent with product; MS (ion spray) 563 (M+1); Anal. Calcd.

409

for $C_{30}H_{38}N_6O_5 \times 2$ HCl: C, 56.69; H, 6.34; N, 13.22. (Found) C, 55.81; H, 6.40; N, 12.68.

Example 198. (Isomer 2) To a solution of the purified isomer in ethyl acetate was added a saturated solution of hydrochloric acid in ethyl acetate. The resulting slurry was concentrated to dryness to yield 1.43 g (38%) of the desired product as an off-white solid: 1 H NMR consistent with structure; MS (ion spray) 563 (M+1); Anal. Calcd. for $C_{30}H_{38}N_{6}O_{5} \times 2$ HCl: C, 56.69; H, 6.34; N, 13.22. (Found) C, 55.71; H, 6.38; N, 12.74.

Preparation 345

10

To a mixture of the product of Preparation 99 (6.0 g, 17.1 mmol) and 10% palladium on carbon (6.0 g) in 15 tetrahydrofuran (100 mL). The reaction mixture was placed under a hydrogen atmosphere (40 psi) using a Parr apparatus for 30 min then filtered through Celite. The resulting solution was then added to a previously 20 prepared mixture of the product of Preparation 1L (6.66 g, 17.1 mmol), 1-hydroxybenzotriazole (2.31 g, 17.1 mmol), and 1,3 dicyclohexylcarbodiimide (3.53 g, 17.1 mmol) in tetrahydrofuran (75 mL). After 16 h at room temperature, the reaction mixture was concentrated and 25 the crude material purified by flash chromatography (silica gel, 4% methanol/dichloromethane) to yield 6.17 g (52%) of the desired product as a brown foam: ¹H NMR consistent with structure; MS (ion spray) 693 (M+1).

10

410

Preparation 346

To a solution of the product Preparation 345 (4.6 g, 6.64 mmol) stirring in tetrahydrofuran (100 mL) at room temperature was added a solution of lithium hydroxide in water (40 mL of 1M). After 30 min, the reaction mixture was acidified with 5N HCl (8.5 mL). The resulting mixture diluted with water and extraced with ethyl acetate. The combined organic extracts were dried over sodium sulfate and concentrated to yield 4.4 g (99%) of the desired product as a yellow foam.


Preparation 347

To a solution of the product Preparation 346 (4.0 g, 6.02 mmol) stirring in tetrahydrofuran (50 mL) at room temperature was added 1-hydroxybenzotriazole (813 mg, 6.02 mmol) and 1,3 dicyclohexylcarbodiimide (1.24 g, 6.02 mmol). After 15 min, dimethylamine (3.0 mL of a 2M soln in tetrahydrofuran, 6.02 mmol) was added. After stirring for 16 h in a sealed flask, the reaction mixture was filtered and concentrated. The resulting crude material was purified by flash chromatography (silica gel, 5% methanol/dichloromethane) to yield 2.79 g (68%) of the desired product as a yellow foam..

PCT/US98/17201

411

Examples 199 and 200



To the product of Preparation 347 (3.4 g, 5.0 mmol) was added a saturated solution of HCl(g)/acetic acid (50 mL).

After 1.5 h, the reaction mixture was concentrated then partitioned between ethyl acetate and saturated sodium bicarbonate. The organic layer was removed, dried over sodium sulfate and concentrated to yield 2.45 g (84%) of the free base as a light yellow foam. The diastereomeric

10 material (2.45 g) was chromatographed on an 8 x 15 cm Prochrom column packed with Kromsil CHI chiral phase using an eluent mixture of 3A alcohol and dimethylethylamine in heptane to provide the individual diastereomers in pure form: ¹H NMR consistent with

product; MS (ion spray) 592 (M+1); Anal. Calcd. for 15 $C_{34}H_{37}N_7O_3$: C, 69.02; H, 6.30; N, 16.57. (Found) C, 67.93; H, 6.29; N, 15.80.

Example 199. (Isomer 1) To a solution of the purified isomer in ethyl acetate was added a saturated solution of hydrochloric acid in ethyl acetate. The resulting slurry was concentrated to dryness to yield 992 mg (37%) of the desired product as an off-white solid: 1H NMR consistent with product; MS (ion spray) 592 (M+1); Anal. Calcd. for $C_{34}H_{37}N_{7}O_{3} \times 2$ HCl: C, 61.44; H, 5.91; N, 14.75.

25 (Found) C, 59.54; H, 5.92; N, 13.76.

20

30

Example 200. (Isomer 2) To a solution of the purified isomer in ethyl acetate was added a saturated solution of hydrochloric acid in ethyl acetate. The resulting slurry was concentrated to dryness to yield 1.17 g (40%) of the desired product as an off-white solid: 1H NMR consistent

412

with structure; MS (ion spray) 592 (M+1); Anal. Calcd. for $C_{34}H_{37}N_7O_3 \times 2$ HCl: C, 61.44; H, 5.91; N, 14.75. (Found) C, 59.03; H, 6.04; N, 13.84.

5

10

15


20

Preparation 348



To a suspension of sodium hydride (1.8 g of a 60 % dispersion in mineral oil, 45 mmol) stirring at 0 oC in diemthylformamide (50 mL) was slowly added 4-nitroimidazole (5.0 g, 44.2 mmol). After 1 h, a solution of 4-fluorobenzyl bromide (5.5 mL, 44.2 mmol) in dimethylformamide (10 mL) was added dropwise over 10 min. After 16 h, the reaction mixture was warmed to room temperature and concentrated. The resultant crude material was then diluted with hot methanol, filtered, allowed to cool, then filtered to yield 5.0 g (51%) of pale yellow crystals: ¹H NMR consistent with structure; MS (ion spray) 222 (M+1); Anal. Calcd. for C₁₀H₈FN₃O₂: C, 54.31; H, 3.65; N, 19.00. Found: C, 53.81; H, 3.92; N, 17.48.

Preparation 349




To a slurry of Raney Nickel (116 mg) in dioxane (15 mL)

was added a slurry of the product of Preparation 348 (582 mg, 2.63 mmol) in dioxane (20 mL). The reaction mixture was placed under a hydrogen atmosphere (60 psi) using a Parr apparatus. After 30 min, the solution was filtered

through Celite and the filtrate immediately added to a previously prepared mixture of the product of Preparation 1d (1.0 g, 2.63 mmol), 1-hydroxybenzotriazole (355 mg, 2.63 mmol), and 1,3 dicyclohexylcarbodiimide (543 mg, 2.63 mmol) in dioxane (25 mL). After 16 h, the reaction mixture was concentrated and the crude material purified by flash chromatography (silica gel, 5% methanol/dichloromethane) to give the desired product. This material was then crystallized from diethyl ether to yield 150 mg (10%) of the desired product as a white powder: ¹H NMR consistent with structure.

Example 201



To the product of Preparation 349 (150 mg, 0.27 mmol) stirring in dichloromethane (5 mL) at room temperature was added anisole (0.05 mL) and trifluoroacetic acid (2 mL). After 3 h, the mixture was concentrated, triturated extensively in diethyl ether, and the product collected via filtration to provide 101 mg (55%) of desired product as a white powder: ¹H NMR consistent with structure; MS (ion spray) 454 (M+1); Anal. Calcd. For C₂₄H₂₈FN₅O₃ x 2TFA: C, 49.34; H, 4.44; N, 10.28. Found: C, 49.64; H, 4.58; N, 10.36.

25

Preparation 350



To a suspension of sodium hydride (1.56 g of a 60 % dispersion in mineral oil, 39 mmol) stirring in

PCT/US98/17201 WO 99/08697

414

DIMETHYLFORMAMIDE (50 mL) at o oC was slowly added 4nitroimidazole (4.0 g, 35.4 mmol). After 1 h, a solution of 4-trifluoro-methoxybenzyl bromide 9.03 g, 35.4 mmol) in dimethylformamide (10 mL) was added dropwise over 10 min. After 16 h, the reaction mixture was warmed to room temperature then concentrated. The resultant crude material was then purified by flash chromatography(silica gel, 100% dichloromethane) and the product subsequently crystallized from diethyl ether to yield 3.95 g (39%) of desired product as white needles: 1H NMR consistent with structure; MS (ion spray) 288 (M+1); Anal. Calcd. for $C_{11}H_8F_3N_3O_3$: C, 45.99; H, 2.79; N, 14.63; F, 19.86. (Found) C, 46.27; H, 2.86; N, 14.86; F, 19.74.

10

15


25

Preparation 351

To a slurry of Raney Nickel (150 mg) in dioxane (15 mL) was added a slurry of the product of Preparation 350 (755 mg, 2.63 mmol) in dioxane (20 mL). The reaction mixture was placed under a hydrogen atmosphere (60 psi) using a 20 Parr apparatus. After 30 min, the solution was filtered through Celite and the filtrate immediately added to a previously prepared mixture of the product of Preparation 1d (1.0 g, 2.63 mmol), 1-hydroxybenzotriazole (355 mg, 2.63 mmol), and 1,3 dicyclohexylcarbodiimide (543 mg, 2.63 mmol) in dioxane (25 mL). After 16 h, the reaction mixture was concentrated and the crude material purified by flash chromatography (silica gel, 5% methanol/ dichloromethane) to provide 171 mg (10%) of the desired 30 product as a white foam: ¹H NMR consistent with structure; used without further characterization.

415

Example 202



To the product of Preparation 351 (171 mg, 0.28 mmol)

5 stirring in dichloromethane (5 mL) at room temperature was added anisole (0.05 mL) and trifluoroacetic acid (2 mL). After 3 h, the mixture was concentrated, triturated extensively in diethyl ether, and the product collected via filtration to provide 84 mg (40%) of desired product

10 as a white powder: ¹H NMR consistent with structure; MS (ion spray) 520 (M+1); Anal. Calcd. For C₂₅H₂₈F₃N₅O₄ x 2

TFA: C, 46.59; H, 4.04; N, 9.37. Found: C, 48.48; H, 4.48; N, 10.04.

15 Preparation 35

20

25

A cold (0-5 °C) solution of 5.0 g (24.37 mmol) of Boc-Dserine in 20 mL of dry N,N-dimethylformamide was added to a stirred suspension of sodium hydride (60% dispersion in mineral oil, 2.2 g, 55 mmol) in 60 mL of dry dimethylformamide at 0-5 ℃ under nitrogen over 5 min. The resulting light yellow brown suspension was stirred at 0-5 $^{\circ}$ C until gas evaluation nearly ceased (~15 min). Then a cold (0-5 $^{\circ}C)$ solution of 4-fluorobenzyl bromide (3.12 mL, 25 mmol) in 10 mL of dry N, N-dimethylformamide (Additional 10 mL of dimethylformamide was used as washings) was added to the stirred mixture over 5 min. The resulting off white suspension was stirred at 0-5 ℃ for 2 h, during which time the mixture turned to a thick

suspension and the magnetic stirring became ineffective. The reaction flask was shaken by hand several times. The cooling bath was removed and the mixture was allowed to warm to room temperature while shaking the reaction flask occasionally by hand. By the time the mixture reached the room temperature, the suspension thinned and the magnetic stirring became effective. The mixture was stirred at ambient temperature for additional 1 h (total 5 hours). The mixture was concentrated and the oily residue was dissolved in 150 mL of water and extracted with ether . The aqueous layer was cooled to 0-5 °C, acidified to pH ~3 with cold (0-5 °C) 0.5N HCl, and quickly extracted with ethyl acetate . The aqueous layer was saturated with sodium chloride and extracted with ethyl acetate . The ethyl acetate extracts were combined, washed with brine , dried over anhydrous sodium sulfate, filtered, and concentrated to dryness to give 6.86 g (89.9%) of the desired product as a light yellow viscous oil which was used in the next step without further purification: $^{1}H-NMR$ (δ , DMSO)1.38 (s, 9H), 3.60-3.70 (m, 2H), 4.15-4.25 (m, 1H), 4.45 (s, 2H), 6.97 (d, J = 8.3Hz, 1H), 7.16 (t, J = 8.7 Hz and 9.0 Hz, 2H), 7.30-7.40(m, 2H), 12.65 (br s, 1H); MS (ion spray) 314 (M+1).

10

15

20

25

30

Preparation 353



Trifluoroacetic acid (40 mL) was added to a stirred solution of 6.5 g (20.75 mmol) of the product of Preparation 352 in 40 mL of dichloromethane and the mixture was stirred at ambient temperature for 1 h, then it was concentrated to dryness. The resulting oily residue was dissolved in 30 mL of deionized water, cooled

PCT/US98/17201 WO 99/08697

417

ice-water bath, neutralized to pH ~7, and the separated white precipitate was filtered. The precipitate was washed with ether (5 \times 10 mL) to remove the yellow brown impurity and the resulting white precipitate was dried to a constant weight to give 2.45 g (55.4%) of the desired product as a white powder: $^{1}H-NMR$ (δ , DMSO)3.37 (dd, J = 3.4, 7.9 Hz, 1H), 3.60 (dd, J = 7.9 Hz, 1H),3.77 (dd, J = 3.4 Hz, 1H), 4.47 (s, 2H), 7.15-7.25 (m,2H), 7.35-7.48 (m, 2H), 7.65 (br s, 1H); MS (ion spray) 214 (M+1); Anal. Calc'd for $C_{10}H_{12}FNO_3$: C, 56.33; H, 5.67; N, 6.57. Found: C, 56.32; H, 5.66; N, 6.39.

Preparation 354

A solution of 1,3-dicyclohexylcarbodiimide (6.18 g, 30 mmol) in 30 mL of tetrahydrofuran was added to a stirred solution of 6.1 g (30 mmol) of Boc-a-methylalanine and 4.14 g (36 mmol) of N-hydroxysuccinimide in 60 mL of tetrahydrofuran at ambient temperature under nitrogen and the mixture was stirred at ambient temperature under 20 nitrogen for overnight. The mixture was cooled to ~0 ℃ and filtered through celite. The precipitate was washed with cold (0-5 °C) tetrahydrofuran. The filtrate was purified concentrated to dryness and by flash chromatography (5% acetone/dichloromethane) to give 7.25 25 g (80.6%) of the desired product as a white powder: 1H-NMR $(\delta, DMSO)$ 1.39 (s, 9H), 1.48 (s, 6H), 2.78 (s, 4H), 7.57 (br s, ~0.5H); MS (ion spray) 301 (M+1); Anal. Calc'd for $C_{13}H_{20}FN_2O_6$: C, 51.99; H, 6.71; N, 9.33. Found: C, 52.70;

H, 6.67; N, 9.30. 30

10

418

Preparation 355

Forty mL of tetrahydrofuran was added to a stirred suspension of 4.26 g (20 mmol) of the product of Preparation 353 in 100 mL 1N sodium bicarbonate and the mixture was stirred for 15 min to give a fine suspension. Forty mL of water was added to the reaction mixture to dissolve the suspension to form a clear solution. To this stirred solution at ambient temperature under nitrogen atmosphere was added a solution of 5.71 g (19 mmol) of 10 354 in the product of Preparation 60 mL of tetrahydrofuran over 3 h and the resulting mixture was stirred at ambient temperature under nitrogen for 4 h. The resulting turbid solution was diluted with 150 mL of 15 water to form a clear solution which was extracted with petroleum ether (2 x 200 mL). The aqueous layer was cooled to 0-5 °C, acidified to pH ~3.0 with cold (0-5 °C) 1N HCl (~100 mL) followed by 0.5N HCl (~15 mL)+, and the separated white precipitate was quickly extracted with 20 ethyl acetate $(1 \times 200 \text{ mL})$. The aqueous layer was saturated with sodium chloride, further acidified to pH ~1, and quickly extracted with ethyl acetate (1 x 200 mL). The ethyl acetate extracts were combined, washed with brine , dried over anhydrous sodium sulfate, filtered and concentrated to dryness to give 7.03 g 25 (92.9%) of the desired product as a white foam: $^{1}H-NMR$ (δ , DMSO) 1.29 (s, 6H), 1.33 (s, 9H), 3.66 (dd, J = 4.0 Hz, 1H), 3.75 (dd, J = 4.3 Hz, 1H), 4.38-4.48 (m, 1H), 4.44(s, 2H), 7.00-7.20 (m, 3H), 7.28-7.42 (m, 3H), 12.8 (br)30 s, ~0.5H); MS (ion spray) 399 (M+1); Anal. Calc'd for $C_{19}H_{27}FN_2O_6$: C, 57.28; H, 6.83; N, 7.03. Found: C, 57.05; H, 6.84; N, 6.87.

419

Preparation 356

7.0 g of 5% palladium on carbon (dry) was added under 5 nitrogen atmosphere to a solution of 6.7 g (22 mmol) the product of Preparation 3 from Examples Part 1 in 35 mL of tetrahydrofuran and the resulting slurry was hydrogenated at ~50 psi of hydrogen for 2 h. The mixture was filtered through celite and the catalyst was washed with tetrahydrofuran (15 x 25 mL). The resulting yellow 10 brown filtrate (~300 mL) was added to a mixture of 6.6 g (16.57 mmol) of the product of Preparation 355 and 2.97 g (22 mmol) of 1-hydroxybenzotriazole. To this stirred brown solution was added 3.7 g (18 mmol) of 1,3dicyclohexylcarbodiimide as solid in one lot and the 15 mixture was stirred at ambient temperature for 20 h, then it was concentrated. The residue was dissolved in 150 mL of ethyl acetate and filtered through celite. The brown filtrate was washed with 1N sodium bicarbonate , brine , dried over anhydrous sodium sulfate, filtered, 20 concentrated to dryness to give 12.3 g of crude product as a tan foam which was purified by flash chromatography (silica gel, 3% methanol in dichloromethane) to give 7.75 g (71.3%) of the desired product as a tan foam: $^{1}\text{H-NMR}$ (δ , DMSO) 1.17 (t, J = 7.2 Hz, 3H), 1.25-1.35 (m, 15H), 3.52-25 3.65 (m, 1H), 3.66-3.72 (m, 1H), 3.76 (s, 3H), 4.21 (q, J)= 7.2, 2H), 4.42 (d, J = 3.4 Hz, 2H), 4.53-4.62 (m, 1H),6.35 (d, J = 3.4 Hz, 1H), 6.98 (d, J = 8.7 Hz, 2H), 7.05-7.30 (m, 5H), 7.32 (d, J = 8.7 Hz), 7.51 (s, 1H), 10.20 30 (br s, 1H); MS (ion spray) 656 (M+1); Anal. Calc'd for

420

 $C_{33}H_{42}FN_5O_8$: C, 60.45; H, 6.46; N, 10.68. Found: C, 61.35; H, 6.57; N, 10.98.

Preparation 357

5

10

15

20

14 mL (14 mmol) of 1M solution of lithium hydroxide was added in one lot to a stirred solution of the product of Preparation 356 (7.2 g, 11 mmol) in 51 mL of dioxane and 30 mL of deionized water. The mixture was stirred at ambient temperature for 20 min and extracted it with ether (3 x 40 mL). The light yellow colored ether extracts were discarded. The brown aqueous layer (pH ~12) was cooled in ice-water bath and acidified to pH ~4 with cold 0.5N HCl and quickly extracted the separated light brown precipitate with ethyl acetate (1 \times 150 mL). The light yellow colored aqueous layer was saturated with sodium chloride, further acidified to pH ~3 and extracted with ethyl acetate . The aqueous layer was discarded. The organic extracts were combined, washed with brine , dried over anhydrous sodium sulfate, filtered, and concentrated to dryness to give 6.9 g (100%)of the desired product as a tan foam: ${}^{1}\text{H-NMR}$ (δ , DMSO) 1.20-1.45 (m. 15H), 3.50-3.60 (m, 1H), 3.65-3.72 (m, 1H), 3.76 (s, 3H), 4.425 (d, J = 3)Hz, 2H), 4.50-4.65 (m, 1H), 6.205 (d, 1.9 Hz, 1H), 6.99 (d, J = 8.7 Hz, 2H), 7.05-7.32 (m, 5H), 7.33 (d, J = 8.7)Hz, 2H), 7.51 (t, J = 1.9 Hz, 1H), 10.20 (br s, 1H), 13.48 (br s, 1H); MS (ion spray) 628 (M+1), 629 (M+2); Anal. Calc'd for $C_{31}H_{38}FN_5O_8$ 0.5 H_2O : C, 58.31; H, 6.33; N, 10.99. Found: C, 58.75; H, 6.26; N, 10.56.

25

421

Preparation 358

To a solution of 7.0 g (11 mmol) of the product of Preparation 357 in 130 mL of N,N-dimethylformamide was added 1.42 mL(12 mmol) of 4-methylpiperidine, 1.62 g (12 mmol) of 1-hydroxybenzotriazole hydrate, and 2.37 g (11.5 mmol) of dicyclohexylcarbodiimide. The reaction mixture was stirred at ambient temperature for 38 h, filtered, the precipitate was washed with ethyl acetate 10 and the filtrate was concentrated on $(2 \times 20 \text{ mL})$ rotavap. The resulting brown syrup was partitioned between ethyl acetate (200 mL) and 0.5N HCl (100 mL) and the layers were quickly separated. The organic extract washed with 100 mL each of saturated sodium 15 bicarbonate and brine. The organic extract was dried over anhydrous sodium sulfate, filtered, and concentrated to give 7 g of crude product as a tan foam which was purified by flash chromatography (silica methanol/ dichloromethane) to give 5.7 g (73%) of the desired product as light brown foam: ${}^{1}H-NMR$ (δ , DMSO) 20 0.12-0.30 (m, 0.5 H), 0.75 (d, J = 6.4 Hz, 1.5 H), 0.88(d, J = 6.0 Hz, 1.5 H), 0.80-1.15 (m, 2H), 1.20-1.40 (m, 2H)15 H), 1.45-1.67 (m, 2.5 H), 2.55-2.75 (m, 1.5 H), 2.93-3.07 (m, 0.5 H), 3.50-3.60 (m, 1H), 3.60-3.70 (m, 1H), 3.75 (d, J = 3.4 Hz, 3H), 3.60-3.85 (m, 1H), 4.30-4.45(m, 1H), 4.425 (d, J = 3.0 Hz, 2H), 4.50-4.65 (m, 1H), 6.62 (d, J = 12 Hz, 1H), 6.98 (t, J = 9.4 Hz, 2H), 7.05-7.45 (m, 10H), 10.13 (br s, 1H); MS (ion spray) 709 (M+1); Anal. Calc'd for $C_{37}H_{49}FN_6O_7$: C, 62.70; H, 6.97; N, 30 11.86. Found: C, 62.44; H, 6.99; N, 11.89.

422

Examples 203 and 204

Trifluoroacetic acid (6 mL) was added to a solution of the product of Preparation 358 (3.88 g, 5.47 mmol) in 15 mL of dichloromethane and the resulting brown solution was stirred at ambient temperature for 4 h. The mixture was poured into 300 mL of saturated sodium bicarbonate and extracted with ethyl acetate (1 x 150 mL). The yellow brown color organic layer was separated and the aqueous layer was saturated with sodium chloride and extracted 10 with fresh ethyl acetate . The aqueous layer was discarded. The organic extracts were combined, washed with brine , dried over anhydrous sodium sulfate, filtered, and concentrated to give 3.2 g (96%) diastereomeric mixture of the desired product as a light 15 brown foam: $^{1}H-NMR$ (δ . DMSO) consistent with structure; MS (ion spray) 609 (M+1); Anal. Calc'd for $C_{32}H_{41}FN_6O_5$ 0.5 H_2O : C, 62.22; H, 6.85; N, 13.60. Found: C, 62.14; H, 6.62; N, 13.62. This material (3.0 g) was purified by HPLC (8 x 15 cm Prochrom column packed with Kromsil CHI-DMP chiral phase, eluted with a mixture of 3A alcohol and dimethylethylamine in heptane) to provide the individual diastereomers which were converted to their respective hydrochloride salts as described below:

Example 203. (Isomer 1) 1.28 g (2.1 mmol) of pure free amine of isomer 1 (lot # CJ4-LFK-53B) was dissolved in 21 mL of 1N HCl, diluted with 21 mL of deionized water, and the resulting solution was freeze-dried to give 1.3 g (96%) of the desired product as a light brown powder: 1 H-30 NMR (δ , DMSO) 0.1-0.25 (m, 0.5 H), 0.75 (d, J = 6.4 Hz, 1.5 H), 0.88 (d, J = 6.0 Hz, 1.5 H), 0.8-1.22 (m, 1.5 H),

423

1.22-1.40 (m, 0.5 H), 1.40-1.75 (m, 8.5 H), 2.55-2.75 (m, 1.5 H), 2.95-3.10 (m, 0.5 H), 3.60-3.75 (m, 3H), 3.76 (d, J = 3.4 Hz, 3H), 4.25-4.45 (m, 1H), 4.50 (s, 2H), 4.65-4.75 (m, 1H), 6.79 (d, J = 12.8 Hz, 1H), 6.95-7.45 (m, 9H), 7.80-7.95 (m, 1H), 8.19 (br s, 3H, exchangeable with deuterium), 8.52 (d, J = 6.8 Hz, 1H, exchangeable with 1H, exchangeable deuterium), 10.86 (br s, deuterium); $t_R = 7.40 \text{ min}$; MS (ion spray) 610 (M+1); Anal. Calc'd for $C_{32}H_{41}FN_6O_5$ 2 HCl 0.5 H_2O : C, 55.73; H, 6.43; N, 12.19. Found: C, 55.50; H, 6.33; N, 12.12. 10 Example 204. (Isomer 2) 1.1 g (1.8 mmol) of pure free amine of isomer 2 (lot # CJ4-LFK-53A) was dissolved in 20 mL of 1N HCl, diluted with 20 mL of deionized water, and the resulting solution was freeze-dried to give 1.1 g (95%) of the desired product as a light brown powder: 1H-15 NMR (δ , DMSO) 0.05-0.25 (m, 0.5 H), 0.75 (d, J = 6.4 Hz, 1.5 H), 0.89 (d, J = 6.0 Hz, 1.5 H), 0.8-1.40 (m, 2H), 1.40-1.70 (m, 8.5 H), 2.55-2.75 (m, 1.5 H), 2.95-3.10 (m, 0.5 H), 3.60-3.75 (m, 3H), 3.77 (d, J = 3.0 Hz, 3H), 4.30-4.45 (m, 1H), 4.49 (s, 2H), 4.68-4.78 (m, 1H), 6.8720 (d, J = 13.6 Hz, 1H), 6.98-7.18 (m, 4H), 7.25-7.45 (m,

(d, J = 13.6 Hz, 1H), 6.98-7.18 (m, 4H), 7.25-7.45 (m, 5H), 8.05-8.20 (m, 1H), 8.27 (br s, 3H, exchangeable with deuterium), 8.57 (d, J = 6.8 Hz, 1H, exchangeable with deuterium), 11.04 (br s, 1H, exchangeable with deuterium);

 $t_R = 9.43$ min; MS (ion spray) 610 (M+1); Anal. Calc'd for $C_{32}H_{41}FN_6O_5$ 2 HCl 0.5 H_2O : C, 55.73; H, 6.43; N, 12.19. Found: C, 55.33; H, 6.40; N, 12.02.

Preparation 359

30

This is Preparation 457 from Examples Part 2C.

424

Preparation 360

5.5 g of 5% palladium on carbon was added under nitrogen atmosphere to a solution of 4.49 g (18 mmol) nitropyrazole in 30 mL of tetrahydrofuran and the resulting slurry was hydrogenated at ~50 psi of hydrogen for 2 h. The mixture was filtered through celite and the catalyst was washed with tetrahydrofuran (10 x 25 mL). The resulting yellow brown filtrate (~300 mL) was added to a mixture of 6.05 g (16 mmol) of the product of 10 2.43 (18 mmol) of 1-1i and a Preparation hydroxybenzotriazole. To this stirred brown solution was added 3.5 g (17 mmol) of 1,3-dicyclohexylcarbodiimide and the mixture was stirred at ambient temperature for 21 h, then concentrated. The residue was dissolved in 150 mL of 15 ethyl acetate and filtered through celite. The brown filtrate was washed successively with 100 mL each of 0.5N HCl, 1N sodium bicarbonate, brine, dried over anhydrous sodium sulfate, filtered, and concentrated to dryness to give 11.6 g of crude product as a tan foam which was 20 purified by flash chromatography (silica gel, 3% methanol in dichloromethane) to give 9.8 g (96%) of the desired product as a tan foam: $^{1}H-NMR$ (δ , DMSO) 1.15 (t, J = 7.1, 6.8 Hz, 3H), 1.20-1.40 (m, 15H), 1.45-1.90 (m, 4H), 2.45-2.65 (m, 2H), 3.77 (s, 3H), 4.10-4.28 (m, 3H), 6.30 (s, 25 1H), 6.98 (d, J = 9.0 Hz, 2H), 7.10-7.30 (m, 6H), 7.36(d, J = 8.7 Hz, 2H), 7.54 (d, J = 5.3 Hz, 1H), 7.69 (d, J)= 0.75 Hz, 1H), 7.82 (be s, 1H), 9.70 (br s, 1H); MS (ion)spray) 636 (M+1); Anal. Calc'd for $C_{34}H_{45}N_{5}O_{7}$: C, 64.23; H, 7.13; N, 11.02. Found: C, 64.50; H, 7.25; N, 11.06. 30

425

Preparation 361

16.5 mL (16.5 mmol) of 1M solution of lithium hydroxide was added in one lot to a stirred solution of the product of Preparation 360 (8.3 g, 13 mmol) in 60 mL of dioxane and 35 mL of deionized water. The mixture was stirred at ambient temperature for 20 min and extracted it with ether (3 x 40 mL). The light yellow colored ether extracts were discarded. The brown aqueous layer (pH ~12) was cooled in ice-water bath and acidified to pH ~4 with 10 cold 0.5N HCl and quickly extracted the separated light brown precipitate with ethyl acetate (1 x 200 mL). The light yellow colored aqueous layer was saturated with sodium chloride, further acidified to pH ~1-2 and extracted with ethyl acetate . The aqueous layer was 15 discarded. The organic extracts were combined, washed with brine , dried over anhydrous sodium sulfate, filtered, and concentrated to dryness to give 7.9 g (100%) the desired product as a tan foam: $^{1}\text{H-NMR}$ (δ , $DMSO/D_2O)$ 1.25-1.35 (m, 15H), 1.40-1.85 (m, 4H), 2.45-2.60 (m, 2H), 3.75 (s, 3H), 4.12-4.27 (m, 1H), 6.14 (s, 1H),6.97 (d, J = 8.7 Hz, 2H), 7.10-7.18 (m, 3H), 7.20-7.30(m. 2H), 7.35 (d. J = 8.7 Hz, 2H), 7.53 (d. J = 6.4 Hz)1H), 7.67 (d, J = 3.0 Hz, 1H), 7.80 (br s, 1H, exchangeable with deuterium), 9.70 25 (br 1H, exchangeable with deuterium), 13.30 (br 1H, S, exchangeable with deuterium); MS (ion spray) 608 (M+1); Anal. Calc'd for $C_{32}H_{41}N_5O_7$: C, 63.25; H, 6.80; N, 11.52. Found: C, 63.18; H, 6.98; N, 10,64.

Preparation 362

To a solution of 6.08 g (10 mmol) of the product of Preparation 361 in 100 mL of N, N-dimethylformamide was added 1.25 mL(15 mmol) of pyrrolidine, 1.62 g (12 mmol) of 1-hydroxybenzotriazole hydrate, and 2.27 g (11 mmol) of dicyclohexylcarbodiimide. The reaction mixture was stirred at ambient temperature for 46 concentrated. The resulting brown syrup was disolved in ethyl acetate (150 mL) and filtered. The filtrate was 10 washed successively with $0.5N\ HCl\ (2\ x\ 25\ mL)$, saturated sodium bicarbonate (2 x 25 mL) and brine (1 x 30 mL). The organic layer was dried over anhydrous sodium sulfate, filtered, and concentrated to dryness to give 5.96 g of crude product as a off white foam which was purified by 15 flash chromatography (silica gel, 3% methanol/ dichloromethane) to give 5.7 g (80.6%) of the desired product as white foam: $^{1}\text{H-NMR}$ ($\delta\text{,}$ DMSO) 1.20-1.40 (m, 15H), 1.45-1.95 (m, 8H), 2.45-2.60 (m, 2H), 2.98-3.10 (m, 1H), 3.25-3.43 (m, 2H), 3.45-3.60 (m, 1H), 3.76 (s, 3H), 20 4.15-4.30 (m, 1H), 6.36 (s, 1H), 6.98 (d, J = 8.7 Hz, 2H), 7.10-7.30 (m, 6H, 1H exchangeable with deuterium), 7.34 (dd, J = 8.7, 1.9 Hz, 2H), 7.50 (d, J = 10.6 Hz,1H), 7.63 (d, J = 6.4 Hz, 1H), 7.70-7.90 (m, exchangeable with deuterium), 9.66 (m, 1H, exchangeable 25 with deuterium); MS (ion spray) 662 (M+1); Anal. Calc'd for $C_{36}H_{48}N_6O_6$: C, 65.43; H, 7.32; N, 12.72. Found: C, 65.14; H, 7.09; N, 12.91.

WO 99/08697

Example 205

427

5

10

20

25

Trifluoroacetic acid (7 mL) was added to a solution of the product of Preparation 363 (4.05 g, 6.13 mmol) in 18 mL of dichloromethane and the resulting clear brown solution was stirred at ambient temperature for 4 h. The mixture was poured into 300 mL of saturated sodium bicarbonate and extracted with ethyl acetate (1 \times 150 mL). The yellow brown color organic layer was separated and the aqueous layer was saturated with sodium chloride and extracted with fresh ethyl acetate . The aqueous layer was discarded. The organic extracts were combined, washed with brine , dried over anhydrous sodium sulfate, filtered, and concentrated to give 3.48 g (99%) of diastereomeric mixture of the desired product (free base 15 form) as a white foam: $^{1}H-NMR$ (δ , DMSO) 1.15-1.25 (m, 6H), 1.45-1.95 (m, 8H), 2.45-2.60 (m, 2H), 2.97-3.07 (m, 1H), 3.25-3.45 (m, 2H), 3.47-3.60 (m, 1H), 3.77 (s, 3H), 4.27-4.37 (m, 1H), 6.36 (s, 1H), 6.98 (d, J = 8.7 Hz, 2H), 7.10-7.18 (m, 3H), 7.20-7.30 (m, 2H), 7.35 (d, J =8.7 Hz, 2H), 7.42 (s, 1H), 7.65 (s, 1H), 8.10 (br s, 1H, exchangeable with deuterium), 10.10 (s, 1H, exchangeable with deutrium); MS (ion spray) 562 (M+1); Anal. Calc'd for $C_{31}H_{40}N_6O_4$ 0.5 H_2O : C, 65.35; H, 7.25; N, 14.75. Found: C, 65.39; H, 7.27; N, 14.12.

Example 206

428

82 mg (0.146 mmol) of the product of Example 205 was dissolved in 2 mL of 1N HCl and diluted with 2 mL of deionized water and the resulting slightly turbid solution was freeze-dried to give 89.5 mg (97%) of the desired product as a off-white crystalline solid: 1 H-NMR (δ , DMSO) consistent with the structure; MS (ion spray) 562 (Free base) (M+1); Anal. Calc'd for $C_{31}H_{40}N_{6}O_{4}$ $H_{2}O$: C, 57.14; H, 6.50; N, 12.90. Found: C, 57.35; H, 6.77; N, 12.89.

10


Preparation 363

To a solution of phthalide (6.0 g, 44.8 mmol) in ethanol (150 mL) was added pyrrolidine (15.0 mL, 180 mmol) and the mixture stirred for 48 h at ambient temperature. The mixture was then concentrated and the residue purified by flash chromatography (silica gel, ethyl acetate) to yield 8.50 g (92 %) of the desired product as an oil.

ESMS: $(M+1)^+$ 206.2. ¹H NMR (DMSO- d_6) δ 7.53-7.45 (m, 1H), 7.42-7.33 (m, 1H), 7.31-7.25 (m, 1H), 7.23-7.17 (m, 1H), 5.11 (t, J = 5.65 Hz, 1H), 4.44 (d, J = 5.65 Hz, 2H), 3.45 (t, J = 6.78 Hz, 2H), 3.09 (t,J = 6.78 Hz, 2H), 1.92 (m, 4H). Anal. Calcd. for $C_{12}H_{15}NO_2$.0.1EtOAc: C, 69.57; H, 7.44; N, 6.54. Found: C, 69.67; H, 7.41; N, 6.54.

25

Preparation 364



10

The compound from Preparation 363 (2.05 g, 10.0 mmol) was combined with 4-nitroimidazole (1.13 g, 10.0 mmol) and triphenylphosphine (2.62 g, 10.0 mmol) in tetrahydrofuran (75 mL) and the mixture cooled in an ice bath. Diethylazodicarboxylate (2.00 mL, 13.0 mmol) was then added via syringe and the mixture stirred overnight while warming to ambient temperature. The mixture was concentrated and the residue purified by flash chromatography (silica gel, ethyl acetate) to provide 1.16 g of the desired product as a yellow solid: ESMS: (M+1)⁺ 301.2; ¹H NMR was consistent with product.

Preparation 365

The product of Preparation 364 (0.50 g, 1.7 mmol) was added to a mixture of 10% palladium/carbon (0.20 g) and palladium/

black (0.05 g) in tetrahydrofuran (30 mL) and the mixture shaken under hydrogen (40 psi) in a Parr apparatus.

- After the reduction was complete, the reaction mixture was filtered through celite and the filtrate immediately combined with 1,3-dicyclohexylcarbodiimide (0.344 g, 1.7 mmol), 1-hydroxybenzotriazole mono-hydrate (0.225 g, 1.7 mmol), and the product of Preparation 1d, (0.633 g, 1.7
- 25 mmol). After stirring overnight at ambient temperature, the mixture was concentrated and the resulting residue slurried in ethyl acetate and filtered. The filtrate was concentrated and the residue purified by flash chromatography (silica gel, chloroform/methanol) which provided 0.68 g (64 %) of product as a white solid.

430

ESMS: $(M+1)^+$ 633.2, 634.2. ¹H NMR was consistent with product. Anal. Calcd. for $C_{34}H_{44}N_6O_6$: C, 64.54; H, 7.01; N, 13.28. Found: C, 64.78; H, 7.21; N, 13.31.

5 Example 207

10

15

20

25

To a solution of the compound from Preparation 365 (0.58 g, 0.92 mmol) stirring in dichloromethane (20 mL) at room temperature was added trifluoroacetic acid (5 mL). After stirring for 1.5 h, the reaction mixture was concentrated and the residue treated with excess aqueous sodium bicarbonate. The mixture was extracted with ethyl acetate and chloroform. The extracts were concentrated and the residue purified by flash chromatography (silica gel, chloroform/methanol) to give 0.40 g of the product. ESMS: (M+1) 533.3, 534.0. ¹H NMR was consistent with product.

Anal. Calcd. for $C_{29}H_{36}N_{6}O_{4}$. 0.23CHCl₃: C, 62.68; H, 6.52; N, 15.00. Found: C, 62.84; H, 6.56; N, 14.62.

Preparation 366

To a solution of phthalide (2.10 g, 15.7 mmol) in ethanol (50 mL) was added dimethyl amine (40 % aqueous, 5.0 mL, 40 mmol) and the mixture stirred for 72 hours at ambient temperature. The mixture was then concentrated and the residue purified by flash chromatography (silica gel,

431

ethyl acetate) to yield 1.46 g (52 %) of the desired product as an oil. ESMS: $(M+1)^+$ 180.1. ¹H NMR was consistent with product.

Preparation 367

5

م م

To the compound from Preparation 366 (1.40 g, 9.0 mmol) was dissolved in dichloromethane (25 mL) followed by the addition of triethylamine (1.25 mL, 9.0 mmol). Then methane sulfonylchloride (0.70 mL, 9.0 mmol) was added 10 dropwise via syringe and the resulting mixture stirred for 0.5 h at ambient temperature. The mixture was then concentrated and the residue combined with potassium carbonate (1.24 g, 18.0 mmol) and 4-nitroimidazole (1.02 g, 9.0 mmol) in dimethylformamide (25 mL) and this 15 mixture stirred overnight at ambient temperature. The mixture was concentrated and combined with water followed by extraction with ethyl acetate. Concentration of the combined extracts left crude product which was purified by flash chromatography (silica gel, chloroform/methanol) 20 to give 0.40 g (15%) of the desired product: ESMS: $(M+1)^{+}$ 275.3. ¹H NMR (300 MHz, DMSO- d_6) δ 8.33 (d, J = 1.51 Hz, 1H), 7.85(d, J = 1.51 Hz, 1H), 7.50-7.37 (m, 3H), 7.35-7.28(m, 1H), 5.24 (s, 2H), 2.97 (s, 3H), 2.65 (s, 3H). 25 Anal. Calcd. for $C_{13}H_{14}N_4O_3$: C, 56.92; H, 5.14; N, 20.43. Found: C, 57.17; H, 5.16; N, 20.40.

Preparation 368

432

The product of Preparation 367 (1.70 g, 6.2 mmol) was added to a mixture of 10% palladium/carbon (1.2 g) and palladium/

5 black (0.5 g) in tetrahydrofuran (100 mL) and the mixture shaken under hydrogen (38 psi) in a Parr apparatus. After the reduction was complete, the reaction mixture was filtered through celite and the filtrate immediately combined with 1,3-dicyclohexylcarbodiimide (1.28 g, 6.2 mmol), 1-hydroxybenzotriazole mono-hydrate (0.840 g, 6.2 10 mmol), and the product of Preparation 1j, (2.35 g, 6.2 mmol). After stirring overnight at ambient temperature, the mixture was concentrated and the resulting residue slurried in ethyl acetate and filtered. The filtrate was concentrated and the residue purified by flash 15 chromatography (silica gel, chloroform/methanol) which provided 3.35 g (88 %) of product as a white foam. ESMS: $(M+1)^+$ 605.4, 606.5. ¹H NMR was consistent with product.

20

Example 208

To a solution of the compound from Preparation 368 (3.30 g, 5.5 mmol) stirring in dichloromethane (40 mL) at room temperature was added trifluoroacetic acid (10 mL).

WO 99/08697

10

433

PCT/US98/17201

After stirring for 2.5 h, the reaction mixture was concentrated and the residue treated with excess aqueous sodium bicarbonate. The mixture was extracted with ethyl acetate and the combined extracts were dried over sodium sulfate and concentrated. The residue was purified by flash chromatography (silica gel, chloroform/methanol) to give 1.10 g of the product as an off white solid: ESMS: (M+1)⁺ 505.2, 506.4. ¹H NMR was consistent with product. Anal. Calcd. for C₂₈H₃₆N₆O₃·0.1CHCl₃: C, 65.34;H, 7.04; N, 16.27. Found: C, 65.67; H, 7.08; N, 15.94.

Preparation 369



Methyl-3-bromomethyl benzoate (6.08 g, 26.5 mmol) was

combined with 4-nitroimidazole (3.00 g, 26.5 mmol) and
potassium carbonate (9.04g, 26.5 mmol) in
dimethylformamide (75 mL) and the mixture stirred
overnight at ambient temperature. The mixture was then
concentrated and the residue taken up in water and
extracted with chloroform.

The combined extracts were dried over sodium sulfate and concentrated leaving a residue which was purified by flash chromatography (silica gel, chloroform/methanol) to give

- 25 5.77 g (83 %) of product as an off white solid. ESMS: $(\text{M+1})^+ \ \ 262.2; \ ^1\text{H NMR} \ (300 \ \text{MHz}, \ \text{DMSO-}d_6) \ \delta \ \ 8.54$ (s, 1H), 8.05 (s, 1H), 8.02 (d, 1H), 7.95 (d, J = 7.7 Hz, 1H), 7.69 (d, J = 7.7 Hz, 1H), 7.57 (t, J = 7.7 Hz, 1H), 5.41 (s, 2H) 3.87 (s, 3H). Anal. Calcd. for:
- 30 $C_{12}H_{11}N_3O_4^{-0}.15CHCl_3$: C, 65.54; H, 7.33; N, 13.90. Found: C, 65.41; H, 7.37; N, 13.48.

Preparati n 370



To a solution of Preparation 369 (4.00 g, 15.3 mmol) in ethanol (5 mL) and tetrahydrofuran (10 mL) was added sodium hydroxide (50 mL of 2 N aqueous solution) and the mixture stirred at ambient temperature until hydrolysis was complete. The aqueous mixture was acidified to pH 2.0 with aqueous hydrochloric acid and the organics extracted with ethyl acetate. Concentration left the 10 crude acid which was combined with 1,3-dicyclohexylcarbodiimide (3.15 g, 15.3 mmol), 1-hydroxybenzotriazole mono-hydrate (2.06 g, 15.3 mmol), and dimethylamine (40 % aqueous, 1.75 mL, 14.0 mmol) in tetrahydrofuran (75 mL) and the mixture 15 stirred overnight at ambient temperature. The mixture was concentrated and the resulting residue slurried in ethyl acetate and filtered. The filtrate was concentrated and the residue purified by flash chromatography (silica gel, chloroform/methanol) which provided 2.50 g (60 %) of the 20 desired product as a tan solid. ESMS: (M+1) 275.3; ¹H NMR (300 MHz, DMSO- d_6) δ 8.54 (d, J = 1.25 Hz, 1H), 8.3 (d, J = 1.25 Hz, 1H), 7.50-7.32 (m, 4H), 5.34 (s, 2H), 2.97 (s, 3H), 2.87 (s, 3H). Anal. Calcd. for: 25 $C_{12}H_{11}N_3O_4$: C, 56.93; H, 5.14; N, 20.43. Found: C, 57.21; H, 5.34; N, 20.36.

Preparation 371

435

The product of Preparation 370 (1.0 g, 4.0 mmol) was added to a mixture of 10% palladium/carbon (0.80 g) and palladium/black (0.30 g) in tetrahydrofuran (100 mL) and the mixture shaken under hydrogen (39 psi) in a Parr apparatus.

After the reduction was complete, the reaction mixture was filtered through celite and the filtrate immediately combined with 1,3-dicyclohexylcarbodiimide (0.824 g, 4.0 mmol), 1-hydroxybenzotriazole mono-hydrate (0.540 g, 4.0 mmol), and the product of Preparation 1j, (1.51 g, 4.0 mmol). After stirring overnight at ambient temperature, the mixture was concentrated and the resulting residue slurried in ethyl acetate and filtered. The filtrate was concentrated and the residue purified by flash chromatography (silica gel, chloroform/methanol) which provided 2.05 g (84 %) of the desired product as a

white solid. ESMS: $(M+1)^+$ 605.4, 606.5. ¹H NMR was consistent with product. Anal. Calcd. for $C_{33}H_{44}N_6O_5$: C, 65.54;

H, 7.33; N, 13.90. Found: C, 65.41; H, 7.37; N, 12.73.

Example 209

25

5

10

15

To a solution of the compound from Preparation 371 (2.00 g, 3.3 mmol) stirring in dichloromethane (30 mL) at room

WO 99/08697

436

PCT/US98/17201

temperature was added trifluoroacetic acid (10 mL). After stirring for 2 h, the reaction mixture was concentrated and the residue treated with excess aqueous sodium bicarbonate. The mixture was extracted with ethyl acetate and the combined extracts were dried over sodium sulfate and concentrated. The residue was purified by flash chromatography (silica gel, chloroform/methanol) to give pure free base. This material was dissolved in minimal ethyl acetate/methanol and treated with diethyl ether saturated with hydrochloric acid. Concentration left a semi-solid which was taken up in chloroform, concentrated, and dried to give 0.765 g of a light yellow solid. ESMS: $(M+1)^+$ 505.2, 506.4. ¹H NMR was consistent with product.

Anal. Calcd. for: $C_{28}H_{41}N_6O_3$ ·1CHCl₃: C, 49.98; H, 5.64; 15 N, 12.06. Found: C, 49.51; H, 6.14; N, 11.67.

Triethylphosphonoacetate (13.44 g, 60 mmol) was 20 dissolved in tetrahydrofuran (100 mL) and the mixture cooled, under a nitrogen atmosphere, to -50°C in a dry ice/acetone bath.

Then n-butyllithium (37.5 mL, 1.6 M in hexanes, 60 mmol) was added via syringe and the mixture stirred at -40 to 25 -50°C for 20 min. before the addition of o-tolualdehyde (6.0 g, 50 mmol). After stirring for 3 h and warming to 10°C, the mixture was quenched with water and the product extracted with ethyl acetate. The combined extracts were dried over sodium sulfate and concentrated yielding an oil which was purified by flash chromatography (silica gel, ethyl acetate) to net 9.25 g (97 %) of the desired product. FDMS: (M+) 190.1. 1 H NMR (DMSO- d_{6}) δ 7.88 (d,

PCT/US98/17201 WO 99/08697

437

J = 15.83 Hz, 1H), 7.71 (d, J = 7.91 Hz, 1H), 7.35-7.15(m, 3H),

 $6.50 \, (d, J = 15.83 \, Hz, 1H), 4.19 \, (q, J = 7.16 \, Hz, 2H),$ 2.39 (s, 3H), 1.26 (t, J = 7.16 Hz, 3H).

5 Anal. Calcd. for $C_{12}H_{14}O_2$: 0.1EtOAc: C, 74.82; H, 7.49. Found: C, 74.65; H, 7.75.

Preparation 373



- The product from Preparation 372 (3.00 g, 15.8 mmol) was 10 dissolved in carbontetrachloride (50 mL) and Nbromosuccinimide (2.82 g, 15.8 mmol) and 2,2'azobisisobutyronitrile (cat. 50 mg) were added. This mixture was heated at reflux for 7 h after which the
- mixture was cooled and filtered. The filtrate was 15 concentrated and added together with potassium carbonate(4.40 g, 31.9 mmol) and 4-nitroimidazole (1.78 g, 15.8 mmol) in dimethylformamide (75 mL). After stirring for 72 h, themixture was concentrated, water
- added, and extracted with ethyl acetate. The combined 20 extracts were concentrated and the resulting residue purified by flash chromatography(chloroform/methanol) to give 2.60 g (55%) of the product as an oil. ESMS: (M+1)* 302.3.
- ¹H NMR (300 MHz, DMSO- d_6) δ 8.32 (d, J = 1.5 Hz, 1H), 7.95 25 (d, J = 15.6 Hz, 1H), 7.86 (d, J = 1.5 Hz, 1H), 7.84-7.70(m, 1H), 7.50-7.38 (m, 2H), 7.19-7.12 (m, 1H), 6.54 (d, 1H)J = 15.6 Hz, 1H, 5.59 (s, 2H), 4.20 (q, J = 7.0 Hz, 2H),1.27 (t, J = 7.0 Hz, 3H). Anal. Calcd. for $C_{15}H_{15}N_3O_4$:
- 30 C, 55.31; H, 4.64; N, 12.69. Found: C, 55.37; H, 4.75;

438

N, 12.80.

Preparation 374



To a solution of Preparation 373 (2.50 g, 8.3 mmol) in ethanol (5 mL) and tetrahydrofuran (10 mL) was added sodium hydroxide (30 mL of 2 N aqueous solution) and the mixture stirred at ambient temperature until hydrolysis was complete. The aqueous mixture was acidified to pH 1.8

with aqueous hydrochloric acid and extracted with ethyl acetate. Concentration left the 2.20 g of the acid as a tan solid.ESMS: $(M+1)^+$ 264.2. ¹H NMR(300 MHz, DMSO- d_6) δ 12.58 (bs, 1H), 8.32 (s, 1H), 7.90 (d, J = 15.6 Hz, 1H), 7.87 (s, 1H), 7.80-7.70 (m, 1H), 7.50-7.37 (m, 2H), δ 17-7.10 (m, 1H), δ 44 (d, J = 15.6 Hz, 1H), 5.58 (s

15 7.17-7.10 (m, 1H), 6.44 (d, J = 15.6 Hz, 1H), 5.58 (s, 2H).

The acid (2.00 g, 7.3 mmol) was combined with 1,3-dicyclo-hexylcarbodiimide (1.51 g, 7.3 mmol), 1-hydroxybenzotriazole mono-hydrate (0.98 g, 7.3 mmol), and

dimethylamine (40 % aqueous, 1.00 mL, 8.0 mmol) in tetrahydrofuran (100 mL) and the mixture stirred overnight at ambient temperature. The mixture was concentrated and the resulting residue purified by flash chromatography (silica gel, chloroform/methanol) which

25 provided 1.78 g (81 %) of the desired amide.
 ESMS: (M+1)* 301.2, 302.3. Anal. Calcd. for C₁₅H₁₆N₄O₃:
 C, 59.99; H, 5.37; N, 18.66. Found: C, 59.79; H, 5.34;
 N, 18.51.

PCT/US98/17201 WO 99/08697


439

The product of Preparation 374 (1.20 g, 4.0 mmol) was added to a mixture of 10% palladium/carbon (0.95 g) and palladium/black (0.25 g) in tetrahydrofuran (100 mL) and

mixture shaken under hydrogen (38 psi) in a Parr apparatus.

After the reduction was complete, the reaction mixture was filtered through celite and the filtrate immediately combined with 1,3-dicyclohexylcarbodiimide (0.824 g, 4.0 mmol), 1-hydroxybenzotriazole mono-hydrate (0.540 g, 4.0 mmol), and the product of Preparation 1j, (1.51 g, 4.0 mmol). After stirring overnight at ambient temperature, the mixture was concentrated and the resulting residue slurried in ethyl acetate and filtered. The filtrate was concentrated and the residue purified by flash chromatography (silica gel, chloroform/methanol) which provided 1.75 g (69 %) of product as a yellow solid. ESMS: $(M+1)^+$ 633.5, 634.5. ¹H NMR was consistent with product. Anal. Calcd. for C35H48N6O5'0.2CHCl3: C, 64.38; H, 20 7.40; N, 12.80. Found: C, 64.41; H, 7.63; N, 12.14.

Example 210



5

10


15

25 To a solution of the compound from Preparation 375 (1.6 g, 2.5 mmol) stirring in dichloromethane (20 mL) at room

440

temperature was added trifluoroacetic acid (5 mL). After stirring for 1 h, the reaction mixture was concentrated and the residue treated with excess aqueous sodium The mixture was extracted with ethyl bicarbonate. acetate and the combined extracts were dried over sodium sulfate and concentrated. The residue was purified by flash chromatography (silica gel, dichloromethane/methanol) to give the free base. compound was dissolved in minimal ethyl acetate and excess ether saturated with hydrochloric acid was added. 10 Concentration and drying netted 0.53 g of the hydrochloride salt as a white solid. ESMS: (M+1) 533.2, 534.3. H NMR was consistent with product. Anal. Calcd. for $C_{30}H_{39}N_6O_3$ 3HCl: C, 59.99; H, 5.37; N, 18.66. Found: C, 59.79; H, 5.34; N, 18.51. 15

Preparation 376



dichloromethane(55 mL) was added triethylamine (17 mL, 122 mmol) and 4-dimethylaminopyridine (cat. 50 mg) and the mixture cooled to 0°C under a nitrogen atmosphere. Then methanesulfonyl chloride (9.4 mL, 122 mmol)was added dropwise via syringe and the resulting mixture stirred for 2 h while warming to ambient temperature. The mixture was concentrated in vacuo and the residue taken up in dimethylformamide (100 mL)and solid potassium carbonate (20 g, 146 mmol) added along with 4-nitro-imidazole (2.75 g, 24.3 mmol). This mixture stirred at ambient

temperature

To 3-aminobenzyl alcohol (3.00 g, 24.4 mmol) in

10

WO 99/08697 PCT/US98/17201

441

for 90 h after which the mixture was concentrated and the residue taken up in water and the product extracted with ethyl acetate. The combined extracts were washed with water and dried over sodium sulfate. Concentration left 5 a semi-solid which was tritruated in chloroform and filtered to yield the desired bis-sulfonamide (3.5 g, 38%) as a white solid. ESMS: $(M+1)^+$ 375.2. ¹H NMR (300) MHz, DMSO- d_6) δ 8.51 (d, J = 1.13 Hz, 1H), 8.02 (d, J = 1.13 Hz, 1H), 7.65-7.45 (m, 4H), 5.36 (s, 2H) 3.54 (s, 6H).



To a solution of Preparation 376 (0.75 g, 2.0 mmol) in ethanol (3 mL) and tetrahydrofuran (3 mL) was added sodium 15 hydroxide (10 mL of 2 N aqueous solution) and the mixture stirred at ambient temperature until hydrolysis was complete. The aqueous mixture was acidified to pH 3.5 with aqueous hydrochloric acid and the organics extracted with ethyl acetate. Concentration left a residue which 20 was

purified by flash chromatography (silica gel, chloroform/ methanol) which afforded 0.41 g (69 %) of pure product as a yellow solid. ESMS: $(M+1)^{+}$ 297.4; ¹H NMR (300 MHz,

DMSO- d_6) δ 9.81 (s, 1H), 8.47 (d, J = 1.51 Hz, 1H), 7.98 25 (d, J = 1.51 Hz, 1H), 7.40-7.30(m, 1H), 7.20-7.00(m,3H), 5.31 (s, 2H), 3.00 (s, 3H). Anal. Calcd. for $C_{11}H_{12}N_4O_4S$:

C, 44.59; H, 4.08; N, 18.91. Found: C, 44.12; H, 4.28; 30 N, 18.90.

442

Preparation 378

The product of Preparation 377 (0.55 g, 1.86 mmol) was added to a mixture of 10% palladium/carbon (0.40 g) and palladium/black (0.15 g) in tetrahydrofuran (50 mL) and the mixture shaken under hydrogen (40 psi) in a Parr apparatus. After the reduction was complete, the reaction mixture was filtered through celite and the filtrate immediately combined with 1,3-

10 dicyclohexylcarbodiimide (0.384 g,

1.86 mmol), 1-hydroxybenzotriazole mono-hydrate (0.251 g,

1.86 mmol), the product of Preparation 1j, (0.70 g,

1.85 mmol), and additional tetrahydrofuran (30 mL).

After stirring overnight at ambient temperature, the

15 mixture was concentrated and the resulting residue slurried in ethyl acetate and filtered. The filtrate was concentrated and

the residue purified by flash chromatography (silica gel, chloroform/methanol) which provided 0.66 g (57 %) of the

desired product as a white solid. ESMS: (M+1)* 627.4, 628.5. ¹H NMR was consistent with product.

Example 211

To a solution of the compound from Preparation 379(0.60 g, 0.96 mmol) stirring in dichloromethane (25 mL) at room temperature was added trifluoroacetic acid (5 mL). After stirring for 2 h, the reaction mixture was concentrated

PCT/US98/17201 WO 99/08697

443

and the residue treated with excess aqueous sodium bicarbonate. The mixture was extracted with ethyl acetate and the combined extracts were dried over sodium sulfate and concentrated. The residue was purified by

flash 5

> chromatography (silica gel, chloroform/methanol) to give 0.25 g of the desired pure product as a white solid. ESMS: $(M+1)^+$ 527.2, 528.3. ¹H NMR was consistent with product.

Anal. Calcd. for $C_{26}H_{34}N_6O_4S$: C, 59.30; H, 6.51; N, 15.96. 10 Found: C, 59.13; H, 6.65; N, 15.66.

Preparation 380

- The product of Preparation 377 (0.60 g, 2.03 mmol) was 15 added to a mixture of 10% palladium/carbon (0.50 g) and palladium/black (0.10 g) in tetrahydrofuran (40 mL) and the mixture shaken under hydrogen (39 psi) in a Parr apparatus. After the reduction was complete, the
- reaction mixture was filtered through celite and the 20 filtrate immediately combined with 1,3dicyclohexylcarbodiimide (0.418 g,
 - 2.03 mmol), 1-hydroxybenzotriazole mono-hydrate(0.274 g,
 - 2.03 mmol), the product of Preparation 1d, (0.77 g,
- 2.03 mmol), and additional tetrahydrofuran (40 mL). 25 After stirring overnight at ambient temperature, the mixture was concentrated and the resulting residue slurried in ethyl acetate and filtered. The filtrate was concentrated and the residue purified by flash
- chromatography (silica gel, chloroform/methanol) which 30

444

provided 0.28 g (22 %) of the desired product as a light tan solid. ^{1}H NMR was consistent with product.

Example 212

5

To a solution of the compound from Preparation 380 (0.28 g, 0.44 mmol) stirring in dichloromethane (10 mL) at room temperature was added trifluoroacetic acid (2.5 mL).

After stirring for 2.5 h, the reaction mixture was

concentrated and the residue treated with excess aqueous sodium bicarbonate. The mixture was extracted with ethyl acetate and the combined extracts were dried over sodium sulfate and concentrated. The residue was purified by flash

chromatography (silica gel, chloroform/methanol) to give 45 mg of the desired product as an off white solid. ESMS: (M+1)⁺ 529.2. ¹H NMR was consistent with product. Anal. Calcd. for C₂₅H₃₂N₆O₅S.2H₂O: C, 53.18; H, 6.43; N, 14.88. Found: C, 53.40; H, 6.30; N, 13.53.

20

Preparation 381

To a solution of 3-isochromanone (3.00 g, 20.0 mmol) in ethanol (40 mL) was added dimethyl amine (40 % aqueous, 3.0 mL, 24 mmol) and the mixture stirred for 72 hours at ambient temperature. The mixture was then concentrated and the residue purified by flash chromatography (silica gel, chloroform/methanol) to yield 3.16 g (81 %) of the desired product as an oil. ESMS: (M+1)* 194.3. ¹H NMR

445

 $(DMSO-d_6)$ δ 7.40-7.34 (m, 1H), 7.25-7.13 (m, 2H), 7.09-7.03 (m, 2H), 5.04 (t, J = 5.3 Hz, 1H), 4.44 (d, J = 5.3 Hz, 1H)Hz, 2H), 3.72 (s, 2H), 3.02 (s, 3H), 2.85 (s, 3H).

Preparation 382

To the compound from Preparation 381 (2.00 g, 10.4 mmol) was dissolved in dichloromethane (40 mL) followed by the addition of triethylamine (1.74 mL, 12.5 mmol). Then methane sulfonylchloride (0.96 mL, 12.5 mmol) was added dropwise via syringe and the resulting mixture stirred for 2.5 h at ambient temperature. The mixture was then concentrated and the residue combined with potassium carbonate (2.85 g, 20.6 mmol) and 4-nitroimidazole (1.17 g, 10.4 mmol) in dimethylformamide (50 mL) and this mixture stirred overnight at ambient temperature. The mixture was concentrated and combined with water followed by extraction with ethyl acetate. Concentration of the combined extracts left crude product which was purified by flash chromatography (silica gel, chloroform/methanol) to give 0.88 g of the desired product as a light solid. ESMS: $(M+1)^+$ 289.1. ¹H NMR (300 MHz, DMSO- d_6) δ 8.32 (s, 1H), 7.88 (s, 1H), 7.30-7.05 (m, 4H), 5.27 (s, 2H), 3.85 (s, 2H), 3.06 (s, 3H), 2.82 (s, 3H). Anal. Calcd. for $C_{14}H_{16}N_4O_3$:0.05CHCl₃: C, 57.35; H, 5.50; N, 19.04. 25 Found:

C, 57.10; H, 5.53; N, 18.80.

5

10

15

20

446

Preparation 383

The product of Preparation 382 (0.75 g, 2.6 mmol) was added to a mixture of 10% palladium/carbon (0.60 g) and palladium/black (0.15 g) in tetrahydrofuran (40 mL) and the mixture shaken under hydrogen (37 psi) in a Parr apparatus.

After the reduction was complete, the reaction mixture was filtered through celite and the filtrate immediately combined with 1,3-dicyclohexylcarbodiimide (0.537 g, 2.6 10 mmol), 1-hydroxybenzotriazole mono-hydrate (0.351 g, 2.6 mmol), and the product of Preparation 1j, (0.983 g, 2.6 mmol). After stirring overnight at ambient temperature, the mixture was concentrated and the resulting residue slurried in ethyl acetate and filtered. The filtrate was 15 concentrated and the residue purified by flash chromatography (silica gel, chloroform/methanol) which provided 0.87 g (84 %) of product as a tan foam. ESMS: $(M+1)^+$ 619.7, 620.8. ¹H NMR was consistent with 20 product.

Example 213

To a solution of the compound from Preparation 383 (0.80 g, 1.3 mmol) stirring in dichloromethane (15 mL) at room temperature was added trifluoroacetic acid (3 mL). After stirring for 1 h, the reaction mixture was concentrated

PCT/US98/17201 WO 99/08697


447

and the residue treated with excess aqueous sodium bicarbonate. The mixture was extracted with ethyl acetate and the combined extracts were dried over sodium sulfate and concentrated. The residue was purified by flash chromatography (silica gel, dichloromethane/methanol) to give 0.14 g of the desired product as a white solid. ESMS: $(M+1)^+$ 519.3, 520.4. ¹H NMR was consistent with product. Anal. Calcd. for C₂₉H₃₈N₆O₃·0.2CHCl₃: C, 65.48;

H, 7.23; N, 15.69. Found: C, 65.06; H, 7.29; N, 15.62.

10

25



To a solution of 5-flouro-2-methylbenzoic acid (9.00 g, 58.4 mmol) in methanol (200 mL) was added tosic acid 15 mono-hydrate (cat. 0.50 g) and the mixture heated overnight at reflux. The mixture was then concentrated and the residue purified by flash chromatography (silica gel, chloroform/ methanol) to yield 5.80 g of the desired ester as an oil. 20

ESMS: $(M+1)^+$ 167.9; ¹H NMR (300 MHz, DMSO- d_6) δ 7.56 (dd, J = 9.4 and 2.6 Hz, 1H), 7.40-7.33 (m, 2H), 3.83 (s, 3H), 2.48 (s, 3H).

Preparation 385



The product from Preparation 384 (5.30 g, 31.5 mmol) was dissolved in carbontetrachloride (75 mL) and Nbromosuccin-

448

imide (5.60 g, 31.5 mmol) and 2,2'-azabisisobuytronitrile (cat. 70 mg) were added. This mixture was heated at reflux for 4 h after which the mixture was cooled and filtered. The filtrate was concentrated and added together with potassium carbonate (5.00 g, 36.2 mmol) and 4-nitroimidazole (3.56 g, 31.5 mmol)in dimethylformamide (50 mL). After stirring for overnight at ambient temperature, the mixture was concentrated, water added, and extracted with ethyl acetate. The combined extracts were concentrated and the resulting residue purified by 10 flash chromatography (chloroform/methanol) to give 6.25 g of product as a white solid. ESMS: (M+1) 280.1. H NMR (300 MHz, DMSO d_6) δ 8.36 (d, J = 1.51 Hz, 1H), 7.90 (d, J = 1.51 Hz, 1H), 7.73 (dd, J = 9.42 and 2.64 Hz, 1H), 7.57-7.45 (m, 15 1H), 7.35-7.25 (m, 1H), 5.60 (s, 2H), 3.87 (s, 3H). Anal. Calcd. for $C_{12}H_{10}N_3O_4F$: C, 51.62; H, 3.61; N, 15.05. Found: C, 52.19; H, 3.75; N, 14.87.

20 Preparation 386

25

To a solution of Preparation 385 (3.25 g, 11.6 mmol) in ethanol (5 mL) and tetrahydrofuran (5 mL) was added sodium hydroxide (25 mL of 2 N aqueous solution) and the mixture stirred at ambient temperature until hydrolysis was complete. The aqueous mixture was acidified to pH 2.5

with aqueous hydrochloric acid and the organics extracted with ethyl acetate. Concentration left the crude acid as a solid. The acid (2.30 g, 8.7 mmol) was combined with 1,3-dicyclo-hexylcarbodiimide (1.79 g, 8.7 mmol),

PCT/US98/17201 WO 99/08697

449

1-hydroxybenzotriazole mono-hydrate (1.17 g, 8.7 mmol), and dimethylamine (40 % aqueous, 2.00 mL, 40.0 mmol) in tetrahydrofuran (60 mL) and the mixture stirred overnight at ambient temperature. The mixture was concentrated and the resulting residue slurried in ethyl acetate and filtered. The filtrate was concentrated and the residue purified by flash chromatography (silica gel, chloroform/methanol) which provided for recovery of the desired amide, 1.85 g.


10 ESMS: $(M+1)^+$ 293.1. ¹H NMR (300 MHz, DMSO- d_6) δ 8.32 (d, J = 1.51 Hz, 1H), 7.84 (d, J = 1.51 Hz, 1H), 7.55-7.45 (m,1H), 7.35-7.20 (m, 2H), 5.21 (s, 2H), 2.96 (s, 3H), 2.65 (s, 3H).

15

20

25

Preparation 386A



The product of Preparation 386 (0.60 g, 2.0 mmol) was added to a mixture of 10% palladium/carbon (0.50 g) and palladium/black (0.10 g) in tetrahydrofuran (100 mL) and the mixture shaken under hydrogen (38 psi) in a Parr apparatus. After the reduction was complete, the reaction mixture was filtered through celite and the filtrate immediately combined with 1,3dicyclohexylcarbodiimide (0.422 g, 2.0 mmol), 1hydroxybenzotriazole mono-hydrate (0.277 g, 2.0 mmol), and the product of Preparation 1j, (0.775 g, 2.0 mmol). After stirring 72 h at ambient temperature, the mixture was concentrated and the resulting residue slurried in ethyl acetate and filtered. The filtrate was 30 concentrated and the residue purified by flash

450

chromatography (silica gel, chloroform/methanol) which provided 0.78 g (62 %) of product. ESMS: $(M+1)^+$ 623.5. ¹H NMR was consistent with product.

5

20

Example 214

To a solution of the compound from Preparation 386A (0.65 g, 1.0 mmol) stirring in dichloromethane (20 mL) at room temperature was added trifluoroacetic acid (4 mL). After stirring for 2 h, the reaction mixture was concentrated and the residue treated with excess aqueous sodium bicarbonate. The mixture was extracted with ethyl acetate and the combined extracts were dried over sodium sulfate and concentrated. The residue was purified by flash chromatography (silica gel, dichloromethane/methanol) to give 0.05 g of the desired product. ESMS: (M+1)* 523.4, 524.5. ¹H NMR was consistent with product. Anal. Calcd. for C20H35N6O3F.0.4CHCl3: C, 63.86; H, 6.70; N, 15.93.

Preparation 387

Found: C, 64.28; H, 6.96; N, 15.54.



Potassium carbonate (6.90 g, 50 mmol) was combined with 4-nitroimidazole (5.65 g, 50 mmol) and 2-cyanobenzyl bromide (9.80g, 50 mmol) in dimethylformamide (100 mL) and the mixture stirred overnight at ambient temperature.

451

The mixture was then concentrated and the residue taken up in water and extracted with ethyl acetate. The combined extracts were dried over sodium sulfate and concentrated leaving a tan solid which was purified by flash chromatography (silica gel, chloroform/methanol) to give 9.76 g (85 %) of product. ESMS: (M+1)⁺ 229.3. ¹H

(300 MHz, DMSO- d_6) δ 8.42 (d, J = 1.51 Hz, 1H), 7.95 (d, J = 1.51 Hz, 1H), 7.92 (dd, J = 7.91 Hz, 1.13 Hz, 1H), 7.77-7.69 (m,, 1H), 7.60-7.52 (m, 1H), 7.31 (d, J = 7.91 Hz, 1H), 5.57 (s, 2H). Anal. Calcd. for $C_{11}H_9N_4O_2$: C, 57.89; H, 3.53; N, 24.55. Found: C, 57.65; H, 3.53; N, 24.33.

15

20

25

30

oil.

Preparation 388



The compound of Preparation 387 (2.50 g, 11 mmol) was dissolved in tetrahydrofuran (40 mL) and borane added (1.0 M in tetrahydrofuran, 15 mL, 15 mmol) and the mixture stirred overnight at ambient temperature. The reaction was then quenched by the slow addition of methanol. After hydrogen evolution had ceased, the mixture was concentrated and the residue purified by flash chromatography (silica gel, chloroform/methanol). Recovered 0.61 g (17%) of the desired amine as a light

ESMS: $(M+1)^+$ 233.1. ¹H NMR (300 MHz, DMSO- d_6) δ 8.42 (d, J = 1.13 Hz, 1H), 7.96 (d, J = 1.13 Hz, 1H), 7.43 (d, J = 7.16 Hz, 1H), 7.38-7.15 (m,, 2H), 7.09 (d, J = 7.16 Hz, 1H), 5.42 (s, 2H), 3.81 (s,2H), 3.30 (bs, 2H).

PCT/US98/17201

452

Preparation 389



The amine from Preparation 388 (0.90 g, 3.88 mmol) was dissolved in tetrahydrofuran (40 mL) and methyl

isocyanate 5

> (0.50 mL, 8.5 mmol) was added dropwise via syringe and the mixture stirred overnight at ambient temperature. The resulting precipitate was filtered and dried which netted 0.89 g, (79%) of the desired urea.ESMS: (M+1)*

290.2. ¹H NMR (300 MHz, DMSO- d_6) δ 8.35 10 (d, J = 1.51 Hz, 1H), 7.92 (d, J = 1.51 Hz, 1H), 7.37-7.20 (m, 3H), 7.12 (d, J = 7.54 Hz, 1H), 6.45-6.35 (m,1H),


5.90-5.78 (m, 1H), 5.39 (s, 2H), 4.28 (d, J = 6.03 Hz,

15

2.56 (d, J = 4.90 Hz, 3H). Anal. Calcd. for $C_{13}H_{15}N_5O_3$: C, 53.97; H, 5.23; N, 24.21. Found: C, 53.71; H, 5.14; N, 24.10.

20

Preparation 390



The product of Preparation 389 (0.50 g, 1.73 mmol) was added to a mixture of 10% palladium/carbon (0.40 g) and palladium/

black (0.10 g) in tetrahydrofuran (80 mL) and the mixture shaken under hydrogen (39 psi) in a Parr apparatus. After

453

6 h, the reaction mixture was filtered through celite to remove catalyst and undissolved starting material (0.15 g, poor solubility). The filtrate was immediately combined

with 1,3-dicyclohexylcarbodiimide (0.356 g, 1.73 mmol), 1-hydroxybenzotriazole mono-hydrate (0.233 g, 1.73 mmol), and the product of Preparation 1d, (0.657 g, 1.73 mmol). After stirring for 72 h at ambient temperature, the mixture was concentrated and the resulting residue slurried in ethyl acetate and filtered. The filtrate was concentrated and the residue purified by flash chromatography (silica gel, chloroform/methanol) which provided 0.25 g of product.

ESMS: (M+1) + 622.5. H NMR was consistent with product.

15

Example 215

To a solution of the compound from Preparation 390 (0.24 g, 0.34 mmol) stirring in dichloromethane (10 mL) at room temperature was added trifluoroacetic acid (1.5 mL). After stirring for 2 h, the reaction mixture was concentrated and the residue treated with excess aqueous sodium bicarbonate. The mixture was extracted with ethyl acetate. The extracts were concentrated and the residue purified by flash chromatography (silica gel, chloroform/methanol) to give 0.065 g of product as a white solid. ESMS: (M+1)⁺ 522.3, 523.5. ¹H NMR was consistent with product.

454



To a solution of 2-aminobenzylamine, 3.0 g (24.6 mmol) in 50 mL of methanol and 15 mL of water at 0°C was added 2.1 g of sodium bicarbonate. The resulting slurry was stirred for 10 min. then 3.9 g of the product of Preparation 306 was added. The reaction mixture was stirred overnight while slowly warming to ambient temperature and was then concentrated to dryness. The resulting residue was absorbed onto a silica pad and was chromatographed using 80% ethyl acetate/hexanes as eluant to yield 2.66 g (50%) of the desired product as an orange solid. ¹H-NMR is consistent with structure; MS (ion spray) 219.2 (M+1); Anal. Calc'd for C₁₀H₁₀N₄O₂: C, 55.04; H, 4.62; N, 25.67. Found: C, 55.31; H, 4.72; N, 25.76.


15

20

25

10

Preparation 392



To a solution of the product of Preparation 391, 2.5 g (11.5 mmol) and triethylamine, 4 mL (28.8 mmol) in 200 mL of dichloromethane at 0°C was added 2.25 mL (28.8 mmol) of methanesulfonyl chloride. The resulting slurry was stirred overnight, slowly warming to ambient temperature and was then concentrated to dryness. The residue was slurried in water and filtered, then slurried in chloroform, filtered and dried under vacuum to yield 3.1 g (72%) of the desired product as a white solid. ¹H-NMR is consistent with structure; MS (ion spray) 375.2 (M+1);

455

Anal. Calc'd for $C_{12}H_{14}N_4O_6S_2$: C, 38.50; H, 3.77; N, 14.96. Found: C, 38.45; H, 3.66; N, 14.74.

Preparation 393



5

To a slurry of the prodcut of Preparation 392, 1.7 g (4.6 mmol) in 60 mL of absolute ethanol and 60 mL of tetrahydrofuran was added 23 mL (23 mmol) of 1N sodium hydroxide. The resulting solution was stirred 90 min then was concentrated to dryness. The residue was 10 partitioned between ethyl acetate and water and acidified to pH = 3 with 1N HCl. The mixture was extracted with ethyl acetate. The combined organics were washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. The residue was chromatographed 15 on silica gel using a gradient of 3-10% methanol/chloroform as eluant to yield 0.7 g (51%) of the desired product as a yellow solid. H-NMR is consistent with structure; MS (ion spray) 297.4 (M+1); Anal. Calc'd for $C_{11}H_{12}N_4O_4S$: C, 44.59; H, 4.08; N, 18.91. Found: C, 20 44.38; H, 4.23; N, 18.65.

Preparation 394

25 A solution of the product of Preparation 393, 0.59 g (2.0 mmol) in 40 mL of tetrahydrofuran was added to a slurry of 10% palladium on carbon (1.0 g) in 40 mL of tetrahydrofuran. The mixture was hydrogenated at 40 psi

PCT/US98/17201 WO 99/08697

456

for 40 min, then filtered through celite. To this solution was added 0.76 g (2.0 mmol) of the product of Preparation 1d, 0.3 g (2.2 mmol) of 1hydroxybenzotriazole and 0.46 g (2.2 mmol) of dicylohexylcarbodiimide. The reaction mixture was stirred overnight at ambient temperature and was then concentrated to dryness. The residue was partitioned between ethyl acetate and water and was then extracted with ethyl acetate. The combined organics were washed with brine, dried over sodium sulfate, filtered and 10 concentrated to dryness. The residue was chromatographed on silica gel using a gradient of 3-10% methanol/chloroform to yield 0.45 g (36%) of the desired product as a tan foam. ¹H-NMR is consistent with structure; MS (ion spray) 629.3 (M+1). 15

To a solution of the product of Preparation 394, 0.35 g (0.56 mmol) in 12 mL of dichloromethane was added 4 mL of trifluoroacetic acid. The resulting mixture was stirred for 1 h, then was concentrated to dryness. The residue was partitioned between ethyl acetate and saturated sodium bicarbonate and was then extracted with ethyl acetate. The combined organics were washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. To a solution of the residue in chloroform was added HCl-saturated ether. The slurry was concentrated to dryness to yield 0.34 g (100%) of the desired product 30 as a tan solid. 1H-NMR is consistent with structure; MS (ion spray) 529.2 (M+1); Anal. Calc'd for


20

25

457

 $C_{25}H_{32}N_6O_5S^2.3HC1$: C, 49.03; H, 5.64; N, 13.72. Found: C, 48.94; H, 5.62; N, 13.39.

Preparation 395



5

10

15

20

To a solution of 4-nitroimidazole (12.76g, 112.8 mmol) in anhydrous dimethylformamide (350 mL) was added sodium hydride (4.96 g, 124.1 mmol, 60% in mineral oil), followed by ethyl bromoacetate dropwise over a period of five minutes, and the reaction was stirred at room temperature overnight. The solution was concentrated and the residue was dissolved in a solution of 20% isopropanol in chloroform, washed with 1N HCl, dried over magnesium sulfate, and concentrated to give 35.0 g crude product as an orange solid. These solids were treated with anhydrous diethyl ether (100 mL), sonicated, heated on a steam bath, and cooled in a 0°C refrigerator overnight. The mixture was filtered and the solids washed with ice cold diethyl ether, then dried to give 16.56 g of the title compound as a white solid, 74% vield: ^{1}H NMR ($^{6}-DMSO$, δ): 1.19 (t, J=6.1 Hz, 3H), 4.15 (q, J = 6.1 Hz, 2H), 5.02 (s, 2H), 7.80 (s, 1H), 8.32 (s, 2H), 7.80 (s, 2H), 8.32 (s, 21H); MS (ion spray) 200 ($M^+ + 1$).

25

Preparation 396

To a mixture of 4.0 g 10% palladium on carbon in 100 mL tetrahydrofuran was added the product of Preparation 395 (9.96 g, 50.0 mmol). The mixture was subjected to 60 psi hydrogen on a Parr apparatus for 2 h, then filtered

through celite to remove the catalyst. The filtrate was and the resulting oil was dissolved in concentrated anhydrous dimethylformamide (10 mL). This solution was added to a solution of the product of Preparation 1d from (20.92 g, 55.0 mmol) and Examples Part 2A anhydroushydroxybenzotriazole (8.42, 55.0 mmol) in dimethylformamide (150 mL) which had been cooled to -13°C in an ice/acetone bath. Dicyclohexyl carbodiimide (11.35 g, 55.0 mmol) was then added, and the stirred solution was allowed to warm to room temperature overnight. 10 solvent was removed, ethyl acetate was added, and the insoluble dicyclohexyl urea was removed by filtration. The filtrate was washed with 1N HCl, saturated aqueous sodium bicarbonate solution, brine, dried over magnesium and concentrated. Purification 15 sulfate, by flash gel (eluting with 5% chromotography on silica methanol/dichloromethane) gave 5.39 g of the title compound as a tan solid, 31% yield: ¹H NMR (d^6 -DMSO, δ): 1.17 (t, J = 6.1 Hz, 3H), 1.25 (m, 15H), 3.55 (q, J = 3.7Hz, 1H), 3.65 (m, 1H), 4.08 (q, J = 3.7 Hz, 2H), 4.40 (s, 20 2H), 4.56 (brs, 1H), 4.84 (s, 2H), 7.15 (s, 1H), 7.22 (m, 6H), 7.38 (s, 1H), 7.42 (m, 1H), 10.10 (brs, 1H): MS (ion spray) 532 $(M^{+} +1)$.

25

30

Preparation 396A

To a solution of the product of Preparation 396 (7.17 g, 13.49 mmol) in dioxane (100 mL) was added a solution of lithium hydroxide (623 mg, 14.84 mmol) in water (50 mL). After 1h, more lithium hydroxide (60 mg, 1.42 mmol) was added. After 30 min, the reaction was quenched with 1N

459

HCl and concentrated. The resulting oil was dissolved in a 20% isopropanol/chloroform solution, washed with 1N HCl, water, dried over magnesium sulfate, and concentrated to give 6.12 g of the title compound as a golden yellow solid, 90% yield: ¹H NMR (d⁶-DMSO, δ): 1.32 (m, 15H), 3.62 (m, 1H), 3.73 (m, 1H), 4.48 (s, 2H), 4.62 (brs, 1H), 4.85 (s, 2H), 7.19 (s, 1H), 7.30 (m, 6H), 7.51 (m, 1H), 7.59 (s, 1H), 10.24 (brs, 1H): MS (ion spray) 504 (M⁺ +1).

10

Preparation 397



To a solution of L-proline (5.38 g, 25.0 mmol) 15 anhydrous tetrahydrofuran (75 mL) was added carbonyl diimidazole (4.05 g 25.0 mmol), and the solution stirred at room temperature. After 30 minutes, benzylamine (2.73 mL, 25.0 mmol) was added dropwise, and the reaction stirred at room temperature overnight. The solvent was 20 removed and the residue dissolved dichloromethane, washed with 1N HCl , saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, concentrated to give 6.29 g of a white solid. solid dissolved in diethyl ether saturated with dry HCl(g) 25 (35 mL, ~3N in HCl) and stirred vigorously at room temperature for 2 h. The volatiles were removed and the residue dissolved in 20% isopropanol/chloroform, washed with saturated aqueous sodium bicarbonate solution, dried over magnesium sulfate, and concentrated to give 2.7 q of the title compound as a yellow oil, 64% yield: 1H NMR $(d^6-DMSO, \delta): 1.55 (m, 1H), 1.62 (m, 1H), 1.90 (m, 1H),$ 2.75 (m, 2H), 3.00 (brs, 1H), 3.23 (brs, 1H), 3.48 (m,

460

1H), 4.20 (d, J = 5.4 Hz, 2H), 7.17 (m, 3H), 7.22 (t, J = 5.0 Hz, 2H), 8.34 (s, 1H); MS(ion spray) 205 (M⁺ +1).

Preparation 398

5

10

15

20

25

To a 0°C solution of the product of Preparation 396A (1.01 g, 2.0 mmol) in anhydrous tetrahydrofuran (10 mL) was added the product of Preparation 397 (409 mg, mmol), 1-hydroxybenzotriazole (322 mg, 2.1 mmol), dicyclohexyl carbodiimide (433 mg, 2.1 mmol). The stirred reaction was allowed to warm to room temperature The insoluble dicyclohexyl urea was removed by filtration, and the filtrate was concentrated. resulting foam was dissolved in ethyl acetate, washed 1N HCl, saturated aqueous sodium bicarbonate with solution, brine, dried over magnesium sulfate, concentrated. Purification by flash chromotography on silica gel (eluting with 4-7% methanol/dichloromethane) gave 636 mg of the title compound as a faintly yellow solid, 46% yield: ¹H NMR (d^6 -DMSO, δ): 1.32 (m, 15H), 1.93 (m, 4H), 2.07 (m, 1H), 3.50 (m, 1H), 3.64 (m, 1H), 3.72 (m, 1H), 4.28 (d, J = 6.3 Hz, 2H), 4.34 (m, 2H), 4.47 (s, 2H)2H), 4.63 (brs, 1H), 4.93 (s, 2H), 7.21 (d, J = 12.5 Hz, 2H), 7.30 (m, 10H), 7.45 (s, 1H), 8.38 (m, 1H), 10.16 (brs, 1H): MS(ion spray) 690 (M $^+$ +1). Anal. (C₃₆H₄₇N₇O₇): H,N;C: calcd 62.68, found 62.13.

461

Example 217

To a solution of the product of Preparation 398 (627 mg, 0.91 mmol) was added acetic acid saturated with $HCl_{(g)}$ (20 mL, ~3N in HCl) and the solution stirred vigorously at room temperature for 2h. The solution was concentrated, toluene was added and the mixture concentrated to assist in removal of acetic acid. The resulting white solid was treated with diethyl ether, sonicated, and 700 mg yellow solid was isolated by filtration. The solid was dried overnight to give 626 mg of the title compound as a yellow powder, 100% yield: ¹H NMR (d⁶-DMSO, δ): consistent with structure; MS(ion spray) 590 (M⁺ +1 of free base); Anal. (C₃₁H₄4N₇O₇Cl): C,H; N: calcd. 14.81, found 13.83.

15

20

10

Preparation 399

To a solution of L-proline (5.38 g, 25.0 mmol) in anhydrous tetrahydrofuran (75 mL) was added carbonyl diimidazole (4.05 g 25.0 mmol), and the solution stirred at room temperature. After 30 min, aniline (2.28 mL, 25.0 mmol) was added dropwise, and the reaction stirred at room temperature overnight. The solvent was removed and the residue dissolved in dichloromethane, washed with 1N HCl, saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and concentrated to give 6.30 g of a white solid. The solid was dissolved in dichloromethane (50 mL) and treated with trifluoroacetic

462

acid (15 mL). The solution was stirred vigorously at room temperature for 1h. The volatiles were removed and the residue dissolved in dichloromethane, washed with saturated aqueous sodium bicarbonate solution, dried over magnesium sulfate, and concentrated to give 1.62 g of the title compound as a white solid, 39% yield: $^1\text{H NMR}$ (d^6-DMSO, δ): consistent with structure; MS(ion spray) 191 (M^+ +1).

10

15

20

25

Preparation 400

To a solution of the product of Preparation 396A (504 mg, 1.00 mmol) in anhydrous tetrahydrofuran (20 mL) was added the product of Preparation 399 (173mg, 0.91 mmol), PyBOP® (473mg, 0.91 mmol), and diisopropylethyl amine (0.48 mL, 2.73 mmol). The reaction was stirred at room temperature overnight. The solution was concentrated and the resulting residue was dissolved in ethyl acetate, washed with 1N HCl, saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and reconcentrated. Purification by flash chromotography on silica gel (eluting with 4-7% methanol/dichloromethane) gave 244 mg of the title compound as an orange solid, 40% yield: 1 H NMR (d 6 -DMSO, δ): consistent with structure; MS(ion spray) 676 (M * +1); Anal. (C $_{35}$ H₄₅N₇O₇): C,H,N.

463

Example 218

To the product of Preparation 400 (234 mg, 0.35 mmol) was added acetic acid saturated with $HCl_{(g)}$ (10 mL, ~3N in HCl) and the solution stirred vigorously at room temperature for 1h. The solution was concentrated, toluene was added and the mixture concentrated to assist in removal of acetic acid. The residue was then treated with diethyl ether, sonicated, and 235 mg a light tan solid was isolated by filtration. The solid was dried to give 230 mg of the title compound, 100% yield: ¹H NMR (d⁶-DMSO, δ): consistent with structure; MS(ion spray) 576 (M⁺ +1 of free base); Anal. (C₃₀H₃₉N₇O₆Cl₂): C,H; N: theory 14.71, found 14.13.

15

20

25

10

Preparation 401

To a solution of L-proline (5.38 g, 25.0 mmol) in anhydrous tetrahydrofuran (75 mL) was added carbonyl diimidazole (4.05 g 25.0 mmol), and the solution stirred at room temperature. After 30 minutes, phenethylamine (3.14 mL, 25.0 mmol) was added dropwise, and the reaction stirred at room temperature overnight. The solvent was removed and the residue dissolved in dichloromethane, washed with 1N HCl , saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and concentrated to give 6.77 g of a white solid. The solid was dissolved in anhydrous dichloromethane (50 mL) and

464

treated with trifluoroacetic acid (15 mL). The solution was stirred vigorously at room temperature for 1h. The volatiles were removed and the residue dissolved in dichloromethane, washed with saturated aqueous sodium bicarbonate solution , dried over magnesium sulfate, and concentrated to give 1.08 g of the title compound as a white solid, 23% yield: ¹H NMR (d⁶-DMSO, δ): consistent with structure; MS(ion spray) 219 (M⁺ +1).

10

Preparation 402

To a solution of the product of Preparation 396A (504 mg, 1.00 mmol) in anhydrous tetrahydrofuran (20 mL) was added the product of Preparation 401 (198 mg, 0.91 mmol),

- PyBOP® (473 mg, 0.91 mmol), and diisopropylethyl amine (0.48 mL, 2.73 mmol). The reaction was stirred at room temperature overnight. The solution was concentrated and the resulting orange residue was dissolved in ethyl acetate, washed with 1N HCl, saturated aqueous sodium
- bicarbonate solution, brine, dried over magnesium sulfate, and concentrated. Purification by flash chromotography on silica gel (eluting with 4-7% methanol/dichloromethane) gave 272 mg of the title compound as an light orange solid, 42% yield: ¹H NMR (d⁶-
- 25 DMSO, δ): consistent with structure; MS(ion spray) 704 (M*+1); Anal. (C₃₇H₄₉N₇O₇): C,H,N.

WO 99/08697

10

PCT/US98/17201

465

Example 219

To the product of Preparation 402 (262 mg, 0.37 mmol) was added acetic acid saturated with $HCl_{(g)}$ (15 mL, ~3N in HCl) and the solution stirred vigorously at room temperature for 1h. The solution was concentrated, toluene was added and the mixture concentrated. The residue was then treated with diethyl ether to give 243 mg of the desired compound, 94% yield: ¹H NMR (d⁶-DMSO, δ): consistent with structure; MS(ion spray) 603 (M⁺ +1 of free base); Anal. (C₃₂H₄₅N₇O₆Cl₂): C,H,N.

Preparation 403

To a solution of the product of Preparation 396A (504 mg, 1.00 mmol) in tetrahydrofuran (20 mL) was added S-alphamethylbenzylamine (0.14 mL, 0.91 mmol), PyBOP® (473mg, 0.91 mmol), and diisopropylethyl amine (0.48 mL, 2.73 mmol). The reaction was stirred at room temperature overnight. The solution was concentrated and the resulting orange residue was dissolved in dichloromethane, washed with 1N HCl, saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and concentrated. Purification by flash chromotography on silica gel (eluting with 5%

466

methanol/dichloromethane) gave 232 mg of the title compound as a white solid, 42% yield: ¹H NMR (d⁶-DMSO, δ): consistent with structure; MS(ion spray) 607 (M⁺ +1); Anal. (C₃₂H₄₂₉N₆O₆): H,N; C: theory 63.35, found 62.64.

5

Example 220

To the product of Preparation 403 (222 mg, 0.37 mmol) was added acetic acid saturated with HCl_(g) (15 mL, ~3N in HCl) and the solution stirred vigorously at room temperature for 1h. The solution was concentrated, toluene was added and the mixture concentrated. The residue was then treated with diethyl ether and sonicated to give 218 mg of the title compound as a tan solid, 100% yield: ¹H NMR (d⁶-DMSO, δ): consistent with structure; MS(ion spray) 607 (M⁺ +1 of free base); Anal. (C₂₇H₃₆N₆O₄Cl₂): C,H,N.

Preparation 404

20

25

To a solution of the product of Preparation 396A (504 mg, 1.00 mmol) in tetrahydrofuran (20 mL) was added S-alphamethylbenzylamine (0.14 mL, 0.91 mmol), PyBOP® (benzotriazol-1-yl-oxytripyrrolidinephosphonium hexafluorophosphate) (473 mg, 0.91 mmol), and

WO 99/08697

10

467

PCT/US98/17201

diisopropylethyl amine (0.48 mL,2.73 mmol). The reaction was stirred at room temperature overnight. The solution was concentrated and the resulting orange residue was dissolved in dichloromethane, washed with 1N HCl, saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and reconcentrated. Purification by flash chromotography on silica gel (eluting with 15% methanol/1:1 diethyl ether:hexane) gave 168 mg of the title compound as a white solid, 30% yield: 1 H NMR (d 6 -DMSO, δ): consistent with structure; MS(ion spray) 607 (M $^{+}$ +1).

Example 221

- To the product of Preparation 404 (164 mg, 0.237 mmol) was added acetic acid saturated with $HCl_{(g)}$ (15 mL, ~3N in HCl) and the mixture stirred vigorously at room temperature for 1h. The solution was concentrated , toluene was added and the mixture concentrated to assist in removal of acetic acid . The residue was then treated with diethyl ether, sonicated, and 161mg of the title compound was isolated by filtration as a yellow solid, 100% yield: ¹H NMR (d⁶-DMSO, δ): consistent with structure; MS(ion spray) 507 (M⁺ +1 of free base); Anal.
- 25 $(C_{27}H_{36}N_6O_4Cl_2)$: C,H; N: theory 14.06, found 13.15.

PCT/US98/17201 WO 99/08697

468

Preparation 405

To a solution of the product of Preparation 396A (504 mg, 1.00 mmol) in tetrahydrofuran (20 mL) was added 1,2,3,4tetrahydroisoguinoline (0.13 mL, 0.91 mmol), PyBOP® (473 mg, 0.91 mmol), and diisopropylethyl amine (0.48 mL, 2.73 mmol). The reaction was stirred at room temperature overnight. The solution was concentrated and the resulting orange residue was dissolved in dichloromethane, washed with 1N HCl , saturated aqueous 10 sodium bicarbonate solution, brine, dried over magnesium sulfate, and reconcentrated. Purification by flash chromotography on silica gel (eluting with 5% methanol/dichloromethane) gave 560 mg of the title compound as a yellow solid, 99% yield: ^{1}H NMR (d $^{6}\text{-DMSO}$, δ): 15 consistent with structure; MS (ion spray) 619 (M+1).

Example 222

To the product of Preparation 405 (583 mg, 0.94 mmol) was 2.0 added acetic acid saturated with $HCl_{(g)}$ (15 mL, ~3N in HCl) and the mixture stirred vigorously at room temperature for 1h. The solution was concentrated, toluene was added and the mixture concentrated to assist 25 in removal of acetic acid . The residue was then treated

469

with diethyl ether, sonicated, and 573 mg of the title compound was isolated by filtration as a yellow solid, 100% yield: 1H NMR (d^6 -DMSO, δ): consistent with structure; Ion spray MS (M^+ +1 of free base): 519; Anal. ($C_{27}H_{37}N_6O_4Cl_3$): C,N; H: calcd 5.94, found 6.68.

Preparation 406

To a solution of the product of Preparation 396A (504 mg, 1.00 mmol) in anhydrous tetrahydrofuran (20 mL) was added 10 indoline (0.10 mL, 0.91 mmol), PyBOP® (473 mg, 0.91 mmol), and diisopropylethyl amine (0.48 mL, 2.73 mmol). The reaction was stirred at room temperature overnight. The solution was concentrated and the resulting orange 15 residue was dissolved in dichloromethane, washed with 1N HCl , saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and reconcentrated. Purification by flash chromotography on silica gel (eluting with 4-7% methanol/dichloromethane) gave 543mg 20 of the title compound as a yellow solid, 99% yield: 1H NMR (d^6-DMSO, δ) : consistent with structure; MS(ion spray) 605 $(M^{+} + 1)$.

Example 223

5

470

To the product of Preparation 406 (534 mg, 0.88 mmol) was added acetic acid saturated with HCl_(g) (15 mL, ~3N in HCl) and the mixture stirred vigorously at room temperature for 1h. The solution was concentrated, toluene was added and the mixture concentrated to assist in removal of acetic acid. The residue was then treated with diethyl ether, sonicated, and 336 mg of the title compound was isolated by filtration as a yellow solid, 62% yield: ¹H NMR (d⁶-DMSO, δ): consistent with structure; MS(ion spray) 505 (M⁺ +1 of free base); Anal. (C₂₇H₃₅N₆O₄Cl₃): C,H,N.

Preparation 407

90

15 To a solution of D-proline (1.08 g, 5.0 mmol) in anhydrous tetrahydrofuran (20 mL) was added carbonyl diimidazole (0.81 g, 5.0 mmol), and the solution stirred at room temperature. After 30 min, benzylamine (0.55 mL, 5.0 mmol) was added dropwise, and the reaction stirred at 20 room temperature overnight. The solvent was removed and the residue dissolved in dichloromethane, washed with 1N HCl , saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and concentrated to give 1.31 g of a white solid. This solid was dissolved in dichloromethane (20 mL) and treated with trifluoroacetic acid (5 mL). The reaction was stirred vigorously at room temperature for 1h. The mixture was concentrated and the residue dissolved in 20% isopropanol/chloroform, washed with saturated aqueous 30 sodium bicarbonate solution, dried over magnesium sulfate, and concentrated to give 600 mg of the title compound as a yellow oil, 59% yield: ¹H NMR (d^6 -DMSO, δ): consistent with structure; MS(ion spray) 205 (M+1).

471

Preparation 408

To a solution of the product of Preparation 396A (504 mg, 1.00 mmol) in tetrahydrofuran (20 mL) was added the product of Preparation 407 (186 mg, 0.91 mmol), PyBOP® (473mg, 0.91 mmol), and disopropylethyl amine (0.48 mL, 2.73 mmol). The reaction was stirred at room temperature overnight. The solution was concentrated and the resulting orange residue was dissolved in 20% 10 isopropanol/chloroform, washed with 1N HCl , saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and reconcentrated. Purification by flash chromotography on silica gel (eluting with 5-10% 15 methanol/dichloromethane) gave 454 mg of the title compound as a white solid, 72% yield: ¹H NMR (d^6 -DMSO, δ): consistent with structure; MS(ion spray) 690 (M+1); Anal. $(C_{36}H_{47}N_7O_7)$: H,N;C: calcd 62.68, found 61.72.

20

25

Example 224

To the product of Preparation 408 (441 mg, 0.64 mmol) was added acetic acid saturated with $HCl_{(g)}$ (20 mL, ~3N in HCl) and the solution stirred vigorously at room temperature for 2h. The solution was concentrated,

472

toluene was added and the mixture concentrated. The residue was then treated with diethyl ether, sonicated, and 459 mg of white solid was isolated by filtration. The solid was dried to give 431 mg of the title compound as a white powder, 96% yield: ¹H NMR (d⁶-DMSO, δ): consistent with structure; MS(ion spray) 590 (M⁺ +1); Anal. (C₃₁H₄₂N₇O₅Cl₃): C,H,N.

Preparation 409

10

To a solution of D-proline (1.08 g, 5.0 mmol) in anhydrous tetrahydrofuran (20 mL) was added carbonyl diimidazole (0.81 g, 5.0 mmol), and the solution stirred at room temperature. After 30 minutes, aniline (0.46 mL, 5.0 mmol) was added dropwise, and the reaction stirred at 15 room temperature overnight. The solvent was removed and the residue dissolved in dichloromethane, washed with 1N HCl , saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and concentrated to give 1.20 g of a white solid. This solid was dissolved 20 in anhydrous dichloromethane (10 mL) and treated with trifluoroacetic acid (10 mL). The reaction was stirred vigorously at room temperature for 1h. The volatiles were removed and the residue dissolved in 20% isopropanol/chloroform, washed with saturated aqueous 25 sodium bicarbonate solution , dried over magnesium sulfate, and concentrated to give 281 mg of the title compound as a white solid, 46% yield: ¹H NMR (d^6 -DMSO, δ): 1.60 (p, J = 8.6 Hz, 2H), 1.73 (m, 1H), 1.98 (m, 1H), 30 2.82 (t, J = 7.6 Hz, 2H), 3.30 (brs, 1H), 3.63 (dd, J =7.6, 4.3 Hz, 1H), 6.99 (t, J = 8.2 Hz, 1H), 7.22 (t, J =

473

8.2 Hz, 2H), 7.59 (d, J = 8.2 Hz, 1H); MS(ion spray) 191 (M⁺ +1).

Preparation 410

5

To a solution of the product of Preparation 396A (619 mg, 1.23 mmol) in dimethylformamide (25 mL) was added the product of Preparation 409 (281 mg, 1.48 mmol), PyBOP® (640 mg, 1.23 mmol), and diisopropylethyl amine (0.44 mL, 10 2.50 mmol). The reaction was stirred at room temperature overnight. The solution was concentrated and the resulting residue was dissolved in dichloromethane, washed with 0.1N HCl, saturated aqueous sodium bicarbonate solution, brine, dried over magnesium 15 sulfate, and reconcentrated. Purification by flash chromotography on silica gel (eluting with 3-6% methanol/dichloromethane) gave 265 mg of the title compound as a yellow solid, 32% yield: ¹H NMR (d^6 -DMSO, δ): consistent with structure. MS(ion spray) 676 (M+1).

20

25

Example 225

To the product of Preparation 410 (262 mg, 0.39 mmol) was added acetic acid saturated with $HCl_{(g)}$ (15 mL, ~3N in HCl) and the mixture stirred vigorously at room

474

temperature for 1h. The solution was concentrated, toluene was added and the mixture. The residue was then treated with diethyl ether, sonicated, to provide 267 mg of the desired product by a tan solid which was dried to give 225 mg of the title compound, 87% yield: ¹H NMR (d⁶-DMSO, δ): consistent with structure; MS(ion spray) 576 (M⁺ +1 of free base); Anal. (C₃₀H₄₁N₇O₆Cl₂): C,H,N.

Preparation 411

10

15

20

30

To a solution of L-proline (1.08 g, 5.0 mmol) in anhydrous tetrahydrofuran (20 mL) was added carbonyl diimidazole (0.81 g, 5.0 mmol), and the solution stirred at room temperature. After 30 minutes, phenethylamine (0.63 mL, 5.0 mmol) was added dropwise, and the reaction stirred at room temperature overnight. The solvent was removed and the residue dissolved in dichloromethane, washed with 1N HCl , saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and concentrated to give 1.39 g of a white solid. This solid was dissolved in anhydrous dichloromethane (10 mL) and treated with trifluoroacetic acid (10 mL). reaction was stirred vigorously at room temperature for The volatiles were removed and the residue dissolved in 20% isopropanol/chloroform, washed with saturated aqueous sodium bicarbonate solution, dried over magnesium sulfate, and concentrated to give 386 mg of the title compound as a yellow oil, 35% yield: ¹H NMR (d⁶-DMSO, δ): consistent with structure; MS(ion spray) 219 (M⁺ +1).

Preparation 412

To a solution of the product of Preparation 396A (360 mg, 0.71 mmol) in dimethylformamide (25 mL) was added the product of Preparation 411 (156 mg, 0.71 mmol), PyBOP® (372 mg, 0.71 mmol), and diisopropylethyl amine (0.25 mL, 1.42 mmol). The reaction was stirred at room temperature overnight. The solution was concentrated and the resulting orange residue was dissolved in 10 dichloromethane, washed with 0.1N HCl, saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and reconcentrated. Purification by flash chromotography on silica gel (eluting with 4-8% methanol/dichloromethane) gave 304 mg of the title compound as a light tan solid, 60% yield: 1H NMR (d6-DMSO, 15 δ): consistent with structure; MS(ion spray) 704 (M⁺ +1).

Example 226

To the product of Preparation 412 (262mg, 0.37 mmol) was added acetic acid saturated with HCl_(g) (15 mL, ~3N in HCl) and the mixture stirred vigorously at room temperature for 1h. The solution was concentrated, toluene was added and the mixture concentrated. The residue was then treated with diethyl ether, sonicated,

and 281 mg of the title compound was isolated by filtration as a yellow solid, 92% yield: 1H NMR (d^6 -DMSO, δ): consistent with structure; MS(ion spray) 604 (M^* +1 of free base); Anal. ($C_{32}H_{44}N_7O_5Cl_3$): C,H; N: calcd 13.75, found 12.96.

5

Preparation 413

To a solution of L-proline (5.38 g, 25.0 mmol) in 10 anhydrous tetrahydrofuran (75 mL) was added carbonyl diimidazole (4.05 g, 25.0 mmol), and the solution stirred at room temperature. After 30 min, p-methoxybenzylamine (3.26 mL, 25.0 mmol) was added dropwise, and the reaction stirred at room temperature overnight. The solvent was 15 removed and the residue dissolved in dichloromethane, washed with 1N HCl , saturated sodium bicarbonate solution, dried over magnesium sulfate, and concentrated. The resulting solids were dissolved in anhydrous dichloromethane (50 mL) and treated with trifluoroacetic acid (20 mL). The reaction was stirred vigorously at 20 room temperature for 1h. The volatiles were removed and the residue dissolved in dichloromethane, washed with saturated aqueous sodium bicarbonate solution , dried over magnesium sulfate, and concentrated to give 0.95 g 25 of the title compound as a yellow oil, 16% yield: 1H NMR (d^6-DMSO, δ) : consistent with structure; MS(ion spray) 235 $(M^+ +1)$.

PCT/US98/17201 WO 99/08697

477

Preparation 414

To a solution of the product of Preparation 396A (1.01 g, 2.0 mmol) in anhydrous tetrahydrofuran (10 mL) was added the product of Preparation 413 (469 mg, 2.0 mmol), 1hydroxybenzotriazole (322 mg, 2.1 mmol), and dicyclohexyl carbodiimide (433 mg, 2.1 mmol). The reaction was stirred at room temperature overnight then filtered and concentrated. The resulting orange residue was dissolved in dichloromethane, washed with 1N HCl, saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and reconcentrated. Purification by flash chromotography on silica gel (eluting with 10% methanol/dichloromethane) gave 370 mg of the title compound as a light yellow foam, 26% yield: 1H NMR (d6-15 DMSO, δ): consistent with structure; MS(ion spray) 720 (M⁺ +1).; Anal. $(C_{37}H_{49}N_7O_8)$: H,N; C: calcd 61.74, found 60.96.

Example 227

20

25

5

10


To the product of Preparation 414 (360 mg, 0.50 mmol) was added acetic acid saturated with $HCl_{(g)}$ (15 mL, ~3N in HCl) and the solution stirred vigorously at room temperature for 1h. The solution was concentrated, toluene was added and the mixture concentrated. The residue was then treated with diethyl ether and sonicated

478

to give 349 mg of the title as a yellow solid, 99% yield: 1 H NMR (d 6 -DMSO, δ): consistent with structure; MS(ion spray) 620 (M $^{+}$ +1 of free base); Anal. (C₃₂H₄₆N₇O₇Cl): C,H; N: calcd 14.16, found 13.26.

5

Preparation 415



To a solution of 5-amino-2-napthol (7.96 g, 50.0 mmol) in anhydrous tetrahydrofuran (200 mL) was added benzyl 10 chloroformate (7.14 mL, 50.0 mL) and diisopropylethylamine (9.6 mL, 55.0 mmol). The reaction was stirred at room temperature overnight. The reaction was quenched with 0.1N HCl, concentrated, treated with anhydrous diethyl ether (150 mL), and 11.49 g of gray solid was isolated by filtration. This material was 15 dissolved in anhydrous tetrahydrofuran (120 mL), cooled to -13°C in an ice/acetone bath, treated with sodium hydride (60% dispersion in mineral oil, 1.88 g, 47.0 mmol), and stirred for 30 min. To this solution was 20 added a solution of methyl iodide (2.6 mL, 41.0 mmol) in tetrahydrofuran (30 mL) and the reaction was allowed to warm to room temperature overnight. The reaction was quenched with 0.1N HCl and concentrated. The resulting residue was dissolved in dichloromethane, washed with 1N 25 HCl, saturated aqueous sodium bicarbonate solution, dried over magnesium sulfate, and reconcentrated. Purification by flash chromotography on silica gel (eluting with 20% ethyl acetate/hexane) followed by sonication in diethyl ether gave 2.66 g of the title compound as a white solid,

479

17% yield: ^{1}H NMR (d^{6} -DMSO, d): consistent with structure; MS(ion spray) 308 (M^{*} +1).

Preparation 416

Ö.

5

10

15

To a solution of 5% palladium on carbon (1.00 g) in anhydrous tetrahydrofuran (45 mL) and acetic acid (45 mL) was added the product of Preparation 415 (2.66 g, 8.65 mmol). The mixture was subjected to 60 psi hydrogen on a Parr apparatus for 2 h, then filtered through celite to remove the catalyst. The solvent was removed and the resulting black oil was dissolved in dichloromethane, washed with saturated aqueous sodium bicarbonate solution, dried over magnesium sulfate, and concentrated. Purification by flash chromotography on silica gel (eluting with 20% ethyl acetate/hexanes) gave 537mg of the title compound as a purple solid, 36% yield: ¹H NMR (d⁶-DMSO, d): consistent with structure; MS(ion spray) 619 (M*+1).

20

25

Preparation 417

To a solution of the product of Preparation 396A (1.01 g, 2.0 mmol) in tetrahydrofuran (20 mL) was added the product of Preparation 416 (346 mg, 2.0 mmol), 1-hydroxybenzotriazole (322 mg, 2.1 mmol), and dicyclohexyl carbodiimide (433 mg, 2.1 mmol). The reaction was stirred at room temperature overnight. The insoluble dicyclohexyl urea was removed by filtration, and the

480

filtrate concentrated. The resulting residue was dissolved in ethyl acetate, washed with 1N HCl , saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and reconcentrated. Purification by flash chromotography on silica gel (eluting with 5% methanol/dichloromethane) gave 372 mg of the title compound as a foam, 28% yield: 1 H NMR (6 -DMSO, d): consistent with structure; MS(ion spray) 659 (6 +1); Anal. (6 -DMsO,): C,H,N.

10

481

Example 228

To a solution of the product of Preparation 417 (362 mg, 0.55 mmol) was added acetic acid saturated with HCl_(g) (15 mL, ~3N in HCl) and the mixture stirred vigorously at room temperature for 1h. The solution was concentrated, toluene was added and the mixture concentrated. The residue was then treated with diethyl ether and sonicated to give 351 mg of the title compound as a lavender solid, 100% yield: ¹H NMR (d⁶-DMSO, d): consistent with structure; MS(ion spray) 559 (M^{*} +1 of free base); Anal. (C₃₀H₃₉N₆O₇Cl): C,H,; N: calcd 13.32, found 11.61.

Preparation 418

15

20

25

To a mixture of 1.0 g 10% palladium on carbon in 100 mL tetrahydrofuran was added the product of Preparation 395 (1.78 g, 8.94 mmol). The mixture was subjected to 60psi hydrogen on a Parr apparatus for 2h, then filtered through celite to remove the catalyst. The majority of solvent was removed (50 mL of tetrahydrofuran remained) and to the resulting solution was added the product of Preparation from Examples Part 2A (3.78 g, 10.0 mmol), 1-hydroxybenzotriazole(1.53 g, 10.0 mmol), and dicyclohexyl carbodiimide (2.06 g, 10.0 mmol). The solution was stirred at room temperature overnight. The insoluble dicyclohexyl urea was removed by filtration and the solvent was removed. The resulting oil was dissolved in

482

ethyl acetate, washed with 1N HCl, saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and concentrated. Purification by flash chromotography on silica gel (eluting with 6% methanol/dichloromethane) gave 3.17 g of the title compound as a yellow solid, 67% yield: ¹H NMR (d⁶-DMSO, d): consistent with structure; MS(ion spray) 530 (M⁺ +1); Anal. (C₂₇H₃₉N₅O₆): C,H; N: calcd 13.22, found 12.69.

10

15

Preparation 419

A solution of the product of Preparation 418 (3.17 g, 5.99 mmol) in dioxane (50 mL) was treated with a solution of lithium hydroxide (301 mg, 7.18 mmol) in water (20 mL) After 1h, the reaction was and stirred vigorously. quenched with 1N HCl, and concentrated. The resulting oil was dissolved in ethyl acetate, washed with 1N HCl, water, brine, dried over magnesium sulfate, concentrated to give 2.88 g of the title compound as a white foam, 96% yield: ^{1}H NMR (d^{6} -DMSO, δ): consistent with structure; MS(ion spray) 502 (M+1).

Preparation 420

To a solution of the product of Preparation 419 (1.0 g, 2.0 mmol) stirring in tetrahydrofuran (15 mL) was added the product of Preparation 397 (409 mg, 2.0 mmol), 1-

483

hydroxybenzotriazole (337 mg, 2.2 mmol), and dicyclohexyl carbodiimide (454 mg, 2.2 mmol). The reaction was allowed to warm to room temperature overnight. insoluble dicyclohexyl urea was removed by filtration, and the filtrate was concentrated. The resulting foam was dissolved in ethyl acetate, washed with 1N HCl ,saturated aqueous sodium bicarbonate solution, brine, magnesium dried over sulfate, and concentrated. Purification by flash chromotography on silica gel (eluting with 4-7% methanol/dichloromethane) gave 794 mg of the title compound as a white solid, 58% yield: 1H NMR (d^6-DMSO, δ) : consistent with structure; MS(ion spray) 688 $(M^{+} + 1)$.

15

20

10

To the product of Preparation 420 (785 mg, 1.14 mmol) was added acetic acid saturated with HCl(g) (20 mL, ~3N in and the mixture stirred vigorously HCl) The solution was concentrated , temperature for 1h. toluene was added and the mixture concetnrated. residue was then treated with diethyl ether and sonicated to precipitate give 695 mg of the title compound as a light pink powder after drying, 90% yield: ^{1}H NMR ($d^{6}-$ DMSO, δ): consistent with structure; MS(ion spray) 588 (M⁺ 25 +1 of free base); Anal. $(C_{32}H_{45}N_7O_5Cl_2)$: H,N; C: calcd 56.63, found 57.53.

484

Example 230

To a solution of the product of Preparation 396A (504 mg; 1.0 mmol) in anhydrous acetonitrile (20 mL) was added 5 benzylamine (0.11 mL, 1.0 mmol), 1-hydroxy-7-azabenzo-(150 1.1 mmol), and 1-ethyl-3-(3triazole ma, dimethylaminopropyl)carbodiimide hydrochloride (211mg, 1.1 mmol). The reaction was stirred at room temperature overnight. The solution was concentrated and the 10 resulting residue was dissolved in dichloromethane, washed with 1N HCl, saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, reconcentrated. Purification by flash chromotography on silica gel (eluting with 7% methanol/dichloromethane) 15 gave 356mg of a white solid. The solids were dissolved in acetic acid saturated with $HCl_{(g)}$ (20 mL, ~3N in HCl) and the mixture stirred vigorously at room temperature for 1h. The solution was concentrated, toluene was added and the mixture concentrated. The residue was then 20 treated with diethyl ether and sonicated to precipitate 302 mg of the title compound as a light tan solid, 53% yield: ^{1}H NMR (d^{6} -DMSO, δ): consistent with structure; MS(ion spray) 493 $(M^+ +1)$ of free base); Anal. $(C_{26}H_{34}N_6O_4Cl_2)$: C,H; N: calcd 14.86, found 14.21.

485

Preparation 421

To a solution of the product of Preparation 396A (504mg, 1.0 mmol) in anhydrous acetonitrile (20 mL) was added (S)-(-)-N, alpha-dimethylbenzylamine (0.15 mL, 1.0 mmol), 1-hydroxy-7-azabenzo-triazole (150 mg, 1.1 mmol), and 1ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (211 mg, 1.1 mmol). The reaction was stirred at room temperature overnight. The solution was concentrated and the resulting residue was dissolved in ethyl acetate, washed with 1N HCl, saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, concentrated. Purification by flash chromotography on silica gel (eluting with 5% methanol/1:1 diethyl ether:hexane) gave 289 mg of the title compound as a 15 white solid, 47% yield: 1 H NMR (6 -DMSO, δ): consistent with structure; MS(ion spray) 621 (M+1); $(C_{33}H_{44}N6_{7}O_{6}): C, H, N.$

20

10

Example 231

To a solution of the product of Preparation 421 (279 mg, 0.237 mmol) was added acetic acid saturated with HCl_(g) (20 mL, ~3N in HCl) and the mixture stirred vigorously at room temperature for 1h. The solution was concentrated, 25 toluene was added and the mixture concentrated.

5

residue was then treated with diethyl ether and sonicated to precipitate 235 mg of the title compound as a light red solid, 88% yield: 1H NMR (d^6 -DMSO, δ): consistent with structure; MS(ion spray) 521 (M^+ +1 of free base); Anal. ($C_{28}H_{38}N_6O_4Cl_2$): C,H; N: calcd 14.16, found 13.24.

Preparation 422

To a solution of the product of Preparation 396A (400 mg, 0.79 mmol) in anhydrous acetnitrile (20 mL) was added N-10 benzylmethylamine (0.11 mL, 0.87 mmol), 1-hydroxy-7azabenzo-triazole (119 mg, 0.87 mmol), and 1-ethyl-3- 3dimethylaminopropyl) carbodiimide hydrochloride (168mg, 0.87 mmol). The reaction was stirred at room temperature overnight. The solution was concentrated and the 15 resulting residue was dissolved in dichloromethane, washed with 1N HCl, saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and reconcentrated. Purification by flash chromotography on silica gel (eluting with 2-7.5% methanol/dichloromethane) 20 gave 122 mg of a white solid: ¹H NMR (d^6 -DMSO, δ): consistent with structure; MS(ion spray) 607 (M+1); Anal. $(C_{32}H_{42}N_6O_6)$: H,N; C: calcd 63.35, found 62.77.

25

Example 232

487

To a solution of the product of Preparation 422 (110 mg, 0.18 mmol) was added acetic acid saturated with $HCl_{(g)}$ (20 mL, ~3N in HCl) and the mixture stirred vigorously at room temperature for 1h. The solution was concentrated, toluene was added and the mixture concentrated. The residue was then treated with diethyl ether and sonicated to precipiate 95 mg of the title compound as a white solid, 90% yield: ¹H NMR (d⁶-DMSO, δ): consistent with structure; MS(ion spray) 507 (M⁺ +1 of free base); Anal. (C₂₇H₃₆N₆O₄Cl₂): C,H; N: calcd 14.50, found 13.24.

Preparation 423

To a solution of the product of Preparation 396A (504 mg, 1.0 mmol) in anhydrous tetrahydrofuran (20 mL) was added p-methoxybenzylamine (0.13 mL, 1.0 mmol), 1hydroxybenzotriazole (184 mg, 1.2 mmol), and dicyclohexyl carbodiimide (248 mg, 1.2 mmol). The reaction was stirred at room temperature overnight. Dicyclohexyl urea was removed by filtration and the filtrate was concentrated. The resulting residue was dissolved in dichloromethane, washed with 1N HCl, saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate, and reconcentrated. Purification by flash chromotography on silica gel (eluting with 2-7% 25 methanol/dichloromethane) gave 236 mg of the title compound as a white solid, 38% yield: ^{1}H NMR ($d^{6}-DMSO$, δ): consistent with structure; MS(ion spray) 623 (M+1); Anal. $(C_{32}H_{42}N_6O_7)$: C, H, N.

10

488

Example 233

To the product of Preparation 423 (226 mg, 0.36 mmol) was added acetic acid saturated with $HCl_{(g)}$ (20 mL, ~3N in HCl) and the mixture stirred vigorously at room temperature for 1h. The solution was concentrated, toluene was added and the mixture concentrated. The residue was then treated with diethyl ether and sonicated to precipitate 139 mg of the title compound as a white solid, 65% yield: ¹H NMR (d⁶-DMSO, δ): consistent with structure; MS(ion spray) 523 (M⁺ +1 of free base).

489

EXAMPLES PART 2C

Preparation 424

To a stirring slurry of sodium hydride (0.66 g of a 60% dispersion in mineral oil, 16.5 mmol) in N,Ndimethylformamide (30 mL) at 0° C, was added a solution of 2-methyl-5-nitroimidazole (2.0 g, 15.7 mmol) in N, N-dimethylformamide (10 mL). solution stirred at 0°C for 10 min then a solution of the product of Preparation 5 from Examples Part 1 (4.21 g, 17.3 mmol) in N, N-dimethylformamide (10 mL) was added. The reaction stirred for 3 h at room temperature, at which time it was quenched with brine and extracted with ethyl acetate. combined extracts were washed with brine, dried (sodium sulfate) and evaporated in vacuo to provide a yellow oil. Flash chromatography (silica gel, 10%-50% ethyl acetate/hexanes) yielded the desired product as a light-yellow solid (2.53 g, 60%): NMR consistent with structure; MS (IS) m/e 288 (M -1); Anal. Calc'd for $C_{14}H_{15}N_3O_4$: C, 58.13; H, 5.23; N, 14.53. Found: C, 58.18; H, 5.26; N, 14.56.

490

Preparation 425

To a suspension of 5% palladium on carbon (1.50 g) and dioxane (120 mL), in a Parr reaction bottle, was added the product of Preparation 424 (2.70 g, 9.33 mmol) as a solid. The reaction bottle was placed on a Parr shaker, and shaken at room temperature for 2 h under a hydrogen atmosphere (35 psi). reaction was filtered through a pad of Celite 521 and the filtrate was then added to a previously prepared mixture of the product of Preparation 1d from Examples Part 1 (2.79 g, 7.34 mmol) and 1hydroxybenzotriazole hydrate (1.10 g, 8.10 mmol) in This solution 50 mL dioxane at room temperature. 15 min at which time stirred for dicyclohexylcarbodiimide (1.66 g, 8.10 mmol) was added. The reaction was stirred for 16 h at room temperature, then the solvent was evaporated under The residue was dissolved in reduced pressure. ethyl acetate and filtered to remove the 1,3-The filtrate was purified by dicyclohexylurea. gel, 50% flash chromatography (silica acetate/hexanes - ethyl acetate) to give the desired product as a tan solid foam (3.10 g, 68%): 1H NMR consistent with structure; MS (IS) m/e 622 (M + 1); Anal. Calc'd for $C_{33}H_{43}N_5O_7$: C, 63.75; H, 6.97; N, 11.26. Found: C, 63.46; H, 6.92; N, 11.54.

491

Preparation 426

To a solution of the product of Preparation 425 (2.85 g, 4.59 mmol) in dioxane (100 mL) and water (50 mL) at room temperature was added lithium hydroxide (0.55 g, 13.0 mmol). The reaction stirred 30 min at room temperature, at which time the dioxane was evaporated under reduced pressure. The residue was diluted with water and extracted with diethyl ether (the ether extracts were discarded). The aqueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl ether and ethyl acetate. The combined organic layers were washed with brine, dried (sodium sulfate) and concentrated under reduced pressure to provide the desired product as a light tan solid foam which was used without further purification (2.64 g, 97%): ¹H NMR consistent with structure; MS (IS) m/e 594 (M + 1); Anal. Calc'd for $C_{31}H_{39}N_5O_7$: C, 62.72; H, 6.62; N, 11.80. Found: C, 60.94; H, 6.40; N, 11.50.

492

Preparation 427

To a solution of the product of Preparation 426 1.18 mmol) in anhydrous N,Ndimethylformamide (20 mL) at room temperature was added 1-hydroxybenzotriazole hydrate (0.17 g, 1.22 mmol) and 4-methylpiperidine (0.13 mL, 1.11 mmol). This mixture stirred for 15 min at room temperature, at which time 1,3-dicyclohexylcarbodiimide (0.25 g, 1.22 mmol) was added. The reaction was stirred at room temperature for 15 h, at which time it was quenched with brine and extracted with ethyl The combined extracts were washed with acetate. brine, dried (sodium sulfate) and evaporated to provide a light-brown solid foam. The foam was dissolved in ethyl acetate, the 1,3-dicyclohexylurea was filtered away and the filtrate was purified by flash chromatography (silica gel, 75% acetate/hexanes - 10% methanol/ethyl acetate) to give the desired product as a light tan solid foam (0.475 g, 60%): ¹H NMR consistent with structure; MS (IS) m/e 673 (M - 1); Anal. Calc'd for $C_{37}H_{50}N_6O_6$: C, 65.85; H, 7.47; N, 12.45. Found: C, 65.20; H, 7.26; N, 12.67.

493

Example 234

To a stirring solution of the product of Preparation 427 (0.41 g, 0.61 mmol) and anisole (0.05 mL, 0.47 mmol) in anhydrous dichloromethane (25 mL) at 0°C was added trifluoroacetic acid (3 mL) via syringe. reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate - 5% methanol/ethyl acetate - 5% triethylamine/10% methanol/85% ethyl acetate) to provide the desired product as a white solid foam (0.26 g, 75%): NMR consistent with structure; MS (IS) m/e 574 (M + 1); Anal. Calc'd for $C_{32}H_{42}N_6O_4$: C, 66.88; H, 7.37; N, 14.62. Found: C, 66.64; H, 7.38; N, 14.34.

494

Preparation 428

To a solution of the product of Preparation 427 1.18 mmol) in g, anhydrous N,Ndimethylformamide (25 mL) at room temperature was added 1-hydroxybenzotriazole hydrate (0.17 g, 1.22 mmol) and pyrrolidine (0.09 mL, 1.11 mmol). This mixture stirred for 15 min at room temperature, at which time 1,3-dicyclohexylcarbodiimide (0.25 g, 1.22 mmol) was added. The reaction was stirred at room temperature for 15 h, at which time it was quenched with brine and extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide a light-brown solid foam. The foam was dissolved in ethyl acetate, the 1,3-dicyclohexylurea was filtered away and the filtrate was purified by flash chromatography (silica ael, 75% acetate/hexanes - 10% methanol/ethyl acetate) to give the desired product as a light tan solid foam (0.58 g, 76%): ¹H NMR consistent with structure; MS (IS) m/e 647 (M + 1); Anal. Calc'd for $C_{35}H_{46}N_6O_6$: C, 65.00; H, 7.17; N, 12.99. Found: C, 63.16; H, 6.81; N, 12.91.

495

Example 235

To a stirring solution of the product of Preparation 428 (0.49 g, 0.76 mmol) and anisole (0.05 mL, 0.47 mmol) in anhydrous dichloromethane (25 mL) at 0°C was added trifluoroacetic acid (3 mL) via syringe. reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. The combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate - 5% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide the desired product as a white solid foam (0.27 g, 65%): NMR consistent with structure; MS (IS) m/e 547 (M + Anal. Calc'd for $C_{30}H_{38}N_6O_4$: C, 65.91; H, 7.01; N, 15.37. Found: C, 65.37; H, 6.81; N, 14.83.

496

Preparation 429



To a 0°C solution of 4-methylimidazole (5.00 g, 61 mmol) and concentrated nitric acid (8.22 ml, 183 mmol) was added concentrated sulfuric acid (8.28 ml, 155 mmol), with stirring. After addition was complete, the ice bath was removed and the reaction was heated to a gentle reflux for 2 h. The reaction was allowed to cool and then poured into ice water. The bright yellow precipitate was collected by vacuum filtration and washed thoroughly with water to provide the desired product as a pale yellow solid (5.28 g, 68%): ¹H NMR consistent with structure; MS (FD) m/e 127 (M+); Anal. Calc'd for $C_4H_5N_3O_2$: C, 37.80; H, 3.96; N, 33.06. Found: 37.73; H, 3.79; N, 33.32.

Preparation 430

To a stirring slurry of sodium hydride (1.70 g of a 60% dispersion in mineral oil, 43 mmol) in N,N-dimethylformamide (50 mL) at 0°C, was added a solution of the product of Preparation 429 (5.15 g, 41 mmol) in N,N-dimethylformamide (50 mL). This solution stirred at 0°C for 10 min then a solution of the product of Preparation 5 from Examples Part 1 (10.84 g, 45 mmol) in N,N-dimethylformamide (40 mL)

PCT/US98/17201

was added. The reaction stirred for 1.5 h at room temperature, at which time it was quenched with brine and extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide a yellow oil, which contained a 9:1 mixture of the desired regioisomer and the undesired 5-nitro-4-methyl regioisomer. Flash chromatography (silica gel, 25%-50% ethyl acetate/hexanes) yielded the desired product as a light-yellow solid (7.14 g, 60%): ¹H NMR consistent with structure; MS (IS) m/e 290 (M + 1); Anal. Calc'd for C₁₄H₁₅N₃O₄: C, 58.13; H, 5.23; N, 14.53. Found: C, 58.38; H, 5.33; N, 14.48.

Preparation 431

To a suspension of 5% palladium on carbon (1.10 g) and dioxane (55 mL), in a Parr reaction bottle, was added the product of Preparation 430 (1.55 g, 5.35 mmol) as a solid. The reaction bottle was placed on a Parr shaker, and shaken at room temperature for 3 h under a hydrogen atmosphere (40 psi). The reaction was filtered through a pad of Celite 521 and the filtrate was then added to a previously prepared mixture of the product of Preparation 1d from Examples Part 2A(2.03 g, 5.35 mmol) and 1-hydroxybenzotriazole hydrate (0.80 g, 5.88 mmol) in 25 mL dioxane at room temperature. This solution

15 which time 1,3stirred for min at dicyclohexylcarbodiimide (1.21 g, 5.88 mmol) was The reaction was stirred at room temperature for 15 h, then the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate and the 1,3-dicyclohexylurea was filtered away. The filtrate was purified by flash (silica gel, 50% ethyl chromatography acetate/hexanes - ethyl acetate) to give the desired product as a light tan solid foam (1.91 g, 57%): 1H NMR consistent with structure; MS (IS) m/e 622 (M + 1); Anal. Calc'd for $C_{33}H_{43}N_5O_7$: C, 63.75; H, 6.97; N, 11.26. Found: C, 62.75; H, 6.89; N, 11.76.

Preparation 432

To a solution of the product of Preparation 431 (1.51 g, 2.41 mmol) in dioxane (75 mL) and water (35 mL) at room temperature was added lithium hydroxide (0.31 g, 7.24 mmol). The reaction stirred 25 min at room temperature, at which time the dioxane was evaporated under reduced pressure. The residue was diluted with water and extracted with diethyl ether (the ether extracts were discarded). The aqueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl ether and ethyl acetate. The combined organic layers were washed with brine, dried (sodium sulfate) and concentrated under

499

reduced pressure to provide the desired product as a light tan solid foam that was used without further purification (1.18 g, 83%): ^{1}H NMR consistent with structure; MS (IS) m/e 594 (M + 1).

WO 99/08697

500

Preparation 433

To a solution of the product of Preparation 432 0.84 mmol) in anhydrous dimethylformamide (20 mL) at room temperature was added 1-hydroxybenzotriazole hydrate (0.126 g, 0.93 mmol) and 4-methylpiperidine (0.10 mL, 0.84 mmol). This mixture stirred for 15 min at room temperature, at which time 1,3-dicyclohexylcarbodiimide (0.19 g, 0.93 mmol) was added. The reaction was stirred at room temperature for 15 h, at which time it was quenched with brine and extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide a tan solid foam. The foam was dissolved in ethyl acetate, the 1,3-dicyclohexylurea was filtered and the filtrate was purified by flash chromatography (silica gel, 90% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to give the desired product as an off-white solid foam (0.24 g, 44%): ¹H NMR consistent with structure; MS (IS) m/e 675 (M + 1); Anal. Calc'd for $C_{37}H_{50}N_6O_6$: C, 65.85; H, 7.47; N, 12.45. Found: C, 64.70; H, 6.86; N, 12.49.

501

Example 236

To a stirring solution of the product of Preparation 433 (0.18 g, 0.27 mmol) and anisole (0.05 mL, 0.47 mmol) in anhydrous dichloromethane (10 mL) at 0°C was added trifluoroacetic acid (2 mL) via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. The organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate - 5% methanol/ethyl acetate -5% triethylamine/10% to provide the desired methanol/ethyl acetate) product as an off-white solid foam (0.12 g, 75%): ¹H NMR consistent with structure; MS (IS) m/e 573 (M - 1); Anal. Calc'd for $C_{12}H_{42}N_6O_4$: C, 66.88; H, 7.37; N, 14.62. Found: C, 65.86; H, 7.18; N, 13.82.

Preparation 434

To a solution of the product of Preparation 433 0.84 mmol) in anhydrous N, Ndimethylformamide (20 mL) at room temperature was added 1-hydroxybenzotriazole hydrate (0.13 g, 0.93 mmol) and pyrrolidine(0.07 mL, 0.84 mmol). mixture stirred for 15 min at room temperature, at which time 1,3-dicyclohexylcarbodiimide (0.19 0.93 mmol) was added. The reaction was stirred at room temperature for 15 h, at which time it was quenched with brine and extracted with ethyl The combined extracts were washed with acetate. brine, dried (sodium sulfate) and evaporated to provide a light-brown solid foam. The foam was dissolved in ethyl acetate, the 1,3-dicyclohexylurea was filtered away and the filtrate was purified by flash chromatography (silica gel, 90% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to give the desired product as a off-white solid foam (0.28 g, 50%): ¹H NMR consistent with structure; MS (IS) m/e 647 (M + 1); Anal. Calc'd for $C_{35}H_{46}N_6O_6$: C, 65.00; H, 7.17; N, 12.99. Found: C, 64.08; H, 7.31; N, 12.81.

Example 237

To a stirring solution of the product of Preparation 434 (0.20 g, 0.30 mmol) and anisole (0.05 mL, 0.47 mmol) in anhydrous dichloromethane (10 mL) at 0°C was added trifluoroacetic acid (2 mL) via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate - 5% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide the desired product as an off-white solid foam (0.12 g, 73%): ¹H NMR consistent with structure; MS (FD) m/e 546 (M+); Anal. Calc'd for $C_{30}H_{38}N_6O_4$: C, 65.91; H, 7.01; N, 15.37. Found: C, 65.10; H, 7.10; N, 14.97.

504

Preparation 435



To a 0°C solution of 2-ethylimidazole (10.0 g, 104 mmol) and concentrated nitric acid (15 ml, 312 mmol) was added concentrated sulfuric acid (14 ml, 265 mmol), with stirring. After addition was complete, the ice bath was removed and the reaction was heated to a gentle reflux for 1.5 h. The reaction was allowed to cool and was then poured into ice water and basified with 1N NaOH (pH~6). The aqueous mixture was extracted with ethyl acetate and the combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide an offwhite solid. The crude solid was recrystallized in methanol to give the desired product as a white solid (6.26 g, 43%): ¹H NMR consistent with structure; MS (IS) m/e 142 (M + 1); Anal. Calc'd for $C_5H_7N_3O_2$: C, 42.55; H, 5.00; N, 29.77. Found: C, 42.43; H, 4.99; N, 29.48.

Preparation 436

To a stirring slurry of sodium hydride (01.49 g of a 60% dispersion in mineral oil, 37.2 mmol) in N,N-dimethylformamide (30 mL) at 0° C, was added a solution of the product of Preparation 435 (5.0 g, 35.4 mmol) in N,N-dimethylformamide (30 mL). This

505

solution stirred at 0°C for 10 min then a solution of the product of Preparation 5 from Examples Part 1 (9.50 g, 39.0 mmol) in N, N-dimethylformamide (30 mL) was added. The reaction was stirred at room temperature for 15 h,, at which time it was quenched. with brine and extracted with ethyl acetate. combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide a orange oil. Flash chromatography of the crude oil (silica gel, 20%-60% ethyl acetate/hexanes) yielded the desired product as a light-yellow solid (8.48 g, ¹H NMR consistent with structure; m/e 302 (M - 1); Anal. Calc'd for $C_{15}H_{17}N_3O_4$: C, 59.40; H, 5.65; N, 13.85. Found: C, 59.49; H, 5.43; N, 13.93.

Preparation 437

To a suspension of 5% palladium on carbon (0.75 g) and dioxane (60 mL), in a Parr reaction bottle, was added the product of Preparation 436 (0.93 g, 3.07 mmol) as a solid. The reaction bottle was placed on a Parr shaker, and shaken at room temperature for 2 h under a hydrogen atmosphere (35 psi). The reaction was filtered through a pad of Celite 521 and the filtrate was then added to a previously prepared mixture of the product of Preparation 1d from Examples Part 2A (1.16 g, 3.07 mmol) and 1-

506

hydroxybenzotriazole hydrate (0.46 g, 3.37 mmol) in 30 mL dioxane at room temperature. This solution 15 stirred for min at which time 1,3dicyclohexylcarbodiimide (0.70 g, 3.37 mmol) was The reaction was stirred at room temperature for 15 h,, then the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate and the 1,3-dicyclohexylurea was filtered away. The filtrate was purified by flash chromatography (silica gel, 50% acetate/hexanes - ethyl acetate) to give the desired product as a light orange solid foam (1.18 g, 61%): ¹H NMR consistent with structure; MS (IS) m/e 636 (M + 1); Anal. Calc'd for $C_{34}H_{45}N_5O_7$: C, 64.23; H, 7.13; N, 11.02. Found: C, 63.45; H, 6.70; N, 10.91.

Preparation 438

To a solution of the product of Preparation 437 (1.51 g, 2.38 mmol) in dioxane (75 mL) and water (35 mL) at room temperature was added lithium hydroxide (0.30 g, 7.13 mmol). The reaction stirred 25 min at room temperature, at which time the dioxane was evaporated under reduced pressure. The residue was diluted with water and extracted with diethyl ether (the ether extracts were discarded). The aqueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl ether and ethyl acetate. The

507

combined organic layers were washed with brine, dried (sodium sulfate) and concentrated under reduced pressure to provide the desired product as a light tan solid foam that was used without further purification (1.40 g, 97%): 1 H NMR consistent with structure; MS (IS) m/e 606 (M - 1); Anal. Calc'd for $C_{32}H_{41}N_{5}O_{7}$: C, 63.25; H, 6.80; N, 11.52. Found: C, 61.13; H, 6.44; N, 11.40.

WO 99/08697

508

Preparation 439

To a solution of the product of Preparation 438 0.66 g, mmol) in anhydrous dimethylformamide (20 mL) at room temperature was added 1-hydroxybenzotriazole hydrate (0.10 g, 0.72 mmol) and 4-methylpiperidine (0.08 mL, 0.66 mmol). This mixture stirred for 15 min at room temperature, at which time 1,3-dicyclohexylcarbodiimide (0.15 g, 0.72 mmol) was added. The reaction was stirred at room temperature for 15 h, at which time it was quenched with brine and extracted with ethyl The combined extracts were washed with acetate. brine, dried (sodium sulfate) and evaporated to provide a light-brown solid foam. The foam was dissolved in ethyl acetate, the 1,3-dicyclohexylurea was filtered away and the filtrate was purified by flash chromatography (silica gel, 90% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to give the desried product as a white solid foam (0.372 g, 80%): ¹H NMR consistent with structure; MS (IS) m/e 689 (M + 1); Anal. Calc'd for $C_{38}H_{52}N_6O_6$: C, 66.26; H, 7.61; N, 12.20. Found: C, 65.31; H, 7.15; N, 11.85.

509

Example 238

To a stirring solution of the product of Preparation 439 (0.30 g, 0.43 mmol) and anisole (0.06 mL, 0.58 mmol) in anhydrous dichloromethane (15 mL) at 0°C was added trifluoroacetic acid (3 mL) via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate -10% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide the desried product as an white solid foam (0.19 g, 75%): NMR consistent with structure; MS (FD) m/e 588 (M+); Anal. Calc'd for $C_{33}H_{44}N_6O_4$: C, 67.32; H, 7.53; N, 14.27. Found: C, 67.48; H, 7.32; N, 14.07.

510

Preparation 440

To a solution of the product of Preparation 438 (0.60 g, 1.00 mmol) in anhydrous N,Ndimethylformamide (20 mL) at room temperature was added 1-hydroxybenzotriazole hydrate (0.15 g, 1.09 mmol) and pyrrolidine (0.08 mL, 1.00 mmol). mixture stirred for 15 min at room temperature, at which time 1,3-dicyclohexylcarbodiimide (0.23 g, 1.09 mmol) was added. The reaction was stirred at room temperature for 15 h, at which time it was quenched with brine and extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide a tan solid foam. The foam was dissolved in ethyl acetate, the 1,3-dicyclohexylurea was filtered away and the filtrate was purified by flash chromatography (silica gel, 90% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to give the desired product as a tan solid foam (0.50 g, 76%): ¹H NMR consistent with structure; MS (IS) m/e 661 (M + 1); Anal. Calc'd for $C_{34}H_{45}N_5O_7$: C, 65.43; H, 7.32; N, 12.72. Found: C, 63.85; H, 7.03; N, 12.71.

Example 239

To a stirring solution of the product of Preparation 440~(0.35~g,~0.54~mmol) and anisole (0.07 mL, 0.65 mmol) in anhydrous dichloromethane (15 mL) at 0°C was added trifluoroacetic acid (3 mL) via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate -10% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide the desired product as an off-white solid foam (0.26 g, 86%): ¹H NMR consistent with structure; MS (FD) m/e 560 (M+); Anal. Calc'd for $C_{31}H_{40}N_6O_4$: C, 66.41; H, 7.19; N, 14.99. Found: C, 66.36; H, 7.16; N, 14.78.

Preparation 441

To a 0°C solution of concentrated nitric acid (2.38 ml, 53 mmol) in 85 mL acetic anhydride was added solid 2-phenylimidazole (7.5 g, 52 mmol), with stirring. After addition was complete, the ice bath was removed and the reaction was heated at 95-100°C for 0.5 h. The reaction was allowed to cool and was then poured into ice water and neutralized with 1N The aqueous mixture was extracted with ethyl acetate and the combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide a beige solid. The crude solid was recrystallized in methanol to give the desired product as a off-white solid (1.55 g, 16%): ¹H NMR consistent with structure; MS (IS) m/e 190 (M + 1); Anal. Calc'd for $C_9H_7N_3O_2$: C, 57.14; H, 3.73; N, 22.21. Found: C, 57.17; H, 3.85; N, 21.95.

Preparation 442

To a stirring slurry of sodium hydride (0.30 g of a 60% dispersion in mineral oil, 7.38 mmol) in N,N-dimethylformamide (30 mL) at 0°C, was added a solution of the product of Preparation 441 (1.33 g, 7.03 mmol) in N,N-dimethylformamide (20 mL). This solution stirred at 0°C for 10 min then a solution of

the product of Preparation 6 from Examples Part 1 (1.88 g, 7.73 mmol) in N, N-dimethylformamide (20 mL) was added. The reaction was stirred at room temperature for 15 h,, at which time it was guenched with brine and extracted with ethyl acetate. combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide a yellow oil. Flash chromatography (silica gel, 20%-50% ethyl acetate/hexanes) yielded the desired product as a yellow solid (1.50 g, 61%): ¹H NMR consistent with structure; MS (IS) m/e 352 (M + 1); Calc'd for $C_{19}H_{17}N_3O_4$: C, 64.95; H, 4.88; N, 11.96. Found: C, 65.23; H, 5.04; N, 11.98.

Preparation 443

To a suspension of 5% palladium on carbon (0.68 g) and tetrahydrofuran (75 mL), in a Parr reaction bottle, was added the product of Preparation 442 (1.34 g, 3.81 mmol) as a solid. The reaction bottle was placed on a Parr shaker, and shaken at room temperature for 2 h under a hydrogen atmosphere (40 psi). The reaction was filtered through a pad of Celite 521 and the filtrate was then added to a previously prepared mixture of the product of Preparation 1d from Examples Part 2A(1.45 g, 3.81 mmol) and 1-hydroxybenzotriazole hydrate (0.57 g, 4.20 mmol) in 30 mL tetrahydrofuran at room

temperature. This solution stirred for 15 min at which time 1,3-dicyclohexylcarbodiimide (0.87 4.20 mmol) was added. The reaction was stirred at room temperature for 15 h,, then the solvent was evaporated under reduced pressure. The residue was in dissolved ethyl acetate and the 1,3dicyclohexylurea was filtered away. The filtrate was purified by flash chromatography (silica gel, 50% ethyl acetate/hexanes - ethyl acetate) to give the desired product as an orange-yellow solid foam (2.04 g, 78%): ¹H NMR consistent with structure; MS (IS) m/e 684 (M + 1); Anal. Calc'd for $C_{38}H_{45}N_5O_7$: C, 66.75; H, 6.63; N, 10.24. Found: C, 65.89; H, 6.26; N, 10.12.

Preparation 444

To a solution of the product of Preparation 443 (1.94 g, 2.84 mmol) in dioxane (60 mL) and water (30 mL) at room temperature was added lithium hydroxide (0.48 g, 1.14 mmol). The reaction stirred 25 min at room temperature, at which time the dioxane was evaporated under reduced pressure. The residue was diluted with water and extracted with diethyl ether (the ether extracts were discarded). The aqueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl ether and ethyl acetate. The combined organic layers were washed with brine,

dried (sodium sulfate) and concentrated under reduced pressure to provide the desired product as an orange-yellow solid foam that was used without further purification (1.71 g, 92%): 1 H NMR consistent with structure; MS (IS) m/e 656 (M + 1); Anal. Calc'd for $C_{36}H_{41}N_5O_7$: C, 65.94; H, 6.30; N, 10.68. Found: C, 63.03; H, 5.86; N, 10.60.

Preparation 445

To a solution of the product of Preparation 444 0.99 (0.65)q, mmol) in anhydrous N, Ndimethylformamide (20 mL) at room temperature was added 1-hydroxybenzotriazole hydrate (0.148 g, 1.09 mmol) and pyrrolidine (0.08 mL, 0.99 mmol). mixture stirred for 15 min at room temperature, at which time 1,3-dicyclohexylcarbodiimide (0.23 1.09 mmol) was added. The reaction was stirred at room temperature for 15 h, at which time it was quenched with brine and extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide a light-tan solid foam. The foam was dissolved in ethyl acetate, the 1,3-dicyclohexylurea was filtered away and the filtrate was purified by flash chromatography (silica gel, 90% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to give the desired product as a white solid (0.31 g,

516

44%): 1 H NMR consistent with structure; MS (IS) m/e 709 (M + 1); Anal. Calc'd for $C_{40}H_{48}N_{6}O_{6}$: C, 67.78; H, 6.83; N, 11.86. Found: C, 67.17; H, 6.92; N, 11.67.

Example 240

To a stirring solution of the product of Preparation 445 (0.27 g, 0.38 mmol) and anisole (0.07 mL, 0.65 mmol) in anhydrous dichloromethane (10 mL) at 0°C was added trifluoroacetic acid (1.5 mL) via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. The organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate -10% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide the desired product as a white solid foam (0.23 g, 99%): NMR consistent with structure; MS (IS) m/e 609 (M + 1); Anal. Calc'd for $C_{39}H_{42}N_6O_8F_6$: C, 55.98; H, 5.06; N, 10.04. Found: C, 56.21; H, 5.14; N, 10.12.

WO 99/08697

Preparation 446

To a stirring slurry of sodium hydride (1.12 g of a 60% dispersion in mineral oil, 28 mmol) in N,Ndimethylformamide (50 mL) at 0°C, was added a solution of 3-nitropyrrole (3.00 g, 27 mmol) in N,Ndimethylformamide (30 mL). This solution stirred at 0°C for 10 min then a solution of the product of Preparation 2 from Examples Part 2A (7.31 g, 27 mmol) in N,N-dimethylformamide (30 mL) was added. The reaction was stirred at room temperature for 15 h, at which time it was quenched with brine and extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide an orange-yellow oil. Flash chromatography (silica gel, 30%-50% acetate/hexanes) yielded the desired product as a yellow oil (6.20 g, 75%): ¹H NMR consistent with structure; MS (IS) m/e 303 (M - 1); Anal. Calc'd for $C_{15}H_{16}N_2O_5$: C, 59.21; H, 5.30; N, 9.21. C, 59.20; H, 5.40; N, 9.03.

518

Preparation 447

To a suspension of 5% palladium on carbon (2.00 g) and tetrahydrofuran (100 mL), in a Parr reaction bottle, was added the product of Preparation 446 (4.10 g, 13.5 mmol) as a solid. The reaction bottle was placed on a Parr shaker, and shaken at room temperature for 2 h under a hydrogen atmosphere (40 The reaction was filtered through a pad of Celite 521 and the filtrate was then added to a previously prepared mixture of the product of Preparation 1j from Examples Part 2A (5.10 g, 13.5 mmol), 1,3-dicyclohexylcarbodiimide (3.06 g, mmol) and 1-hydroxybenzotriazole hydrate (2.02 g, 14.8 mmol) in 50 mL tetrahydrofuran at 0°C. The reaction was stirred for 16 h at room temperature, then the solvent was evaporated under reduced The residue was dissolved in ethyl pressure. acetate and the 1,3-dicyclohexylurea was filtered filtrate was away. The purified chromatography (silica gel, 50% ethyl acetate/hexanes - ethyl acetate) to give the desired product as a light orange solid foam (4.00 g, 47%): ¹H NMR consistent with structure; MS (IS) m/e 635 (M + 1); Anal. Calc'd for $C_{35}H_{46}N_4O_7$: C, 66.23; H, 7.30; N, 8.83. Found: C, 66.30; H, 7.31; N, 9.03.

519

Preparation 448

To a solution of the product of Preparation 447 (3.86 g, 6.08 mmol) in tetrahydrofuran (60 mL) and water (30 mL) at room temperature was added lithium hydroxide (0.64 g, 15.2 mmol). The reaction stirred 25 min at room temperature, at which time the tetrahydrofuran was evaporated under reduced pressure. The residue was diluted with water and extracted with diethyl ether (the ether extracts were discarded). The aqueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl ether and ethyl acetate. The combined organic layers were washed with brine, dried (sodium sulfate) and concentrated under reduced pressure to provide the desired product as a light tan solid foam that was used without further purification (3.54 g, 96%): ¹H NMR consistent with structure; MS (IS) m/e 607 (M + 1); Anal. Calc'd for $C_{33}H_{42}N_4O_7$: C, 65.33; H, 6.98; N, 9.23. Found: C, 64.83; H, 6.76; N, 8.65.

520

Preparation 449

To a solution of the product of Preparation 448 (3.46 g, 5.71 mmol) in anhydrous dichloromethane (75 mL) at 0°C was added N-methylmorpholine (0.76 mL, 6.86 mmol) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (1.11 g, 6.29 mmol). This mixture stirred for 1 h, warming to room temperature, at which time 4methylpiperidine (0.75 mL, 6.28 mmol) was added. The reaction stirred for 2 h at room temperature, 2-chloro-4,6-dimethoxy-1,3,5-triazine (0.20 g, 1.14 mmol) was added. The reaction was stirred for another 1 h and the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate, the solids were filtered and the filtrate was purified by flash chromatography (silica gel, ethyl acetate - 10% methanol/ethyl acetate) to give the desired product as an off-white solid foam (0.3.20 g, 82%): ¹H NMR consistent with structure; MS (IS) m/e 688 (M + 1); Anal. Calc'd for $C_{39}H_{53}N_5O_6$: C, 68.10; H, 7.77; N, 10.18. Found: C, 67.55; H, 7.72; N, 10.28.

Example 241 and 242

To a stirring solution of the product of Preparation 449 (3.10 g, 4.51 mmol) and anisole (0.52 mL, 4.73 mmol) in anhydrous dichloromethane (100 mL) at 0°C was added trifluoroacetic acid (12 mL) via syringe. The reaction was stirred for 4 hours warming to room temperature and then quenched by pouring over icecooled saturated sodium bicarbonate. layer was collected and the aqueous layer was extracted twice with dichloromethane. The combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid The impure foam was purified by flash foam. chromatography (silica gel, ethyl acetate - 10% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide the desired product as an off-white solid foam (2.05 g, 77%). ¹H NMR consistent with structure; MS (IS) m/e 586 (M -Anal. Calc'd for $C_{34}H_{45}N_5O_4$: C, 69.48; H, 7.72; N, 11.92. Found: C, 69.27; H, 7.69; N, 11.70. Diastereomeric separation: The product was resolved by HPLC [Kromasil packing material, 15% 3A alcohol/ 85% heptane (w/ 0.2% DMEA)] to provide diastereomers. The first diastereomer (0.80 g) (retention time = 7.13 min) was dissolved in ethyl acetate (20 mL) and then a saturated solution of

522

hydrochloric acid in diethyl ether (3 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided Example 241 (0.79 g) as a white amorphous solid: ¹H NMR consistent with structure; MS (IS) m/e 588 (M + 1); Anal. Calc'd for $C_{34}H_{45}N_5O_4$ 'HCl: C, 65.42; H, 7.42; N, 11.22. Found: C, 64.26; H, 7.37; N, 10.93. The second diastereomer (0.65 g) (retention time = 8.23 min) was dissolved in ethyl acetate (20 mL) and then a saturated solution of hydrochloric acid in diethyl ether (2 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided Example 242 (0.61 g) as a white amorphous solid: ¹H NMR consistent with structure; MS (IS) m/e 588 (M + Anal. Calc'd for $C_{34}H_{45}N_5O_4$ 'HCl: C, 65.42; H, 7.42; N, 11.22. Found: C, 64.52; H, 7.31; N, 10.72.

Preparation 250



To a 0°C solution of pyrazole (20.0 g, 294 mmol) and fuming nitric acid (40 ml, 881 mmol) was added concentrated sulfuric acid (40 ml, 750 mmol), with stirring. After addition was complete, the ice bath was removed and the reaction was heated to a gentle reflux for 1 h. The reaction was allowed to cool and was then poured into ice water, providing an off-white precipitate. The precipitate was collected by vacuum filtration and the crude solid was recrystallized in methanol to give the desired

523

product as a white solid (20.4 g, 61%): 1 H NMR consistent with structure; MS (IS) m/e 114 (M + 1).

Preparation 451

To a stirring slurry of 2-naphthyl acetic acid in carbon tetrachloride (100 mL) at 0°C was added thionyl chloride (64.01 g, 538 mmol). The mixture was heated at 70°C for 1h, then cooled to 0°C . Nbromosuccinimide (57.4 g, 323 mmol), hydrobromic acid (48% aq, 1.7 mL) and carbon tetrachloride (100 mL) were added, and this mixture was heated to reflux for 2 h. After allowing the reaction mixture to cool to room temperature, the succinimide was filtered away and ethanol (150 mL) was added This solution stirred for 1 h at room dropwise. temperature and was then concentrated in vacuo to provide an orange oil. The crude oil was purified by flash chromatography (silica gel; 10% ethyl acetate/hexanes) to provide a yellow oil which solidified upon sitting. The solid was triturated with 5% ethyl acetate/hexanes to provide the desired product as a white solid (37.5 g, 48%): : ¹H NMR consistent with structure; MS (FD) m/e 292,294 (M+); Anal. Calc'd for $C_{14}H_{13}BrO_2$: C, 57.36; H, 4.47. Found: C, 58.27; H, 4.51.

524

Preparation 452

To a stirring slurry of sodium hydride (1.30 g of a 60% dispersion in mineral oil, 32.5 mmol) in N,Ndimethylformamide (35 mL) at 0°C, was added a solution of the product of Preparation 451 (3.50 g, 31.0 mmol) in N,N-dimethylformamide (25 mL). solution stirred at 0°C for 10 min then a solution of the product of Preparation 451 (9.98 g, 34.04 mmol) in N,N-dimethylformamide (50 mL) was added. reaction stirred for 12 h at room temperature, at which time it was quenched with brine and extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate) and evaporated in vacuo to provide a yellow oil. Flash chromatography (silica gel, 15%-40% ethyl acetate/hexanes) yielded the desired product as a light-yellow oil (5.40 g, 54%): ¹H NMR consistent with structure; MS (IS) m/e 326 (M + 1).

Preparation 453

To a suspension of 5% palladium on carbon (5.00 g) and tetrahydrofuran (100 mL), in a Parr reaction

525

bottle, was added the product of Preparation 452 (5.00 g, 15.4 mmol) as a solution in 50 mL tetrahydrofuran. The reaction bottle was placed on a Parr shaker, and shaken at room temperature for 3.5 h under a hydrogen atmosphere (35 psi). reaction was filtered through a pad of Celite 521 and the filtrate was then added to a previously prepared mixture of the product of Preparation 1d from Examples Part 2A (4.00 g, 10.5 mmol), 1,3dicyclohexylcarbodiimide (2.44 g, 11.8 mmol) and 1hydroxybenzotriazole hydrate (1.61 g, 11.8 mmol) in 50 mL tetrahydrofuran at room temperature. reaction was stirred for 16 h at room temperature, then the solvent was evaporated under reduced The residue was dissolved in ethyl pressure. acetate and the remaining solids were filtered away. The filtrate was purified by flash chromatography (silica gel, 70% ethyl acetate/hexanes - ethyl acetate) to give the desired product as a white solid foam (4.75 g, 69%): ¹H'NMR consistent with structure; MS (IS) m/e 658 (M + 1); Anal. Calc'd for $C_{36}H_{43}N_5O_7$: C, 65.74; H, 6.59; N, 10.65. Found: C, 65.11; H, 6.46; N, 10.66.

Preparation 454

To a solution of the product of Preparation 453 (3.95 g, 6.01 mmol) in tetrahydrofuran (60 mL), water (30 mL) and ethanol (10 mL) at room temperature was added lithium hydroxide (1.01 g, 24.0 mmol). The reaction stirred 25 min at room temperature, at which time the tetrahydrofuran was evaporated under reduced pressure. The residue was diluted with water and extracted with diethyl ether (the ether extracts were discarded). The aqueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl ether and ethyl acetate. combined organic layers were washed with brine, dried (sodium sulfate) and concentrated under reduced pressure to provide the desired product as a light tan solid foam that was used without further purification (3.55 g, 94%): H NMR consistent with structure; MS (IS) m/e 630 (M + 1).

Preparation 455

To a solution of the product of Preparation 454 (0.40 g, 0.64 mmol) in anhydrous dichloromethane (20 mL) at 0°C was added N-methylmorpholine (0.08 mL, 0.76 mmol) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (0.12 g, 0.70 mmol). This mixture stirred for 1 h, warming to room temperature, at which time pyrrolidine (0.06 mL, 0.70 mmol) was added. reaction stirred for 4 h at room temperature, then the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate, the solids were filtered away and the filtrate was purified by flash chromatography (silica gel, ethyl acetate - 10% methanol/ethyl acetate) to give the desired product as an off-white solid foam (0.21 g, 48%): 1 H NMR consistent with structure; MS (IS) m/e683 (M + 1);Anal. Calc'd for $C_{34}H_{46}N_{6}O_{6}$: C, 66.84; H, 6.79; N, 12.31. Found: C, 64.83; H, 6.53; N, 12.64.

528

Example 243

To a stirring solution of the product of Preparation 455 (0.19 g, 0.29 mmol) and anisole (0.03 mL, 0.30 mmol) in anhydrous dichloromethane (12 mL) at 0°C was added trifluoroacetic acid (1.8 mL) via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled saturated sodium bicarbonate. The organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate -10% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide an off-white solid foam (0.11 g, 65%). The free based material was dissolved in ethyl acetate (5 mL) and then a saturated solution of hydrochloric acid in diethyl ether (1 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided the desired product (0.12 g) as a white amorphous solid: ¹H NMR consistent with structure; MS (IS) m/e 583 (M + 1); Anal. Calc'd for $C_{33}H_{38}N_6O_4$ 2HCl: C,

529

60.46; H, 6.15; N, 12.82. Found: C, 61.22; H, 6.50; N, 12.75.

Preparation 456

To a solution of the product of Preparation 454 (0.50 g, 0.80 mmol) in anhydrous dichloromethane (25 mL) at 0°C was added N-methylmorpholine (0.11 mL, 0.95 mmol) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (0.15 g, 0.87 mmol). This mixture stirred for 1 h, warming to room temperature, at which time a 2M solution of N, N-dimethylamine (0.83 mL, 1.67 mmol) was added. The reaction stirred for 4 h at room temperature, then the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate, the solids were filtered away and the filtrate was purified by flash chromatography (silica gel, ethyl acetate - 10% methanol/ethyl acetate) to give the desired productas a light yellow solid foam (0.27 g, 52%): ¹H NMR consistent with structure; MS (IS) m/e 657 (M + 1); Calc'd for $C_{36}H_{44}N_6O_6$: C, 65.84; H, 6.75; N, 12.80. Found: C, 63.89; H, 6.65; N, 13.06.

Example 244

To a stirring solution of the product of Preparation 456 (0.26 g, 0.40 mmol) and anisole (0.04 mL, 0.42 mmol) in anhydrous dichloromethane (15 mL) at 0°C was added trifluoroacetic acid (2.3 mL) via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate -10% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide an off-white solid foam (0.16 g, 64%). The free based material was dissolved in ethyl acetate (5 mL) and then a saturated solution of hydrochloric acid in diethyl ether (1 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided desired product (0.17 g) as an off-white amorphous solid: 1H NMR consistent with structure; MS (IS) m/e 557 (M + 1); Anal. Calc'd for

531

 $C_{31}H_{36}N_6O_4$ 2HCl: C, 59.14; H, 6.08; N, 13.35. Found: C, 59.87; H, 6.22; N, 13.16.

Preparation 457

To a stirring slurry of sodium hydride (1.49 g of a 60% dispersion in mineral oil, 37.1 mmol) in N,Ndimethylformamide (35 mL) at 0°C, was added a solution of Preparation 450 (4.00 g, 35.4 mmol) in N, N-dimethylformamide (25 mL). This solution stirred at 0°C for 10 min then a solution of the product of Preparation 2 from Examples Part 2A (10.6 g, 38.9 mmol) in N,N-dimethylformamide (50 mL) was The reaction stirred for 12 h at room added. temperature, at which time it was guenched with brine and extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide a yellow oil. Flash chromatography (silica gel, 15%-40% ethyl acetate/hexanes) yielded the desired product as a light-yellow oil (9.90 g, 92%): ¹H NMR consistent with structure; MS (IS) m/e 304 (M - 1); Anal. Calc'd for $C_{14}H_{15}N_{3}O_{5}$: C, 55.08; H, 4.95; N, 13.76. Found: C, 55.26; H, 4.89; N, 13.62.

Preparation 458

To a suspension of 5% palladium on carbon (2.50 g) and tetrahydrofuran (60 mL), in a Parr reaction bottle, was added the product of Preparation 457 (2.60 g, 8.55 mmol) as a solution in 30 tetrahydrofuran. The reaction bottle was placed on a Parr shaker, and shaken at room temperature for 3.5 h under a hydrogen atmosphere (35 psi). reaction was filtered through a pad of Celite 521 and the filtrate was then added to a previously prepared mixture of the product of Preparation 1d from Examples Part 2A (3.25 g, 8.55 mmol), 1,3dicyclohexylcarbodiimide (1.94 g, 9.40 mmol) and 1hydroxybenzotriazole hydrate (1.28 g, 9.40 mmol) in 35 mL tetrahydrofuran at room temperature. reaction was stirred for 16 h at room temperature, then the solvent was evaporated under reduced The residue was dissolved in ethyl acetate and the solids were filtered away. filtrate was purified by flash chromatography (silica gel, 70% ethyl acetate/hexanes - ethyl acetate) to give the desired product as a white solid foam (4.14 g, 76%): ¹H NMR consistent with structure; MS (IS) m/e 638 (M + 1); Anal. Calc'd for $C_{33}H_{43}N_5O_8$: C, 62.15; H, 6.80; N, 10.98. C, 62.05; H, 6.85; N, 11.07.

Preparation 459

To a solution of the product of Preparation 458 (3.95 g, 6.21 mmol) in tetrahydrofuran (60 mL), water (30 mL) and ethanol (10 mL) at room temperature was added lithium hydroxide (1.04 g, 24.8 mmol). The reaction stirred 25 min at room temperature, at which time the tetrahydrofuran was evaporated under reduced pressure. The residue was diluted with water and extracted with diethyl ether (the ether extracts were discarded). The aqueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl ether and ethyl acetate. combined organic layers were washed with brine, dried (sodium sulfate) and concentrated under reduced pressure to provide the desired product as an off-white solid foam that was used without further purification (3.53 g, 93%): consistent with structure; MS (IS) m/e 610 (M + 1); Anal. Calc'd for $C_{31}H_{39}N_5O_8$: C, 61.07; H, 6.45; N, 11.49. Found: C, 59.51; H, 6.17; N, 11.86.

WO 99/08697

534

PCT/US98/17201

Preparation 460

To a solution of the product of Preparation 458 (0.60 g, 0.99 mmol) in anhydrous dichloromethane (35 mL) at 0°C was added N-methylmorpholine (0.13 mL, 1.18 mmol) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (0.19 g, 1.09 mmol). This mixture stirred for 1 h, warming to room temperature, at which time pyrrolidine (0.09 mL, 1.04 mmol) was added. The reaction stirred for 2 h at room temperature, then the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate, the solids were filtered away and the filtrate was purified by flash chromatography (silica gel, 90% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to give the desired product as an off-white solid 57%): ¹H NMR consistent with foam (0.375 g, structure; MS (IS) m/e 661 (M - 1); Anal. Calc'd for $C_{35}H_{46}N_{6}O_{7}$: C, 63.43; H, 7.00; N, 12.68. C, 63.03; H, 7.00; N, 12.62.

535

Example 245

To a stirring solution of the product of Preparation 460 (0.26 g, 0.39 mmol) and anisole (0.04 mL, 0.42 mmol) in anhydrous dichloromethane (15 mL) at 0°C was added trifluoroacetic acid (2.3 mL) via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate -10% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide a white solid foam (0.17 g, 80%). The free based material was dissolved in ethyl acetate (5 mL) and then a saturated solution of hydrochloric acid in diethyl ether (1 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided the desired product (0.18 g) as a white amorphous solid: ¹H NMR consistent with structure; MS (IS) m/e 563 (M + 1); Anal. Calc'd for $C_{30}H_{38}N_6O_5$ 2HCl: C,

WO 99/08697

56.69; H, 6.34; N, 13.22. Found: C, 58.29; H, 6.29; N, 13.52.

Preparation 461

To a solution of the product of Preparation 459 (0.60 g, 0.99 mmol) in anhydrous dichloromethane (35 mL) at 0°C was added N-methylmorpholine (0.13 mL, 1.18 mmol) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (0.19 g, 1.09 mmol). This mixture stirred for 1 h, warming to room temperature, at which time a 2M solution of N,N-dimethylamine (0.52 mL, 1.04 mmol) was added. The reaction stirred for 2 h at room temperature, then the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate, the solids were filtered away and the filtrate was purified by flash chromatography (silica gel, 90% ethyl acetate/hexanes methanol/ethyl acetate) to give the desired product as an off-white solid foam (0.35 g, 56%): ¹H NMR consistent with structure; MS (IS) m/e 635 (M - 1); Anal. Calc'd for $C_{33}H_{44}N_6O_7$: C, 62.25; H, 6.97; N, 13.20. Found: C, 61.78; H, 6.89; N, 13.05.

WO 99/08697

Example 246

To a stirring solution of the product of Preparation 461 (0.25 g, 0.39 mmol) and anisole (0.04 mL, 0.42 mmol) in anhydrous dichloromethane (15 mL) at 0°C was added trifluoroacetic acid (2.3 mL) via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate -10% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide a white solid foam (0.19 g, 89%). The free based material was dissolved in (5 mL) ethyl acetate and then a saturated solution of hydrochloric acid in diethyl ether (1 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided the desired product (0.19 g) as a white amorphous solid: ¹H NMR consistent with structure; m/e 537 (M + 1); Anal. Calc'd for $C_{28}H_{36}N_6O_5$ 2HCl: C,

538

55.17; H, 6.28; N,13.79. Found: C, 56.27; H, 6.15; N, 14.06.

Preparation 462

To a suspension of 5% palladium on carbon (0.65 g) and tetrahydrofuran (40 mL), in a Parr reaction bottle, was added the product of Preparation 457 (0.65 g, 2.14 mmol) as a solution in 20 mL tetrahydrofuran. The reaction bottle was placed on a Parr shaker, and shaken at room temperature for 2.5 h under a hydrogen atmosphere (35 psi). The reaction was filtered through a pad of Celite 521 and the filtrate was then added to a previously prepared mixture of the product of Preparation 1j from Examples Part 2A (0.81 g, 2.14 mmol), 1,3dicyclohexylcarbodiimide (0.49 g, 2.35 mmol) and 1hydroxybenzotriazole hydrate (0.32 g, 2.35 mmol) in 30 mL tetrahydrofuran at room temperature. reaction was stirred for 16 h at room temperature, then the solvent was evaporated under reduced The residue was dissolved in ethyl pressure. acetate and the solids were filtered away. The flash chromatography filtrate was purified by (silica gel, 70% ethyl acetate/hexanes - ethyl acetate) to give the desired product as a white solid foam (0.82 g, 60%): ¹H NMR consistent with structure; MS (IS) m/e 636 (M + 1); Anal. Calc'd

539

for $C_{34}H_{45}N_5O_7$: C, 64.23; H, 7.13; N,11.02. Found: C, 64.53; H, 7.04; N, 10.97.

Preparation 463

To a solution of the product of Preparation 462(0.72 g, 1.14 mmol) in tetrahydrofuran (30 mL), water (15 mL) and ethanol (5 mL) at room temperature was added lithium hydroxide (0.19 g, 4.54 mmol). The reaction stirred 25 min at room temperature, at which time the tetrahydrofuran was evaporated under reduced The residue was diluted with water and pressure. extracted with diethyl ether (the ether extracts were discarded). The aqueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl ether and ethyl acetate. The combined organic layers were washed with brine, dried (sodium sulfate) and concentrated under reduced pressure to provide the desired product as a white solid that was used without further purification (0.71 g, 99%): ¹H NMR consistent with structure; MS (IS) m/e 606 (M - 1).

540

Preparation 464

To a solution of the product of Preparation 463 (0.20 g, 0.32 mmol) in anhydrous dichloromethane (20 mL) at 0°C was added N-methylmorpholine (0.05 mL, 0.39 mmol) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (0.07 g, 0.35 mmol). This mixture stirred for 1 h, warming to room temperature, at which time pyrrolidine (0.03 mL, 0.35 mmol) was added. The reaction stirred for 2 h at room temperature, then the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate, the solids were filtered away and the filtrate was purified by flash chromatography (silica gel, 90% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to give the desired product as an off-white solid 85%): ¹H NMR consistent with foam (0.18 g, structure; MS (IS) m/e 661 (M + 1); Anal. Calc'd for $C_{36}H_{48}N_6O_6$: C, 65.43; H, 7.32; N, 12.72. C, 64.37; H, 7.24; N, 12.47.

WO 99/08697

541

PCT/US98/17201

Example 247

To a stirring solution of the product of Preparation 464 (0.21 g, 0.31 mmol) and anisole (0.04 mL, 0.42 mmol) in anhydrous dichloromethane (15 mL) at 0°C was added trifluoroacetic acid (1.8 mL) via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate -10% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide a white solid foam (0.15 g, 86%). The free based material was dissolved in ethyl acetate (5 mL) and then a saturated solution of hydrochloric acid in diethyl ether (1 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided the desired product (0.13 g) as a white amorphous solid: ¹H NMR consistent with structure; MS (IS) m/e 561 (M + 1); Anal. Calc'd for $C_{31}H_{40}N_6O_4$ 2HCl: C,

542

58.76; H, 6.68; N, 13.26. Found: C, 59.73; H, 6.63; N, 13.35.

Preparation 465

To a solution of the product of Preparation 463 (0.32 g, 0.52 mmol) in anhydrous dichloromethane (20 mL) at 0°C was added N-methylmorpholine (0.07 mL, 0.62 mmol) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (0.10 g, 0.57 mmol). This mixture stirred for 1 h, warming to room temperature, at which time a 2M solution of N,N-dimethylamine (0.29 mL, 0.57 mmol) was added. The reaction stirred for 2 h at room temperature, then the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate, the solids were filtered away and the filtrate was purified by flash chromatography 90% ethyl acetate/hexanes -(silica gel, methanol/ethyl acetate) to give the desired product as an off-white solid foam (0.19 g, 58%): ¹H NMR consistent with structure; MS (IS) m/e 635 (M + 1); Anal. Calc'd for $C_{34}H_{46}N_6O_6$: C, 64.33; H, 7.30; N, 13.24. Found: C, 63.64; H, 7.35; N, 12.95.

543

Example 248

To a stirring solution of the product of Preparation 465 (0.19 g, 0.29 mmol) and anisole (0.03 mL, 0.33 mmol) in anhydrous dichloromethane (15 mL) at 0°C was added trifluoroacetic acid (1.8 mL) via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. The combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate -10% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide a white solid foam (0.14 q, 88%). The free based material was dissolved in ethyl acetate (5 mL) and then a saturated solution of hydrochloric acid in diethyl ether (1 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided the desired product (0.12 g) as a white amorphous ¹H NMR consistent with structure; solid: m/e 535 (M + 1); Anal. Calc'd for $C_{29}H_{38}N_6O_4$ 2HCl: C,

544

57.33; H, 6.64; N, 13.83. Found: C, 58.11 H, 6.61; N, 13.78.

Preparation 466

To a suspension of 5% palladium on carbon (2.60 g) and tetrahydrofuran (100 mL), in a Parr reaction bottle, was added the product of Preparation 136 from Examples Part 2A (5.00 g, 15.3 mmol) as a The reaction bottle was placed on a Parr shaker, and shaken at room temperature for 2 h under a hydrogen atmosphere (40 psi). The reaction was filtered through a pad of Celite 521 filtrate was then added to a previously prepared mixture of the product of Preparation 1j from 2A (5.80 g, 15.3 Examples Part mmol), dicyclohexylcarbodiimide (3.48 g, 16.9 mmol) and 1hydroxybenzotriazole hydrate (2.29 g, 16.9 mmol) in 50 mL tetrahydrofuran at 0°C. The reaction was stirred for 16 h at room temperature, then the solvent was evaporated under reduced pressure. residue was dissolved in ethyl acetate and the 1,3dicyclohexylurea was filtered away. The filtrate was purified by flash chromatography (silica gel, 80% ethyl acetate/hexanes - 5% methanol/ethyl acetate) to give the desired product as a light yellow solid foam (7.96 g, 80%): ¹H NMR consistent with structure; MS (IS) m/e 656 (M + 1);

545

Calc'd for $C_{37}H_{45}N_5O_6$: C, 67.77; H, 6.92; N, 10.68. Found: C, 67.49; H, 6.88; N, 11.71.

Preparation 467

To a solution of the product of Preparation 466(8.73 g, 13.3 mmol) in tetrahydrofuran (120 mL) and water (60 mL) at room temperature was added lithium hydroxide (2.23 g, 53.2 mmol). The reaction stirred 35 min at room temperature, at which time the tetrahydrofuran was evaporated under pressure. The residue was diluted with water and extracted with diethyl ether (the ether extracts were discarded). The aqueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl The combined organic ether and ethyl acetate. layers were washed with brine, dried sulfate) and concentrated under reduced pressure to provide the desired product as a light yellow solid foam that was used without further purification (8.18 g, 98%): ¹H NMR consistent with structure; MS (IS) m/e 628 (M + 1); Anal. Calc'd for $C_{35}H_{41}N_5O_6$: C, 66.97; H, 6.58; N, 11.16. Found: C, 66.68; H, 6.75; N, 11.12.

546

Preparation 468

To a solution of the product of Preparation 467 (2.90 g, 4.61 mmol) in anhydrous dichloromethane (40 mL) at 0°C was added N-methylmorpholine (0.61 mL, 5.53 mmol) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (1.05 g, 5.99 mmol). This mixture stirred for 1 h, warming to room temperature, at which time 4methylpiperidine (0.60 mL, 5.07 mmol) was added. The reaction stirred for 2 h at room temperature, 2-chloro-4,6-dimethoxy-1,3,5-triazine then more (0.20 g) was added. The reaction was stirred for another 1 h and the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate, the solids were filtered away and the filtrate was purified by flash chromatography (silica gel, ethyl acetate - 10% methanol/ethyl acetate) to give the desired product as a light yellow solid foam (2.99 g, 92%): ¹H NMR consistent with structure; MS (IS) m/e 709 (M + 1); Anal. Calc'd for $C_{41}H_{52}N_6O_5$: C, 69.47; H, 7.39; N, 11.85. Found: C, 69.30; H, 7.47; N, 11.92.

547

Example 249 and 250

To a stirring solution of the product of Preparation 468 (4.40 g, 6.20 mmol) and anisole (0.71 mL, 6.50 mmol) in anhydrous dichloromethane (140 mL) at 0°C was added trifluoroacetic acid (14 mL) via syringe. The reaction was stirred for 4 h warming to room temperature and then was quenched by pouring over ice-cooled saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, 5% methanol/ethyl triethylamine/10% 5% methanol/ethyl acetate) to provide the desired product as a light yellow solid foam (3.75 g, 99%). ¹H NMR consistent with structure; MS (IS) m/e 609 (M + 1); Anal. Calc'd for $C_{36}H_{44}N_6O_3$: C, 71.03; H, 7.29; N, 13.80. Found: C, 69.83; H, 7.17; N, 13.54.

Diastereomeric separation: the desired product was resolved by HPLC [Kromasil packing material, 15% 3A alcohol/ 85% heptane (w/ 0.2% DMEA)] to provide two diastereomers. The first diastereomer (1.30 g) (retention time = 6.77 min) was dissolved in ethyl acetate (20 mL) and then a saturated solution of

hydrochloric acid in diethyl ether (3 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided Example 249 (1.10 g) as a white amorphous solid: ¹H NMR consistent with structure; MS (IS) m/e 609 (M + 1); Anal. Calc'd for C₃₆H₄₄N₆O₃'HCl: C, 67.02; H, 7.03; N. Found: C, 66.53; H, 6.96; N, 12.80. The second diastereomer (1.50 g) (retention time = 9.17 min) was dissolved in ethyl acetate (20 mL) and then a saturated solution of hydrochloric acid in diethyl ether (3 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided Example 250 (1.47 g) as a white amorphous solid: ¹H NMR consistent with structure; MS (IS) m/e 609 (M Anal. Calc'd for $C_{36}H_{44}N_6O_3$ 'HCl: C, 67.02; H, 7.03; N, 13.03. Found: C, 66.08; H, 6.95; N, 12.71.

Preparation 469

To a solution of the product of Preparation 467 (5.10 g, 8.11 mmol) in anhydrous dichloromethane (75 mL) at 0°C was added N-methylmorpholine (1.07 mL, 9.73 mmol) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (1.85 g, 10.5 mmol). This mixture stirred for 1 h, warming to room temperature, at which time

549

pyrrolidine (0.75 mL, 8.93 mmol) was added. The reaction stirred for 2 h at room temperature, then more 2-chloro-4,6-dimethoxy-1,3,5-triazine (0.20 g) was added. The reaction was stirred for another 1 h and the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate, the solids were filtered away and the filtrate was purified by flash chromatography (silica gel, ethyl acetate - 10% methanol/ethyl acetate) to give the desired product as a light yellow solid foam (5.30 g, 96%): 1 H NMR consistent with structure; MS (IS) m/e 681 (M + 1); Anal. Calc'd for $C_{39}H_{48}N_6O_5$: C, 68.80; H, 7.11; N, 12.34. Found: C, 68.07; H, 7.10; N, 12.85.

550

Example 251 and 252

To a stirring solution of the product of Preparation 469 (5.15 g, 7.55 mmol) and anisole (0.86 mL, 7.93 mmol) in anhydrous dichloromethane (150 mL) at 0°C was added trifluoroacetic acid (15 mL) via syringe. The reaction was stirred for 4 h warming to room temperature and then was quenched by pouring over ice-cooled saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, 5% methanol/ethyl 5% triethylamine/10% acetate methanol/ethyl acetate) to provide the desired product as an offwhite solid foam (4.11 g, 94%). H NMR consistent with structure; MS (IS) m/e 581 (M + 1); Anal. Calc'd for $C_{34}H_{40}N_6O_3$: C, 70.32; H, 6.94; N, 14.47. Found: C, 70.34; H, 6.79; N, 13.70.

Diastereomeric separation: the desired product was resolved by HPLC [Kromasil packing material, 15% 3A alcohol/ 85% heptane (w/ 0.2% DMEA)] to provide two diastereomers. The first diastereomer (1.70 g) (retention time = 7.72 min) was dissolved in ethyl acetate (20 mL) and then a saturated solution of

551

hydrochloric acid in diethyl ether (3 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided Example 251 (1.27 g) as a off-white amorphous solid: ¹H NMR consistent with structure; MS (IS) m/e 581 (M + 1); Anal. Calc'd for $C_{34}H_{40}N_6O_3$ 2HCl: C, 66.17; H, 6.70; N, 13.62. C, 65.65; H, 6.90; N, 13.48. Found: The second diastereomer (1.40 g) (retention time = 10.81) was dissolved in ethyl acetate (20 mL) and then a saturated solution of hydrochloric acid in diethyl ether (3 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided Example 252 (1.47 g) as a off-white amorphous solid: ¹H NMR consistent with structure; MS (IS) m/e 581 (M + 1); Anal. Calc'd for $C_{34}H_{40}N_6O_3$ 2HCl: C, 66.17; H, 6.70; N, 13.62. Found: C, 65.73; H, 7.03; N, 13.31.

Preparation 470

To a suspension of 5% palladium on carbon (2.60 g) and tetrahydrofuran (100 mL), in a Parr reaction bottle, was added the product of Preparation 99 from Examples Part 2A (5.00 g, 15.3 mmol) as a solid. The reaction bottle was placed on a Parr shaker, and

552

shaken at room temperature for 2 h under a hydrogen atmosphere (40 psi). The reaction was filtered through a pad of Celite 521 and the filtrate was then added to a previously prepared mixture of the product of Preparation 1j from Examples Part 2A (5.80 g, 15.3 mmol), 1,3-dicyclohexylcarbodiimide (3.48 g, 16.9 mmol) and 1-hydroxybenzotriazole hydrate (2.29 g, 16.9 mmol) in 50 mL tetrahydrofuran The reaction was stirred for 16 h at room temperature, then the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate and the 1,3-dicyclohexylurea was filtered away. The filtrate was purified by flash (silica chromatography gel, 808 ethyl acetate/hexanes - 5% methanol/ethyl acetate) to give the desired product as a light yellow solid foam (7.96 g, 80%): ¹H NMR consistent with structure; MS (IS) m/e 682 (M + 1); Anal. Calc'd for $C_{39}H_{47}N_5O_6$: C, 68.70; H, 6.95; N, 10.27. Found: C, 68.27; H, 6.86; N, 10.77.

553

Preparation 471

To a solution of the product of Preparation 470 (8.73 g, 13.3 mmol) in tetrahydrofuran (120 mL) and water (60 mL) at room temperature was added lithium hydroxide (2.23 g, 53.2 mmol). The reaction stirred 35 min at room temperature, at which time the tetrahydrofuran was evaporated under reduced The residue was diluted with water and pressure. extracted with diethyl ether (the ether extracts were discarded). The aqueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl ether and ethyl acetate. The combined organic layers were washed with brine, dried (sodium sulfate) and concentrated under reduced pressure to provide the desired product as a light yellow solid foam that was used without further purification (8.18 g, 98%): ¹H NMR consistent with structure; MS (IS) m/e 654 (M + 1); Anal. Calc'd for $C_{37}H_{43}N_5O_6$: C, 67.98; H, 6.63; N, 10.71. Found: C, 66.83; H, 6.59; N, 10.50.

554

Preparation 472

To a solution of the product of Preparation 471 (1.00 g, 1.52 mmol) in anhydrous dichloromethane (30 mL) at 0°C was added N-methylmorpholine (0.20 mL, 1.82 mmol) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (0.35 g, 1.98 mmol). This mixture stirred for 1 h, warming to room temperature, at which time a 2M solution of N,N-dimethylamine (0.84 mL, 1.68 mmol) was added. The reaction stirred for 2 h at room temperature, then more 2-chloro-4,6-dimethoxy-1,3,5triazine (0.08 g) was added. The reaction was stirred for another 1 h and the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate, the solids were filtered away and the filtrate was purified by flash chromatography (silica gel, 90% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to give the desired product as a light yellow solid ¹H NMR consistent with foam (0.83 g, 80%): structure; MS (IS) m/e 681 (M + 1); Anal. Calc'd for $C_{39}H_{48}N_6O_5$: C, 68.80; H, 7.11; N, 12.34. Found: C, 68.23; H, 7.03; N, 12.66.

555

Example 253

To a stirring solution of the product of Preparation 472 (3.10 g, 4.52 mmol) and anisole (0.52 mL, 4.75 mmol) in anhydrous dichloromethane (100 mL) at 0°C was added trifluoroacetic acid (10 mL) via syringe. The reaction was stirred for 4 h warming to room temperature and then was quenched by pouring over ice-cooled saturated sodium bicarbonate. organic layer was collected and the aqueous laver was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, 5% methanol/ethyl 5% triethylamine/10% methanol/ethyl acetate acetate) to provide the desired product as an offwhite solid foam (2.40 g, 91%). H NMR consistent with structure; MS (IS) m/e 581 (M + 1); Calc'd for $C_{34}H_{40}N_6O_3$: C, 70.32; H, 6.94; N, 14.47. Found: C, 69.36; H, 6.71; N, 14.10.

Diastereomeric separation: the desired product was resolved by HPLC [Kromasil packing material, 15% 3A alcohol/ 85% heptane (w/ 0.2% DMEA)] to provide two diastereomers. The second diastereomer (0.76 g) (retention time = 9.98 min) was dissolved in ethyl

556

acetate (15 mL) and then a saturated solution of hydrochloric acid in diethyl ether (2 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided Example 253 (0.70 g) as a white amorphous solid: 1 H NMR consistent with structure; MS (IS) m/e 581 (M + 1); Anal. Calc'd for $C_{34}H_{40}N_6O_3$ HCl: C, 66.18; H, 6.70; N, 13.62. Found: C, 64.39; H, 6.69; N, 13.19.

Preparation 473

To a solution of the product of Preparation 5 from Examples Part 2A (3.60 g, 5.90 mmol) in anhydrous dichloromethane (60 mL) at 0°C was added methylmorpholine (0.78 mL, 7.08 mmol) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (1.35 g, 7.67 mmol). This mixture stirred for 1 h, warming to room temperature, at which time a 2M solution of N,Ndimethylamine (3.30 mL, 6.49 mmol) was added. reaction stirred for 2 h at room temperature, then more 2-chloro-4,6-dimethoxy-1,3,5-triazine (0.30 g) was added. The reaction was stirred for another 1 h and the solvent was evaporated under reduced The residue was dissolved in ethyl pressure. acetate, the solids were filtered away and the filtrate was purified by flash chromatography (silica gel, ethyl acetate - 10% methanol/ethyl

557

acetate) to give the desired product as a light yellow solid foam (2.93 g, 78%): 1 H NMR consistent with structure; MS (IS) m/e 637 (M + 1); Anal. Calc'd for $C_{33}H_{44}N_6O_7$: C, 62.25; H, 6.97; N, 13.20. Found: C, 61.02; H, 6.67; N, 13.72.

Example 254

To a stirring solution of the product of Preparation 473 (3.60 g, 5.65 mmol) and anisole (0.65 mL, 5.93 mmol) in anhydrous dichloromethane (130 mL) at 0°C was added trifluoroacetic acid (13 mL) via syringe. The reaction was stirred for 4 h warming to room temperature and then was quenched by pouring over ice-cooled saturated sodium bicarbonate. The organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, 5% methanol/ethyl 5% triethylamine/10% methanol/ethyl acetate) to provide the desired product as an offwhite solid foam (2.20 g, 73%). ¹H NMR consistent with structure; MS (IS) m/e 537 (M + 1); Anal. Calc'd for $C_{28}H_{36}N_6O_5$: C, 62.67; H, 6.76; N, 15.66. Found: C, 62.53; H, 6.62; N, 15.57.

558

Diastereomeric separation: the product was resolved by HPLC [Kromasil packing material, 15% 3A alcohol/ (w/ 0.2% DMEA)] to provide 85% heptane diastereomers. The second diastereomer (0.45 g) (retention time = 10.70 min) was dissolved in ethyl acetate (10 mL) and then a saturated solution of hydrochloric acid in diethyl ether (2 mL) was added, with stirring. The white precipitate was collected by vacuum filtration and rinsed with diethyl ether. Vacuum drying provided Example 254 (0.40 g) as a white amorphous solid: ¹H NMR consistent with structure; MS (IS) m/e 537 (M + 1); Anal. Calc'd for $C_{28}H_{36}N_6O_5$ 2HCl: C, 55.17; H, 6.28; N, 13.79. Found: C, 56.56; H, 6.38; N, 14.26.

Preparation 474

To a stirring slurry of sodium hydride (0.155 g of a 60% dispersion in mineral oil, 3.86 mmol) in N,N-dimethylformamide (30 mL) at 0oC, was added a solution of 3-nitropyrrole (0.600 g, 3.68 mmol) in N,N-dimethylformamide (10 mL). This solution stirred at 0oC for 10 min then a solution of the product of Preparation 5, Examples Part 1, in N,N-dimethylformamide (10 mL) was added. The reaction was stirred at room temperature for 15 h,, at which

PCT/US98/17201

time it was quenched with brine and extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide a yellow oil, which contained a 9:1 mixture of the desired regioisomer and the undesired 5-nitro-4-methyl regioisomer. Flash chromatography (silica gel, 20%-50% ethyl acetate/hexanes) yielded the desired product as a light-yellow solid (2.53 g, 60%).

Preparation 475

To a stirring solution of the product of Preparation 474 (1.01 g, 1.59 mmol) in dioxane (30 mL) and water (15 mL) at room temperature was added lithium hydroxide (0.26 g, 6.30 mmol). The reaction stirred 25 min at room temperature, at which time the dioxane was evaporated under reduced pressure. The residue was diluted with water and extracted with diethyl ether (the ether extracts were discarded). The aqueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl ether and ethyl acetate. The combined organic layers were washed with brine, dried (sodium sulfate) and concentrated under reduced pressure to provide the desired product as a light tan solid foam that was used without further purification (0.96 g, 99%):

560

Preparation 476

To a stirring solution of Preapation 475 (0.433 g, 1.46 mmol), L-proline methyl ester hydrochloride (0.241 g, 1.46 mmol), 1-hydroxybenzotriazole hydrate (0.200 g, 1.46 mmol) and N, N-diisopropylethylamine (0.660 g, 5.10 mmol) in anhydrous 1,2-dichlormethane (30 mL), at room temperature, was added 1-(3dimethylaminopropyl)-3-ethylcarbodiimide (0.310 1.60 mmol). The reaction was stirred for 18 h at room temperature, quenched with water (50 mL) and the aqueous layer was extracted with ethyl acetate. The combined extracts were washed with 10% citric acid, saturated sodium bicarbonate, water and brine, dried (sodium sulfate) and evaporated to provide a yellow solid. Rotary chromatography (silica gel, ethyl acetate/hexanes 85% ethyl acetate/hexanes) gave 0.250 g (42%) of the desired productas a white foam solid.

561

Preparation 477

To a suspension of 5% palladium on carbon (1.75 g) ethyl acetate (90 mL) and ethanol (10 mL), in a Parr reaction bottle. was added the product Preparation 476 (3.51 g, 11.5 mmol) as a solid. reaction bottle was placed on a Parr shaker, and shaken at room temperature for 45 min under a hydrogen atmosphere (35 psi). The reaction was filtered through a pad of Celite 521. The filtrate was evaporated to an orange solid foam which was then added to a previously prepared mixture of the product of Preparation 1d (4.33 g, 11.5 mmol) and 1hydroxybenzotriazole hydrate (1.72 g, 12.6 mmol) in 50 mL dioxane at room temperature. This solution stirred for 15 min at which time dicyclohexylcarbodiimide (1.66 g, 10.3 mmol) was added. The reaction was stirred at room temperature for 15 h, then the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate and the 1,3-dicyclohexylurea was filtered away. The filtrate was purified by flash chromatography (silica gel, 75% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to

562

give the desired product as a light tan solid foam (4.5 g, 62%):

Example 255

To a stirring solution of the product of Preparation 477 (0.77 g, 1.12 mmol) and anisole (0.05 mL, 0.47 mmol) in anhydrous dichloromethane (20 mL) at 0oC was added trifluoroacetic acid via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. The organic layer was collected and the aqueous layer was extracted twice with dichloromethane. The combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, 100% ethyl acetate 95% 5% methanol/ ethyl acetate triethylamine/10% methanol/ 85% ethyl acetate) to provide the desired product as an off-white solid foam (0.625 g, 95%).

563

Preparation 478

To a stirring solution of the product of Preparation 475 (0.433 g, 1.46 mmol), pyrrolidine (0.241 g, 1.46 mmol), 1-hydroxybenzotriazole hydrate (0.200 g, 1.46 mmol) and N,N-diisopropylethylamine (0.660 g, 5.10 mmol) in anhydrous 1,2-dichlormethane (30 mL), at added room temperature, 1 - (3 was dimethylaminopropyl)-3-ethylcarbodiimide (0.310 The reaction was stirred for 18 h at 1.60 mmol). room temperature, quenched with water (50 mL) and the aqueous layer was extracted with ethyl acetate. The combined extracts were washed with 10% citric acid, saturated sodium bicarbonate, water and brine, dried (sodium sulfate) and evaporated to provide a yellow solid. Flash chromatography (silica gel, 40% ethyl acetate/hexanes - 100% ethyl acetate) gave 0.250 g (42%) of the desired product as a white foam solid.

564

Preparation 479

To a suspension of 5% palladium on carbon (1.75 g) ethyl acetate (50 mL) and ethanol (25 mL), in a Parr reaction bottle, was added Preparation 478 (3.51 g, 11.5 mmol) as a solid. The reaction bottle was placed on a Parr shaker, and shaken at room temperature for 45 min under a hydrogen atmosphere The reaction was filtered through a pad (35 psi). of Celite 521. The filtrate was evaporated to an orange solid foam which was then added to previously prepared mixture of the product of Preparation 1d (4.33)g, 11.5 mmol) 1hydroxybenzotriazole hydrate (1.72 g, 12.6 mmol) in 50 mL dioxane at room temperature. This solution 15 min at which time 1.3stirred for dicyclohexylcarbodiimide (1.66 g, 10.3 mmol) was The reaction was stirred at room temperature for 15 h, then the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate and the 1,3-dicyclohexylurea was filtered away. The filtrate was purified by flash chromatography (silica gel, 50% ethyl acetate/hexanes - 100% ethyl acetate) to give the desired product as a light tan solid foam (4.5 g, 62%).

565

Example 256

To a stirring solution of the product of Preparation 479 (0.77 g, 1.12 mmol) and anisole (0.05 mL, 0.47 mmol) in anhydrous dichloromethane (20 mL) at 0oC was added trifluoroacetic acid via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. The organic layer was collected and the aqueous layer was extracted twice with dichloromethane. The combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, 100% ethyl acetate - ′ 5% methanol/ 95% ethyl acetate 5% triethylamine/10% methanol/ 85% ethyl acetate) to provide the desired product as an off-white solid foam (0.625 g, 95%).

566

Preparation 480

To a stirring slurry of sodium hydride (0.155 g of a 60% dispersion in mineral oil, 3.86 mmol) in N,Ndimethylformamide (30 mL) at 0oC, was added a solution of 3-nitropyrrole (0.600 g, 3.68 mmol) in N, N-dimethylformamide (10 mL). This solution stirred at OoC for 10 min then a solution of the product of Preparation 5, Examples Part 1 in N,Ndimethylformamide (10 mL) was added. The reaction was stirred at room temperature for 15 h,, at which time it was quenched with brine and extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide a yellow oil, which contained a 9:1 mixture of the desired regioisomer and the undesired 5nitro-4-methyl regioisomer. Flash chromatography (silica gel, 20%-50% ethyl acetate/hexanes) yielded the desired product as a light-yellow solid (2.53 g, 60%).

567

Preparation 481

To a stirring solution of the product of Preparation 480 (1.01 g, 1.59 mmol) in dioxane (30 mL) and water (15 mL) at room temperature was added lithium hydroxide (0.26 g, 6.30 mmol). The reaction stirred 25 min at room temperature, at which time the dioxane was evaporated under reduced pressure. residue was diluted with water and extracted with diethyl ether (the ether extracts were discarded). The agueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl ether and ethyl acetate. The combined organic layers were washed with brine, dried (sodium sulfate) and concentrated under reduced pressure to provide the desired product as a light tan solid foam that was used without further purification (0. 96 g, 99%).

Preparation 482

To a stirring solution of the product of Preparation 481 (0.433 g, 1.46 mmol), L-proline methyl ester hydrochloride (0.241 g, 1.46 mmol), hydroxybenzotriazole hydrate (0.200 g, 1.46 mmol) and N,N-diisopropylethylamine (0.660 g, 5.10 mmol) in anhydrous 1,2-dichlormethane (30 mL), at room temperature, was added 1-(3-dimethylaminopropyl)-3ethylcarbodiimide (0.310 g, 1.60 mmol). reaction was stirred for 18 h at room temperature, quenched with water (50 mL) and the aqueous layer was extracted with ethyl acetate. The combined extracts were washed with 10% citric acid, saturated sodium bicarbonate, water and brine, dried (sodium sulfate) and evaporated to provide a yellow solid. Rotary chromatography (silica gel, 35% ethyl acetate/hexanes - 85% ethyl acetate/hexanes) gave 0.250 g (42%) of the desired product as a white foam solid.

569

Preparatin 483

To a suspension of 5% palladium on carbon (1.75 g) methanol (25 mL), was added the product of Preparation 482 (3.51 g, 11.5 mmol) as a solid. reaction was stirred at room temperature for 3-4 h under 1 atmosphere of H2. The reaction was filtered through a pad of Celite 521. The filtrate was evaporated to an off-white solid foam which was then added to a previously prepared mixture of the product of Preparation 1d (4.33 g, 11.5 mmol) and 1hydroxybenzotriazole hydrate (1.72 g, 12.6 mmol) in 50 mL N, N-dimethylformamide at room temperature. This solution stirred for 15 min at which time 1,3dicyclohexylcarbodiimide (1.66 g, 10.3 mmol) was added. The reaction was stirred at room temperature for 15 h,, then guenched with brine and extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate), and evaporated to give a tan solid foam. The crude foam was purified by flash chromatography (silica gel, 75% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to provide the separated diastereomers, Preparation 483A and Preparation 483B as off-white solid foams (4.5 g, 62%).

Example 257

570

To a stirring solution of the product of Preparation 483A (0.77 g, 1.12 mmol) and anisole (0.05 mL, 0.47 mmol) in anhydrous dichloromethane (20 mL) at 0oC was added trifluoroacetic acid via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, 100% ethyl acetate - 5% methanol/ 95% ethyl acetate triethylamine/10% methanol/ 85% ethyl acetate) to provide the desired product as an white solid foam (0.625 g, 95%):

571

Example 258

To a stirring solution of the product of Preparation 483B (0.77 g, 1.12 mmol) and anisole (0.05 mL, 0.47 mmol) in anhydrous dichloromethane (20 mL) at 0oC was added trifluoroacetic acid via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. The organic layer was collected and the aqueous layer was extracted twice with dichloromethane. The combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, 100% ethyl acetate

- 5% methanol/ 95% ethyl acetate - 5% triethylamine/10% methanol/ 85% ethyl acetate) to provide the desired product as an white solid foam (0.625 g, 95%).

572

Preparation 484

To a stirring solution of the product of Preparation 481 (0.433 g, 1.46 mmol), L-proline methyl ester hydrochloride (0.241 g, 1.46 mmol), 1hydroxybenzotriazole hydrate (0.200 g, 1.46 mmol) and N, N-diisopropylethylamine (0.660 g, 5.10 mmol) in anhydrous 1,2-dichlormethane (30 mL), at room temperature, was added 1-(3-dimethylaminopropyl)-3ethylcarbodiimide (0.310 g, 1.60 mmol). reaction was stirred for 18 h at room temperature, quenched with water (50 mL) and the aqueous layer was extracted with ethyl acetate. The combined extracts were washed with 10% citric acid, saturated sodium bicarbonate, water and brine, dried (sodium sulfate) and evaporated to provide a yellow solid. Rotary chromatography (silica gel, 35% acetate/hexanes - 85% ethyl acetate/hexanes) gave 0.250 g (42%) of the desired product as a white foam solid.

573

Preparation 485

To a suspension of 5% palladium on carbon (1.75 g) methanol (25 mL), was added the product of Preparation 484 (3.51 g, 11.5 mmol) as a solid. reaction was stirred at room temperature for 3-4 h under 1 atmosphere of H_2 . The reaction was filtered through a pad of Celite 521. The filtrate was evaporated to an off-white solid foam which was then added to a previously prepared mixture of product of Preparation 1d (4.33 g, 11.5 mmol) and 1hydroxybenzotriazole hydrate (1.72 g, 12.6 mmol) in 50 mL N, N-dimethylformamide at room temperature. This solution stirred for 15 min at which time 1,3dicyclohexylcarbodiimide (1.66 g, 10.3 mmol) was added. The reaction was stirred at room temperature for 15 h,, then quenched with brine and extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate), evaporated to give a tan solid foam. The crude foam was purified by flash chromatography (silica gel, 75% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to provide the desired product as an offwhite solid foam (4.5 g, 62%).

574

Example 259

To a stirring solution of the product of Preparation 485~(0.77~g,~1.12~mmol) and anisole (0.05 mL, 0.47 mmol) in anhydrous dichloromethane (20 mL) at 0oC was added trifluoroacetic acid via syringe. The reaction was stirred for 4 h, warming to room temperature and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. organic layer was collected and the aqueous layer was extracted twice with dichloromethane. The combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, 100% ethyl acetate 5% methanol/ 95% ethyl acetate triethylamine/10% methanol/ 85% ethyl acetate) provide the desired product as a white solid foam (0.625 g, 95%).

575

EXAMPLES PART 2D

Preparation 486

5 To a stirring slurry of sodium hydride (0.750 g of a 60% dispersion in mineral oil, 18.7 mmol) in N,N-dimethylformamide (40 mL) at 0°C, was added a solution of 3-nitropyrrole (2.00 g, 17.8 mmol) in N, N-dimethylformamide (25 mL). This solution 10 stirred at 0°C for 10 min then a solution of the product of Preparation 5 from Examples Part 1 (4.77 g, 19.6 mmol) in N,N-dimethylformamide (10 mL) was added. The reaction was stirred at room temperature for 15 h, at which time it was quenched with brine and extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide a yellow oil. Flash chromatography (silica gel, 20%-50% ethyl acetate/hexanes) yielded the desired product as a light-yellow oil (4.77 g, 98%): ¹H NMR consistent 20 with structure; MS (IS) m/e 275 (M + 1).

Preparation 487

To a stirring solution of the product of Preparation 486 (4.70 g, 17.1 mmol) in dioxane (100 mL) and

576

water (50 mL) at room temperature was added lithium hydroxide (2.88 g, 68.5 mmol). The reaction stirred 25 min at room temperature, at which time the dioxane was evaporated under reduced pressure. The residue was diluted with water and extracted with diethyl ether (the ether extracts were discarded). The aqueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl ether and ethyl acetate. The combined organic layers were washed with brine, dried (sodium sulfate) and concentrated under reduced pressure to provide the desired product as a light tan solid foam that was used without further purification (4.11 g, 98%): ¹H NMR consistent with structure; MS (IS) m/e 247 (M + 1).

15

25

10

Preparation 488

To a stirring solution of the product of Preparation 487 (2.50 g, 10.1 mmol), L-proline methyl ester hydrochloride (2.10 g, 12.7 mmol), hydroxybenzotriazole hydrate (1.72 g, 12.7 mmol) and N, N-diisopropylethylamine (4.40 mL, 25.4 mmol) in dichlormethane (90 anhydrous mL), temperature, was added 1-(3-dimethylaminopropyl)-3ethylcarbodiimide (2.63 g, 13.7 mmol). The reaction was stirred for 18 h at room temperature, then was quenched with water and the aqueous layer was extracted with ethyl acetate. The combined extracts were washed with 10% citric acid, saturated sodium

577

bicarbonate, water and brine, dried (sodium sulfate) and evaporated to provide a yellow solid. Flash chromatography (silica gel, 35% ethyl acetate/hexanes - 85% ethyl acetate/hexanes) gave 2.76 g (76%) of the desired product as a white foam solid: ¹H NMR consistent with structure; MS (IS) m/e 358 (M + 1).

Preparation 489

10

15

20

To a suspension of 5% palladium on carbon (1.30 g) in ethyl acetate (90 mL) and ethanol (10 mL), in a Parr reaction bottle, was added the product of Preparation 488 (1.30 g, 3.63 mmol) as a solid. reaction bottle was placed on a Parr shaker, and shaken at room temperature for 45 min under a hydrogen atmosphere (35 psi). The reaction was filtered through a pad of Celite 521. The filtrate was evaporated to an orange solid foam which was then added to a previously prepared mixture of the product of Preparation 1d from Examples Part 2A (1.38 g, 3.63 mmol) and 1-hydroxybenzotriazole hydrate (0.54 g, 4.00 mmol) in dioxane (50 mL) at room temperature. This solution stirred for 15 min at which time 1,3-dicyclohexylcarbodiimide (0.83 g, 4.00 mmol) was added. The reaction was stirred at room temperature for 15 h, then the solvent was evaporated under reduced pressure. The residue was

578

dissolved in ethyl acetate and the 1,3-dicyclohexylurea was filtered away. The filtrate was purified by flash chromatography (silica gel, 75% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to give the desired product as a tan solid foam (0.91 g, 40%): ¹H NMR consistent with structure; MS (IS) m/e 690 (M + 1).

Example 260

10

15

20

25

To a stirring solution of the product of Preparation 489 (0.88 g, 1.28 mmol) and anisole (0.15 mL, 1.28 mmol) in anhydrous dichloromethane (40 mL) at 0°C was added trifluoroacetic acid (6 mL). The reaction was stirred for 4 h, warming to room temperature, and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. The organic layer was collected and the aqueous layer was extracted twice with dichloromethane. The combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. impure foam was purified by flash chromatography (silica gel, ethyl acetate - 5% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide the desired product as an light yellow solid foam (0.710 g, 94%): ¹H NMR consistent with structure; MS (IS) m/e 590 (M

20

579

Preparation 490

To a solution of the product of Preparation 487 5 (2.50 g, 10.0 mmol), in anhydrous dichloromethane (60 mL) at 0°C was added N-methylmorpholine (1.34 mL, 12.0 mmol) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (1.96 g, 11.0 mmol). This mixture stirred for 1 h, warming to room temperature, at which pyrrolidine (0.93 mL, 11.0 mmol) was added. 10 The reaction was then stirred for an additional 2.5 h at room temperature at which time the solvent was evaporated under reduced pressure. The resulting residue was dissolved in ethyl acetate and the remaining solids were filtered away. The filtrate was purified by flash chromatography (silica gel, 40% ethyl acetate/hexanes - ethyl acetate), to provide the desired product as a light yellow foam solid (2.57 g, 86%): ¹H NMR consistent with structure; MS (IS) m/e 300 (M + 1).

Preparation 491

580

To a suspension of 5% palladium on carbon (0.40 g) in ethyl acetate (50 mL) and ethanol (25 mL), in a Parr reaction bottle, was added the product of Preparation 490 (0.65 g, 2.20 mmol) as a solid. reaction bottle was placed on a Parr shaker, and 5 shaken at room temperature for 1 h under a hydrogen atmosphere (35 psi). The reaction was filtered through a pad of Celite 521. The filtrate was evaporated to an orange solid foam which was then 10 added to a previously prepared mixture of the product of Preparation 1d from Examples Part 2A g, 2.12 mmol) and 1-hydroxybenzotriazole hydrate (0.31 g, 2.30 mmol) in N,N-dimethylformamide (50 mL), at room temperature. This solution stirred 15 for 15 min at which time 1,3dicyclohexylcarbodiimide (0.48 g, 2.30 mmol) was The reaction was stirred at room temperature for 15 h, then was quenched with brine, extracted with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate), and 20 evaporated under reduced pressure to give a brown solid foam. The impure solid was purified by flash (silica chromatography gel, 50% acetate/hexanes - ethyl acetate) to give the desired product as a yellow solid foam (0.46 g, 42%): ¹H NMR 25 consistent with structure; MS (IS) m/e 632 (M + 1).

10

15


20

Example 261

To a stirring solution of the product of Preparation 491 (0.37 g, 0.59 mmol) and anisole (0.03 mL, 0.30 mmol) in anhydrous dichloromethane (13 mL) at 0°C was added trifluoroacetic acid (2 mL). The reaction was stirred for 4 h, warming to room temperature, and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. The organic layer was collected and the aqueous layer was extracted twice with dichloromethane. The combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. impure foam was purified by flash chromatography (silica gel, ethyl acetate - 5% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide the desired product as an light yellow solid foam (0.27 g, 86%): ¹H NMR consistent with structure; MS (IS) m/e 532 (M + 1); Anal. Calc'd for $C_{30}H_{37}N_5O_4$: C, 67.68; H, 7.01; N, 13.17. Found: C, 67.30; H, 7.02; N, 12.80.

582

Preparation 492



To a stirring slurry of sodium hydride (0.74 g of a 60% dispersion in mineral oil, 18.6 mmol) in N,Ndimethylformamide (30 mL) at 0°C, was added a solution of 4-nitropyrazole (2.00 g, 17.7 mmol) (from Preparation 450) in N,N-dimethylformamide (20 mL). This solution stirred at 0°C for 10 min then a solution of the product of Preparation 5 of Examples Part 1 (4.75 g, 19.5 mmol) in N,N-dimethylformamide 10 (20 mL) was added. The reaction was stirred at room temperature for 15 h, at which time it was quenched with brine and extracted with ethyl acetate. combined extracts were washed with brine, dried (sodium sulfate) and evaporated to provide a yellow 15 Flash chromatography (silica gel, 20%-50% ethyl acetate/hexanes) yielded the desired product as a colorless oil (4.76 g, 97%): ¹H NMR consistent with structure; MS (IS) m/e 276 (M + 1).

20

Preparation 493

To a stirring solution of the product of Preparation 492 (4.65 g, 16.9 mmol) in dioxane (100 mL) and water (50 mL) at room temperature was added lithium hydroxide (2.83 g, 67.5 mmol). The reaction stirred

583

25 min at room temperature, at which time the dioxane was evaporated under reduced pressure. The residue was diluted with water and extracted with diethyl ether (the ether extracts were discarded). The aqueous layer was acidified (pH 2-3) with 1N HCl and then extracted with diethyl ether and ethyl The combined organic layers were washed acetate. with brine, dried (sodium sulfate) and concentrated under reduced pressure to provide the desired product as a light orange solid that was used without further purification (4.09 g, 98%): ¹H NMR consistent with structure; MS (IS) m/e 248 (M + 1).

Preparation 494

15

20

25

10

To a stirring solution of the product of Preparation 493 (2.00 g, 8.09 mmol), L-proline methyl ester hydrochloride (1.68 g, 10.1 mmol), 1hydroxybenzotriazole hydrate (1.38 g, 10.1 mmol) and N, N-diisopropylethylamine (3.50 mL, 20.2 mmol) in anhydrous dichlormethane (80 mL), at temperature, was added 1-(3-dimethylaminopropyl)-3ethylcarbodiimide (2.09 g, 10.92 mmol). reaction was stirred for 18 h at room temperature, then was quenched with water and the aqueous layer was extracted with ethyl acetate. The combined extracts were washed with 10% citric acid, saturated sodium bicarbonate, water and brine, dried (sodium sulfate) and evaporated to provide a yellow solid.

584

Flash chromatography (silica gel, 35% ethyl acetate/hexanes - 85% ethyl acetate/hexanes) gave the desired product as a white foam solid (1.80 g, 62%): 1 H NMR consistent with structure; MS (IS) m/e 359 (M + 1).

Preparation 495

5

To a suspension of 5% palladium on carbon (0.20 g) 10 in methanol (25 mL), was added the product of Preparation 494 (0.36 g, 1.01 mmol) as a solid. reaction was stirred at room temperature for 4 h under a hydrogen atmosphere (1 atm). The reaction was filtered through a pad of Celite 521. filtrate was evaporated to an off-white solid foam 15 which was then added to a previously prepared mixture of the product of Preparation 1d from Examples Part 2A (0.30 g, 0.80 mmol) and 1hydroxybenzotriazole hydrate (0.12 g, 0.88 mmol) in N, N-dimethylformamide (15 mL) at room temperature. 20 This solution stirred for 15 min at which time 1,3dicyclohexylcarbodiimide (0.18 g, 0.88 mmol) was The reaction was stirred at room temperature for 15 h, then guenched with brine and extracted with ethyl acetate. The combined extracts were 25 washed with brine, dried (sodium sulfate), and evaporated to give a tan solid foam. The crude foam

585

was purified by flash chromatography (silica gel, 75% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to provide the separated diastereomers, as off-white solid foams: 495A (0.18 g, 33%) ¹H NMR consistent with structure; MS (IS) m/e 691 (M + 1); and 495B (0.19 g, 35%): ¹H NMR consistent with structure; MS (IS) m/e 691 (M + 1).

Example 262

10

15

20

25

To a stirring solution of the product of Preparation 0.22 495A (0.15 g,mmol) in anhydrous 0°C dichloromethane (15 mL) at was trifluoroacetic acid (3 mL). The reaction was stirred for 4 h, warming to room temperature, and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. The organic layer was collected and the aqueous layer was extracted twice with dichloromethane. The combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. impure foam was purified by flash chromatography (silica gel, ethyl acetate - 5% methanol/ethyl acetate -5% triethylamine/10% methanol/ethyl acetate) to provide the desired product as a white solid foam (0.10 g, 77%): ¹H NMR consistent with structure; MS (IS) m/e 591 (M + 1).

Example 263

To a stirring solution of the product of Preparation 5 295B (0.15 g, 0.22 mmol) in anhydrous 0°C dichloromethane (15 mL) at added was trifluoroacetic acid (3 mL). The reaction was stirred for 4 h, warming to room temperature, and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. The organic layer was 10 collected and the aqueous layer was extracted twice with dichloromethane. The combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give an light yellow solid foam. 15 impure foam was purified by flash chromatography (silica gel, ethyl acetate - 5% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide the desired product as a white 20 solid foam (0.08 g, 62%): ¹H NMR consistent with structure; MS (IS) m/e 591 (M + 1).

587

Preparation 496

To a stirring solution of the product of Preparation 493 (0.75 g, 3.03 mmol), pyrrolidine (0.32 mL, 3.79 mmol), 1-hydroxybenzotriazole hydrate (0.52 g, 3.79 mmol) and N,N-diisopropylethylamine (1.10 mL, 6.07 mmol) in anhydrous dichlormethane (50 mL), at room temperature, was added 1-(3-dimethylaminopropyl)-3ethylcarbodiimide (0.79 g, 4.10 mmol). The reaction 10 was stirred for 18 h at room temperature, quenched with water and the aqueous layer was extracted with ethyl acetate. The combined extracts were washed with 10% citric acid, saturated sodium bicarbonate, dried (sodium water and brine, sulfate) and 15 evaporated to provide a yellow solid. Flash chromatography (silica gel, 35% ethyl acetate/hexanes - 85% ethyl acetate/hexanes) gave the desired product as a white foam solid (0.83 g, 91%): 1H NMR consistent with structure; MS (IS) m/e 20 301 (M + 1).

WO 99/08697

Preparation 497

To a suspension of 5% palladium on carbon (0.40 g) in a mixture of ethyl acetate (55 mL) and ethanol (7 mL), was added the product of Preparation 496 (0.71 g, 2.35 mmol) as a solid. The reaction was shaken on a Parr shaker at room temperature for 1.5 h under a hydrogen atmosphere (35 psi). The reaction was filtered through a pad of Celite 521. The filtrate was evaporated to an off-white solid foam which was 10 then added to a previously prepared mixture of the product of Preparation 1d from Examples Part 2A (0.89 g, 2.35 mmol) and 1-hydroxybenzotriazole hydrate (0.351 g, 2.58 mmol) in N, N-15 dimethylformamide (35 mL) at room temperature. solution stirred for 15 min at which time 1,3dicyclohexylcarbodiimide (0.53 g, 2.58 mmol) was added. The reaction was stirred at room temperature for 15 h, then guenched with brine and extracted 20 with ethyl acetate. The combined extracts were washed with brine, dried (sodium sulfate), and evaporated to give a tan solid foam. The crude foam was purified by flash chromatography (silica gel, 75% ethyl acetate/hexanes - 10% methanol/ethyl acetate) to provide the desried product as an off-25 white solid foam (0.97 g, 65%): ¹H NMR consistent with structure; MS (IS) m/e 633 (M + 1).

WO 99/08697

10

15

20

589

Example 264

To a stirring solution of the product of Preparation 497 (0.75 g, 1.19 mmol) and anisole (0.04 mL, 0.39 mmol) in anhydrous dichloromethane (30 mL) at 0°C was added trifluoroacetic acid (4 mL). The reaction was stirred for 4 h, warming to room temperature, and then was quenched by pouring over ice-cooled, saturated sodium bicarbonate. The organic layer was collected and the aqueous layer was extracted twice with dichloromethane. The combined organic layers were washed with sodium bicarbonate, water and brine, then dried (sodium sulfate) and evaporated in vacuo to give a light yellow solid foam. The impure foam was purified by flash chromatography (silica gel, ethyl acetate - 5% methanol/ethyl acetate - 5% triethylamine/10% methanol/ethyl acetate) to provide the desired product as a white solid foam (0.62 g, 98%): ¹H NMR consistent with structure; MS (IS) m/e 533 (M + 1).

590

EXAMPLES PART 3

Preparation of 1-N-alkylated-4-nitroimidazoles

5

10

15

Preparation 1

To a solution of α -bromophenylacetic acid (100 g, 466 mmol) stirring in absolute ethanol (500 mL) at room temperature was added p-toluenesulfonic acid monohydrate (10 g, 53 mmol). This solution was heated to reflux and, after 8 h, concentrated to dryness. The resulting residue was dissolved in ethyl acetate, washed with saturated aqueous sodium bicarbonate, brine, dried over sodium sulfate, filtered, and concentrated to yield 77 g (68 %) of the desired product as an orange oil: $^1\text{H-NMR}$ is consistent with structure; MS (FD) 241.9, 243.9.

20

Preparation 2

To a slurry of sodium hydride (13.6 g of a 60% dispersion in mineral oil, 341 mmol) stirring in N,N-dimethylformamide (240 mL) was carefully added

591

4-nitroimidazole (38.6 g, 341 mmol) such that the temperature during the addition was maintained below 40°C. This resulting slurry was stirred for 1 h and then cooled to 5°C. To this mixture was slowly added a compound of Preparation 1 (76 g, 310 mmol) at a rate such that the reaction temperature was maintained below 20°C. After 4 h, the reaction was concentrated and subsequently extracted with ethyl acetate. The combined organic extracts were filtered, washed with water, brine, dried over sodium sulfate, filtered and concentrated. The resulting residue was purified by silica gel chromatography (methanol/chloroform gradient) to yield the 60.1 g (70%) of the desired product as a white solid: ¹H-NMR is consistent with structure; MS (FD) 275 (M+); Anal. Calc'd. for: C, 56.73; H, 4.73; N, 15.27. Found: C, 56.48; H, 4.78; N, 15.08.

10

15

20

25

30

Preparation 3

Mo Mo

A solution of the product of Preparation 2 (10.00 g, 36.36 mmol) in DMF (50 mL) was added dropwise to a suspension of sodium hydride (1.60 g, 40.00 mmol) in DMF (50 mL) under nitrogen at 0 °C. The mixture was stirred 10 min., then methyl iodide (2.5 mL, 40.00 mmol) was added dropwise. The reaction was stirred thirty minutes at 0°C, then 1 h at ambient temperature. The mixture was


592

quenched with a saturated solution of sodium Ethyl acetate was added and the bicarbonate. mixture washed with bicarbonate followed by brine. The organic layer was dried over sodium sulfate, concentrated in vacuo. filtered, and The purified resulting foam was by flash (300 silica, 2:3 chromatography g ethy1 acetate/hexanes) to yield the desired product (8.81 g, 84%) as a light yellow foam: ¹H NMR (30010 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{14}H_{15}N_3O_4$; 58.13 C, 5.23 H, 14.53 N; found 57.88 C, 5.36 H, 14.39 N; FDMS (M+) - 289.

Preparation 4

15

20



To a solution of 4-methoxyphenylacetic acid, 98 g (590 mmol) in 300 mL of absolute ethanol was added 20 g (105 mmol) of TsOH. The reaction mixture was refluxed for 5 h then concentrated to dryness. The resulting oil was chromatographed on silica gel using 20% ethyl acetate/hexanes as eluant to afford 102 g (89%) of the desired product as a colorless oil. 1 H-NMR (d, DMSO) 1.17 (t, J = 8.7 Hz, 3H), 3.56 (s, 2H), 3.73 (s, 3H), 4.05 (q, J = 7.2 Hz, 2H), 6.87 (d, J = 8.7 Hz, 2H), 7.17 (d, 8.7 Hz, 2H); MS (ion spray) 195.3 (M+1); Anal. Calc'd for $C_{11}H_{14}O_{3}$: C, 68.02; H, 7.27. Found: C, 67.95, 7.17.

25

593

Preparation 5

To a solution of the product of Preparation 4, 40 5 g (200 mmol) in 500 mL of carbon tetrachloride was added 37 g (206 mmol) of N-bromosuccinimide and 4 drops of 48% HBr. The reaction mixture was refluxed for 5 h, filtered and concentrated to dryness. The resulting oil was chromatographed on silica gel using chloroform as eluant to afford 10 51.1 g (94%) of the desired product as a colorless oil. $^{1}H-NMR$ (d, DMSO) 1.19 (t, J = 8.4 Hz, 3H), 3.77 (s, 3H), 4.18 (m, 2H), 5.88 (s, 1H), 6.95 (d, J = 8.4 Hz, 2H), 7.50 (d, J = 8.4 Hz, 2H); MS (FD)272, 274 (M+); Anal. Calc'd for $C_{11}H_{13}BrO_3$: C, 15 48.37; H, 4.80. Found: C, 48.52, 4.77.

Preparation 6

20


25

To a solution of the product of Preparation 5, 49.5 g (181 mmol) in 500 mL of DMF was added 20.5 g (181 mmol) of 4-nitroimidazole and 75 g (543 mmol) of potassium carbonate. The reaction mixture was stirred overnight at ambient temperature, filtered and concentrated to dryness.

594

The resulting oil was partitioned between ethyl acetate and water and extracted with ethyl The combined organics were washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. The resulting oil was absorbed onto a silica pad and chromatographed on silica gel using 30-70% ethyl acetates/hexanes as eluant to yield 33.6 g (61%) of the desired product as an orange oil that solidifies upon sitting. $^{1}H-NMR$ (d, DMSO) 1.17 (t, J = 7.2 Hz, 3H), 3.78 (s, 3H), 4.25 (q, J = 7.2 Hz, 2H), 6.57(s, 1H), 7.02 (d, J = 8.7 Hz, 2H), 7.46 (d, J =8.7 Hz, 2H), 7.92 (s, 1H), 8.38 (s, 1H); MS (ion)spray) 306 (M+1); Anal. Calc'd for $C_{14}H_{15}N_3O_5$: C, 15 55.08; H, 4.95; N, 13.76. Found: C, 54.93; H, 4.89; N, 13.82

Preparation 7



20


25

10

Prepared as in Preparation 3 using the product of Preparation 6 (5.00 g, 16.39 mmol) in DMF (25 mL) and sodium hydride (0.72 g, 18.03 mmol) and methyl iodide (1.12 ml, 18.03 mmol) in DMF (25 mL) to yield the desired product (4.81 g, 92%) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C15H17N3O5; 56.42 C, 5.37 H, 13.16 N; found 56.13 C, 5.35 H, 13.01 N; ISMS (M+) -320.

595

Preparation 7A



5 Biphenylacetic acid (25.2 g, 119 mmol), TsOH (3.3 g, 17 mmol), absolute ethanol (250 mL), as in Preparation 4. 25.4 g (89%) of the desired product as a yellow oil. ¹H-NMR is consistent with structure; MS (FD) 240.1 (M+); Anal. Calc'd for C₁₆H₁₆O₂: C, 79.97; H, 6.71. Found: C, 79.75; H, 6.59.

Preparation 8

The product of Preparation 7A (18.0 g, 75.0 mmol), N-bromosuccinimide (13.7 g, 77.25 mL), 48% HBr (4 drops), carbon tetrachloride (80 mL), as in Preparation 5 gave 22.56 g (94%) of the desired product as a yellow oil. ¹H-NMR is consistent with structure; MS (FD) 318, 320 (M+); Anal. Calc'd for C₁₆H₁₅BrO₂ 0.05CHCl₃: C, 60.21; H, 4.74. Found: C, 59.50; H, 4.75.


596

Preparation 9

To a slurry of 2.42 g (60.5 mmol) of sodium hydride in 200 mL of DMF at ambient temperature 5 was added 6.9 g (60.5 mmol) of 4-nitroimidazole. After 10 min, 17.62 g (55.0 mmol) of the product of Preparation 8 was added. The resulting mixture was stirred overnight at ambient temperature then concentrated to dryness. The residue was slurried 10 in ethyl acetate and filtered. The resulting oil was partitioned between ethyl acetate and water and extracted with ethyl acetate. The combined organics were washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. 15 The resulting oil was absorbed onto a silica pad and chromatographed on silica gel using 30-50% ethyl acetate/hexanes as eluant to yield 12.0 g (62%) of the desired product as a yellow viscous oil. ¹H-NMR is consistent with structure; MS (FD) 20 351 (M+).


597

Preparation 10



Prepared as described in Preparation 2 using the product of Preparation 9 (11.03 g, 31.39 mmol) in DMF (50 mL) and sodium hydride (1.25 g, 31.39 mmol) and methyl iodide (1.9 ml, 31.39 mmol) in DMF (50 mL) to yield the desired product (10.25 g, 89%) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₂₀H₁₉N₃O₄; 65.75 C, 5.26 H, 11.50 N; found 63.84 C, 5.16 H, 10.94 N; ISMS (M+) - 366.

Preparation 11



15

20


A suspension of 2-naphthyl acetic acid (49.37 g, 265.0 mmol) and thionyl chloride (80 mL) in carbon tetrachloride (55 mL) was heated to reflux for 20 minutes at which time all material went into solution. The reaction was cooled to ambient temperature. Carbon tetrachloride (125 mL), N-bromosuccinamide (56.60 g, 318.0 mmol), and hydrobromic acid (48% aq., catalytic, 0.5 mL) were added. The mixture was heated to reflux for 30

598

min., cooled to ambient temperature, filtered, and concentrated in vacuo. The material was redissolved in dichloromethane (200 mL) and excess ethanol (100 mL) was added dropwise. The mixture was stirred at ambient temperature 1 hour, then concentrated in vacuo. The crude material was silica, chromatographed (700 q 30% ethyl acetate/hexane) to yield a crude tan solid. This crude material was dissolved dimethylformamide (200 mL) and 4-nitroimidazole (29.78 g, 10 mmol) and potassium carbonate (72.70 g, 526.8 mmol) were added. The reaction was stirred at ambient temperature, then concentrated in vacuo to Ethyl acetate and water were added and 15 the mixture washed with sodium bicarbonate and The organic layer was dried over sodium brine. sulfate and concentrated in vacuo. The crude material was chromatographed (1.0 kg silica, 30% ethyl acetate/hexane) to yield the desired product 20 (40.2 g, 47%) as a brown foam: ${}^{1}H$ NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{17}H_{15}N_3O_4$; 62.76 C, 4.65 H, 12.92 N; found 60.54 C, 4.35 H, 12.04 N; ISMS (M+) - 326.


599

Preparation 12



Prepared as described in Preparation 3 using the product of Preparation 10 (13.9 g, 42.65 mmol) in DMF (50 mL) and sodium hydride (1.71 g, 42.65 mmol) and methyl iodide (2.64 ml, 42.65 mmol) in DMF (50 mL) to yield the desired product (10.94 g, 77%) as a light yellow oil: ¹H NMR (300 MHz, CDCl₃) 10 - consistent with structure; Anal. calcd. for $C_{18}H_{17}N_{3}O_{4}$; 63.71 C, 5.05 H, 12.38 N; found 63.80 C, 4.98 H, 12.41 N; ISMS (M+) - 340.

15



A solution of the product of Preparation 2 (8.35 20 g, 28.89 mmol) in THF (100 mL) was treated with lithium hydroxide (1.82 g, 43.34 mmol) and water The reaction was stirred at ambient temperature for 30 minutes. Water was added and the mixture washed with diethyl ether. The pH of 25 the aqueous layer was adjusted to 3.0 with 10%

.

WO 99/08697

600

PCT/US98/17201

sodium bisulfate. The mixture was saturated with sodium chloride and washed with ethyl acetate. The ethyl acetate washes were combined, dried over sodium sulfate, filtered, and concentrated vacuo. The resulting crude solid was dissolved in 5 anhydrous dichloromethane (100 mL) under nitrogen. To this solution was added catalytic DMF (0.1 mL) and excess oxalyl chloride (25 g). This mixture was stirred 3 hours, then concentrated in vacuo. The resulting crude foam was dissolved in THF (20 10 mL) and added dropwise to a solution of lithium (4R,5S)-(+)-4-methyl-5-phenyl-2-oxazolidinone[generated by adding n-BuLi (1.6M in hexanes, 19.9 mL, 31.82 mmol) dropwise to a solution of (4R, 5S) - (+) -4-methyl-5-phenyl-2-oxazolidinone (5.64 g, 15 in THF (50 mL) at -78 C under 31.82 mmol) nitrogen. This solution was stirred 20 min., then used without further purification.]. The resulting mixture was stirred at -78 C for 30 min, 20 warmed to 0 C. The mixture was quenched with saturated sodium bicarbonate. Ethyl acetate and water were added and the mixture washed with sodium bicarbonate and brine. The organic layer dried over sodium sulfate, filtered, concentrated in vacuo. The resulting foam was 25 purified by flash chromatography (400 g silica, 5% diethvl ether/dichloromethane) to yield 1 (3.76)31% vield) diastereomer g, (4.32 g, 36%) of the desired diastereomer 2 product as colorless foams: diastereomer 1 - 1H NMR 30 (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{22}H_{20}N_4O_5$; 62.85 C, 4.80 H, 13.33 N; found 60.97 C, 4.64 H, 12.44 N; FDMS (M+) - 420: diastereomer 2 - ¹H NMR (300 MHz, CDCl₃)

601

consistent with structure; Anal. calcd. for $C_{22}H_{20}N_4O_5$; 62.85 C, 4.80 H, 13.33 N; found 62.41 C, 4.82 H, 11.92 N; FDMS (M+) - 420.

Preparation 14

5

Prepared as described in Preparation 13 using the product of Preparation 7 (4.80 g, 15.03 mmol) in THF (50 mL) and lithium hydroxide (1.26 g, 30.06 mmol) in water (25 mL) to give the crude acid. This material was immediately reacted with anhydrous dichloromethane (100 mL), catalytic DMF (0.5 mL), and excess oxalyl chloride (12 mL) to . 15 give the crude acid chloride. This crude product was reacted with THF (20 mL), n-BuLi (1.6M in hexanes, 14.1 mL, 22.54 mmol), and (4R, 5S)-(+)-4methyl-5-phenyl-2-oxazolidinone (4.00 g, mmol) in THF (50 mL) to yield diastereomer 1 (2.79 20 g, 41% yield) and diastereomer 2 (2.80 g, 41%) of the desired product as colorless foams: diastereomer 1 - ¹H NMR (300 MHz, CDCl₃) consistent with structure; Anal. calcd. for 25 $C_{23}H_{22}N_4O_6$; 61.33 C, 4.92 H, 12.44 N; found 60.92 C, 4.82 H, 12.03 N; ISMS (M+) - 451: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{23}H_{22}N_4O_6$; 61.33 C, 4.92 H, 12.44 N; found 61.57 C, 4.98 H, 12.47 N; ISMS (M+) - 451.

602

Preparation 15

5

Prepared as in Preparation 13 using the product of Preparation 10 (10.20 g, 27.92 mmol) in THF (100 mL) and lithium hydroxide (2.34 g, 55.84 mmol) in 10 water (50 mL) to give the crude acid. resulting crude solid was dissolved in anhydrous mL) reacted dichloromethane (150 and catalytic DMF (0.5 mL) and excess oxalyl chloride The resulting crude foam was dissolved (23 mL). in THF (50 mL) and reacted with n-BuLi (1.6M in 15 hexanes, 25.1 mL, 40.28 mmol), (4R, 5S)-(+)-4methyl-5-phenyl-2-oxazolidinone (7.14 g, mmol), and THF (150 mL) to yield diastereomer 1 (6.21 g, 45% yield) and diastereomer 2 (6.20 g, 45%) of the desired product as colorless foams: 1H 20 NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{28}H_{24}N_4O_5$; 66.93 C, 4.99 H, 11.56 N; found 65.32 C, 5.06 H, 10.66 N; ISMS (M+) - $497: {}^{1}H NMR (300 MHz, CDCl_{3}) - consistent with$ 25 structure; Anal. calcd. for $C_{28}H_{24}N_4O_5$; 66.93 C, 4.99 H, 11.56 N; found 65.05 C, 4.92 H, 10.61 N; FDMS (M+) - 497.


603

Preparation 16

Prepared as in Preparation 13 using the product of Preparation 12 (10.9 g, 32.27 mmol) in THF (150 mL) and lithium hydroxide (1.63 g, 38.73 mmol) in water (75 mL) to give the crude acid. resulting crude solid was dissolved in anhydrous dichloromethane (150 mL) and reacted with catalytic DMF (0.5 mL) and excess oxalyl chloride 10 (23 mL). The resulting crude foam was dissolved in THF (50 mL) and reacted with n-BuLi (1.6M in hexanes, 30.1 mL, 48.23 mmol), (4R, 5S)-(+)-4methy1-5-pheny1-2-oxazolidinone (8.55 g, 48.23 mmol), and THF (150 mL) to yield diastereomer 1 15 (6.13 g, 41% yield) and diastereomer 2 (4.82 g, 32%) of the desired product as colorless foams: diastereomer 1 - 1H NMR (300 MHz, CDCl₃) consistent with structure; Anal. calcd. for $C_{26}H_{22}N_4O_5$; 66.38 C, 4.71 H, 11.91 N; found 65.24 C, 20 4.72 H, 11.59 N; ISMS (M+) - 471: distereomer 2 -¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{26}H_{22}N_4O_5$; 66.38 C, 4.71 H, 11.91 N; found 66.45 C, 4.77 H, 12.20 N; ISMS (M+) - 471.25

Preparation 17

604



A solution of the product of Preparation 13, diastereomer 1 (2.30 g, 5.48 mmol) in THF (50 mL) was added to a solution of lithium hydroxide (0.25 g, 6.03 mmol) in water (25 mL). The resulting mixture was stirred at ambient temperature for 30 minutes. Water was added and the mixture washed with diethyl ether. The pH of the aqueous layer was adjusted to 3.0 with 10% aqueous sodium bisulfate. The mixture was saturated with sodium 10 chloride and washed with ethyl acetate. The ethyl acetate washes were combined, dried over sodium sulfate, filtered, and concentrated in vacuo. resulting crude solid was dissolved in anhydrous dichloromethane (50 mL) under nitrogen. 15 To this solution was added catalytic DMF (0.1 mL) excess oxalyl chloride (5 g). This mixture was stirred 3 hours, then concentrated in vacuo. resulting crude foam was dissolved in anhydrous dichloromethane (50 mL) and cooled to 0 °C. 4-20 (catalytic, Dimethylaminopyridine 10 mg) and pyrrolidine (1.8 mL, 18.74 mmol) were added and the resulting solution stirred for 18 hours. Dichloromethane was then added and the mixture washed with sodium bicarbonate and brine. 25 organic layer was dried over sodium sulfate, filtered, and concentrated in vacuo. The crude foam was purified by flash chromatography (silica, 100 g, 5% methanol/dichloromethane) to yield the 30 desired product (1.73 g, 88% yield) as a colorless


605

foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{16}H_{18}N_4O_3$; 61.14 C, 5.77 H, 17.82 N; found 60.67 C, 5.78 H, 16.03 N; FDMS (M+) - 314.

5

10

Preparation 18




Prepared as in Preparation 17 using the product of Preparation 13, diastereomer 1 (1.88 g, 5.44 mmol) in THF (50 mL) and lithium hydroxide (0.23 g, 5.63 mmol) in water (25 mL) to give the crude acid. The resulting crude solid was dissolved in anhydrous dichloromethane (50 mL) and reacted with catalytic DMF (0.1 mL) and excess oxalyl chloride 15 (5 g) to give the crude acid chloride. resulting crude foam was dissolved in anhydrous and reacted with 4dichloromethane (50 mL) Dimethylaminopyridine (catalytic, 10 mg), proline methyl ester hydrochloride (0.90 g, 5.44 20 mmol), and N, N-diisopropylethylamine (2.8 16.31 mmol) to yield the desired product (1.39 g, 69% yield) as a colorless foam: 1H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{18}H_{20}N_4O_5$; 58.60 C, 5.41 H, 15.04 N; found 57.95

25 C, 5.40 H, 13.45 N; FDMS (M+) - 372.

606


Preparation 19



Prepared as in Preparation 17 using the product of Preparation 13, diastereomer 1 (1.88 g, 5.44 mmol) in THF (50 mL) and lithium hydroxide (0.23 g, 5.63 mmol) in water (25 mL) to give the crude acid. The resulting crude solid was dissolved in anhydrous dichloromethane (50 mL) and reacted with catalytic DMF (0.1 mL) and excess oxalyl chloride 10 (5 g) to give the crude acid chloride. The resulting crude foam was dissolved in anhydrous dichloromethane (50 mL) and reacted with 4-Dimethylaminopyridine (catalytic, 10 mg), proline methyl ester hydrochloride (0.90 g, 5.44 15 mmol), and N, N-diethylisopropylamine (2.8 mL, 16.31 mmol) to yield the desired product (1.21 g, 65% yield) as a colorless foam: 1H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{18}H_{22}N_4O_3$; 63.14 C, 6.48 H, 16.36 N; found 63.29 20 C, 6.45 H, 15.29 N; FDMS (M+) - 342.

607

Preparation 20



Prepared as in Preparation 17 using the product of Preparation 14, diastereomer 1 (1.25 g, 2.78 mmol) in THF (50 mL) and lithium hydroxide (0.14 g, 3.33 mmol) in water (25 mL) to give the crude acid. The resulting crude solid was dissolved anhydrous dichloromethane (50 mL) and reacted with catalytic DMF (0.1 mL) and excess oxalyl chloride 10 (5 g) to give the crude acid chloride. resulting crude foam was dissolved in anhydrous dichloromethane (50 mL) and reacted with 4-Dimethylaminopyridine (catalytic, 10 mg) pyrrolidine (0.24 mL, 2.89 mmol) to yield the 15 desired product (0.78 g, 86% yield) as a colorless foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{17}H_{20}N_4O_4$; 59.59 C, 5.85 H, 16.27 N; found 59.59 C, 5.96 H, 16.19 N; ISMS (M+) - 345.20

Preparation 21

Prepared as in Preparation 17 using the product of Preparation 13, diastereomer 1 (2.31 g, 5.15 mmol)

608


in THF (50 mL) and lithium hydroxide (0.26 g, 6.18 mmol) in water (25 mL) to give the crude acid. The resulting crude solid was dissolved anhydrous dichloromethane (50 mL) and reacted with catalytic DMF (0.1 mL) and excess oxalyl chloride (5 g) to give the crude acid chloride. resulting crude foam was dissolved in anhydrous dichloromethane (50 mL) and reacted with 4-Dimethylaminopyridine (catalytic, 10 ma) dimethylamine (2.0 M in THF, 7.7 mL, 15.46 mmol) to yield the desired product (1.57 g, 96% yield) as a colorless foam: ¹H NMR (300 MHz, CDCl₃) consistent with structure; Anal. calcd. $C_{15}H_{18}N_4O_4$; 56.60 C, 5.70 H, 17.60 N; found 57.04 C, 6.09 H, 16.82 N; ISMS (M+) - 319.

5

10

15

Preparation 22




Prepared as in Preparation 17 using the product of 20 Preparation 14, diastereomer 1 (1.00g, 2.22 mmol) in THF (50 mL) and lithium hydroxide (0.10 g, 2.44 mmol) in water (25 mL) to give the crude acid. resulting crude solid was dissolved anhydrous dichloromethane (50 mL) and reacted with 25 catalytic DMF (0.1 mL) and excess oxalyl chloride (5 g) to give the crude acid chloride. resulting crude foam was dissolved in anhydrous dichloromethane (50 mL) and reacted with Dimethylaminopyridine (catalytic, 10 mg) and 4-30

609

methylpiperidine (0.34 mL, 2.71 mmol) to yield the desired product (0.38 g, 50% yield) as a colorless foam: ^{1}H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₁₉H₂₄N₄O₄; 61.28 C, 6.50 H, 15.05 N; found 61.38 C, 6.40 H, 15.11 N; FDMS (M+) - 372.

Preparation 23




10

Prepared as in Preparation 17 using the product of Preparation 15, diastereomer 1 (1.00 g, 2.02 mmol) in THF (20 mL) and lithium hydroxide (0.13 g, 3.09 mmol) in water (10 mL) to give the crude acid. resulting crude solid was dissolved 15 anhydrous dichloromethane (20 mL) and reacted with catalytic DMF (0.1 mL) and excess oxalyl chloride (5 g). The resulting crude foam was dissolved in anhydrous dichloromethane (20 mL) and reacted with 20 4-Dimethylaminopyridine (catalytic, 10 ma) and pyrrolidine (0.65 mL, 7.76 mmol) to yield the desired product (0.80 g, 98% yield) as a colorless foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{22}H_{22}N_4O_3$; 67.68 C, 5.68 H, 14.34 N; found 65.36 C, 5.54 H, 13.43 N; ISMS 25 (M+) - 391.


610

Preparation 24



Prepared as in Preparation 17 using the product of Preparation 15, diastereomer 1 (0.50 g, 1.00 mmol) in THF (20 mL) and lithium hydroxide (0.05 g, 1.10 mmol) in water (10 mL) to give the crude acid. The resulting crude solid was dissolved in anhydrous dichloromethane (50 mL) and reacted with catalytic DMF (0.1 mL) and excess oxalyl chloride (5 g) to give the crude acid chloride. 10 resulting crude foam was dissolved in anhydrous dichloromethane (50 mL) and reacted with 4-Dimethylaminopyridine (catalytic, 10 mg), Nmethylmorpholine (0.33 mL, 3.00 mmol), dimethylamine hydrochloride (0.13 g, 1.50 mmol) to 15 yield the desired product (0.30 g, 82% yield) as a colorless foam: ¹H NMR (300 MHz, CDCl₃) consistent with structure; Anal. calcd. for $C_{20}H_{20}N_4O_3$; 65.92 C, 5.53 H, 15.37 N; found 64.17 C, 20 5.41 H, 14.15 N; ISMS (M+) - 365.

Preparation 25




Prepared as in Preparation 17 using the product of Preparation 15, diastereomer 1 (0.40 g, 0.80 mmol) in THF (20 mL) and lithium hydroxide (0.04 g, 0.96 mmol) in water (10 mL) to give the crude acid. The resulting crude solid was dissolved in anhydrous dichloromethane (50 mL) and reacted with catalytic DMF (0.1 mL) and excess oxalyl chloride (5 g) to give the crude acid chloride. resulting crude foam was dissolved in anhydrous dichloromethane (50 mL) and reacted with 4-Dimethylaminopyridine (catalytic, 10 mg) and 4methylpiperidine (0.24 mL, 2.89 mmol) to yield the desired product (0.30 g, 90% yield) as a colorless foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{24}H_{26}N_4O_3$; 68.88 C, 6.26 H, 13.39 N; found 67.40 C, 6.72 H, 12.45 N; FDMS (M+) - 419.

20

10

15

Preparation 26



Prepared as in Preparation 17 using the product of Preparation 16, diastereomer 1 (1.00 g, 2.13 mmol) in THF (20 mL) and lithium hydroxide (0.10 g, 2.33 mmol) in water (10 mL) to give the crude acid.

612

The resulting crude solid was dissolved anhydrous dichloromethane (20 mL) and reacted with catalytic DMF (0.1 mL) and excess oxalyl chloride The resulting crude foam was dissolved in anhydrous dichloromethane (20 mL) and reacted with 4-Dimethylaminopyridine (catalytic, 10 mg) pyrrolidine (0.61 mL, 6.39 mmol)to yield the desired product (0.42 g, 54% yield) as a colorless foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{20}H_{20}N_4O_3$; 65.92 C, 5.53 H, 15.38 N; found 61.50 C, 5.41 H, 13.91 N; ISMS (M+) - 365.

Preparation of Dipeptide Acids

15

20

25

30

10

Preparation 27

To a solution of boc-(OBz)-D-Ser-OH (25.0 g, 84.7 mmol) stirring in anhydrous N, N-dimethylformamide (500 mL) at room temperature was added sodium bicarbonate (14.2 g, 169 mmol) followed by methyl iodide (26.4 mL, 424 mmol). After 18 h, reaction mixture was concentrated to approximately 100 mL. Ethyl acetate was added and the mixture washed with aqueous sodium bicarbonate and brine. The organic extract was dried and concentrated to give the desired compound (25 g, 96%) as a light yellow oil: 1 H NMR (300 MHz, CDCl₃) d 1.45 (s, 9H), 3.70 (m, 1H), 3.75 (s, 3H), 3.85 (m, 1H), 4.50 (m, 3H), 7.30 (m, 5H); MS (FD) m/e 310; Anal. calc'd for C₁₆H₂₃NO₅: C, 62.12; H, 7.49; N, 4.53. Found: C, 62.31; H, 7.49; N, 4.43.

Preparation 28

To a solution of the product of Preparation 27 (5.0 g, 16 mmol) stirring in dichloromethane (40 and anisole (1 mL) at 0 °C was added trifluoroacetic acid (10 mL). After 4 h at room temperature, saturated sodium bicarbonate solution was added and the mixture extracted with ethyl The combined organic extracts were 10 acetate. washed with brine, dried over sodium sufate, and concentrated. The crude product was used in the next step without further purification.

15 Preparation 29

20

To a solution of the product of Preparation 28 (65.4 mmol), boc-a-aminoisobutyric acid (13.2 g, 65.4 mmol), 1-hydroxybenzotriazole (8.8 g, 65.4 mmol), and N, N-diisopropylethylamine (22.8 mL, 130.7 mmol) stirring in dichloromethane (500 mL) at 0 °C was added 1-(3-dimethylaminopropyl)-3ethylcarbodiimide (12.3 g, 71.9 mmol). After 18 h, ethyl acetate and saturated ammonium chloride were added and the mixture extracted with ammonium chloride, sodium bicarbonate, and brine. organic extracts were dried over sodium sulfate and concentrated. Purification by silica gel chromatography (25% ethyl acetate/hexanes) yielded 30 the desired compound (21.6 g, 83%) as a white

614

solid: ¹H NMR (300 MHz, CDCl₃) d 1.39 (s, 9H), 1.48 (s, 6H), 3.62 (dd, J = 3.4, 9.1 Hz, 1H), 3.70 (s, 3H), 3.85 (dd, J = 3.4, 9.1 Hz, 1H), 4.48 (dd, J = 12.5, 22.7 Hz, 2H), 4.75 (m, 1H), 4.92 (s, 1H), 7.11 (d, J = 8.6 Hz, 1H), 7.35 (m, 5H); MS (FD) m/e 395; Anal. calc'd for C₂0H₃0N₂O₆: C, 60.90; H, 7.67; N, 7.10. Found: C, 61.02; H, 7.78; N, 7.10.

10 Preparation 30

$$\bigcirc \bigcap_{HO_2C} \stackrel{H}{\overset{}{\overset{}{\bigvee}}}_{O}_{NHBoc}$$

To a solution of the product of Preparation 29 (5.30 g, 13.4) stirring in dioxane (100 mL)/water (50 mL) at room temperature was added lithium hydroxide (2.80 g, 67.3 mmol). After 18 h, water 15 was added and the solution concentrated. The resulting mixture was extracted with diethyl ether. Sodium chloride was added to the aqueous layer and the pH adjusted to 3.5 with 1 N HCl. The resulting mixture was extracted with ethyl 20 acetate and the combined organic extracts dried over sodium sulfate then concentrated to yield the title compound (4.40 g, 86%) as a white foam: 1 H NMR (300 MHz, CDCl₃) d 1.39 (s, 9H), 1.45 (s, 3H), 1.47 (s, 3H), 3.68 (m, 1H), 3.95 (m, 1H), 4.54 (s, 25 2H), 4.70 (m, 1H), 5.51 (bs, 1H), 7.18 (d, J = 9.1Hz, 1H), 7.25 (m, 5H), 9.90 (bs, 1H); MS (FD) m/e 381; Anal. calc'd for C19H28N2O6: C, 59.99; H, 7.42; N, 7.36. Found: C, 59.74; H, 7.26; N, 7.30.

615

Preparation 31

A solution of sodium ethoxide was generated by the addition of sodium metal (52.89 grams, 2.3007 mol) over 3 hours to ethanol (1500 mL). To the sodium ethoxide solution at ambient temperature was added a solution of diethylacetamidomalonate (499.75 grams, 2.3007 mol) dissolved in ethanol (225 mL). The reaction mixture was stirred for 1.5 hours at 10 ambient temperature. 1-bromo-3-phenylpropane (458.07 grams, 2.3007 mol) was added over 15 minutes and the reaction mixture was refluxed until complete as determined by hplc (16 hours). The reaction mixture was concentrated to dryness 15 and the residue partitioned between ethyl acetate $(1 \times 1500 \text{ mL} \text{ and } 2 \times 500 \text{ mL})$ and water (1500 mL). The ethyl acetate layers were combined, washed with saturated sodium chloride solution (4 x 500 mL), dried using sodium sulfate, and concentrated 20 to give 752.1 grams (98%) of the desired product as a light yellow solid. A 1.0 gram sample was recrystallized from hexane:ethyl acetate (19:1, v:v) to give a mp 84-86°C. ¹H nmr (CDCl₃): δ 1.18-1.23 (t, 6H), 1.37-1.50 (m, 2H), 2.02 (s, 25 3H), 2.34-2.41 (m, 2H), 2.58-2.62 (t, 2H), 4.16-4.24 (q, 4H), 6.76 (s, broad, 1H), 7.11-7.28 (m, 5H). ¹³C nmr (CDCl₃): δ 13.95, 23.03, 25.67, 35.45, 62.46, 66.49, 125.40, 128.27, 128.35, 141.77, 168.11, 168.94. MS (FIA) 30 m/z 336.3 ([M+H]⁺). IR (KBr, cm⁻¹) 1645.98

(amide), 1744.76 (C=O). Anal. Calcd. for $C_{18}H_{25}NO_5$:

616

C, 64.46; H, 7.51; N, 4.17. Found: C, 64.60; H, 7.37; N, 4.39.

Preparation 32

5

slurry consisting of the product Preparation 31 (249.15 grams, 0.7428 mol) and 2.5 N sodium hydroxide solution was heated at 100°C for 10 three hours. The reaction mixture was cooled to 30°C and the pH adjusted to 5.0 using concentrated hydrochloric acid. The solution was heated to 100 °C and the pH was held at 5.0 using concentrated hydrochloric acid as needed until the reaction was 15 complete as determined by hplc. The solution was filtered while hot through diatomaceous earth. The filtrate was cooled to 5-10 °C and the pH adjusted to 1.0 using concentrated hydrochloric acid. The resulting slurry was stirred for 1 hour at 5 °C, filtered, and dried in vacuum at 50 °C to 20 give 160.34 grams (92%) of (DL)-N-acetyl-2-amino-5-phenylpentanoic acid as a white powder, mp 145-148 °C. ¹H nmr (DMSO-d₆): δ 1.60-1.71 (m, 4H), 1.86 (s, 3H), 2.56-2.59 (m, 2H), 4.19-4.23 (m, 1H), 7.16-7.30 (m, 5H), 8.14 (d, 1H). ¹³C nmr (25 DMSO- d_6): δ 23.17, 28.25, 31.55, 35.51, 52.55, 126.60, 129.14, 142.64, 170.25, 174.65. MS (FIA) m/z 236.2 (M⁺). IR (KBr, cm⁻¹) 1609.17 (amide), 1741.12 (C=O). Anal. Calcd. for $C_{13}H_{17}NO_3$: C, 30 66.36; H, 7.28; N, 5.95. Found: C, 66.41; H, 7.15; N, 5.96.

617

Preparation 33

A solution consisting of (DL)-N-acetyl-2-amino-5phenylpentanoic acid (438.0 grams, 1.862 mol), chloride (1.10 grams), 2N potassium hydroxide solution (931 mL, 1.862 mol), and water (8000 mL) was adjusted to a pH of 8.0 by the addition of 2N potassium hydroxide solution. То added the reaction mixture was Acylase I (Aspergillus melleus, 39.42 grams) and vigorously 10 stirred for 24 hours at 40 °C while maintaining a pH of 8.0 by addition of 2N potassium hydroxide. The resulting slurry was filtered. The filtrate was adjusted to a pH of 2.0 giving a thick slurry. The product was isolated by filtration, washed 15 with hexane (2000 mL) and dried in vacuum at 50 °C to give 188.52 grams (43%) of (D)-N-acetyl-2amino-5-phenylpentanoic acid. ^{1}H nmr (DMSO- d_{6}): δ 1.59-1.74 (m, 4H), 1.86 (s, 3H), 2.57-2.60 (m, 20 · 2H), 4.22-4.26 (m, 1H), 7.16-7.30 (m, 5H), 8.02 (d, 1H), 12.39 (s, broad, 1H). ¹³C nmr (DMSO-d₆): 23.18, 28.13, 31.66, 35.54, 52.58, 126.56, 129.10, 142.67, 170.12, 174.48. MS (FIA) m/z 236.1 (M^{+}) . IR (KBr, cm^{-1}) 1625.08 (amide), Anal. Calcd. for $C_{13}H_{17}NO_3$: $C_{13}H_{17}NO_3$: 1700.24 (C=O). 25 66.36; H, 7.28; N, 5.95. Found: C, 66.49; H,

7.00; N, 6.03.

618

Preparation 34

A solution consisting of (D)-N-acetyl-2-amino-5phenylpentanoic acid (188.8 grams, 0.8024 mol), ethanol (535 mL), and concentrated hydrochloric 5 acid (268 mL, 3.21 mol) was warmed to 85 °C and monitored by hplc. The reaction was determined to be incomplete by hplc at 14.5 hours and additional concentrated hydrochloric acid (50 mL) 10 was added. The reaction was determined to be complete by hplc after 22.5 hours. Water was azeotropically distilled from the reaction by continuous addition and distillation of 8000 mL of ethanol. The ethanol was azeotropically distilled from the reaction by the continuous addition and 15 distillation of ethyl acetate (2000 mL). °C cooling the solution to 0 the product crystallized. The solution containing the product was stirred for 1 hour at 0 °C, filtered, and the cake dried in vacuum at 40 °C to give 199.0 grams 20 (96%) of 2-amino-5-phenylpentanoic acid, ethyl ester hydrochloride, mp 117-121 °C. ¹H nmr (DMSOd6): δ 1.15-1.21 (t, 3H), 1.50-1.89 (m, 4H), 2.48-2.67 (m, 2H), 3.92-3.98 (t, 1H), 4.08-4.25 25 (m, 2H), 7.12-7.29 (m, 5H), 8.76 (s, broad, 3H).¹³C nmr (DMSO-d₆): δ 13.90, 25.97, 29.52, 34.41, 61.56, 124.91, 125.81, 128.24, 141.27, 51.71. 169.35. MS (FIA) m/z 222.3 (M⁺). IR (KBr, cm⁻¹) 1741.14 (C=O). $[\alpha]^{20}_{p} = -11.17$ (c = 30.62 mg / 3mL, MeOH). Anal. Calcd. for $C_{13}H_{20}NO_2Cl$: C, 30

619

60.58; H, 7.82; N, 5.43. Found: C, 60.45; H, 7.67; N, 5.55.

Preparation 35

5

A slurry consisting of N-t-BOC- α -aminoisobutyric acid (90.64 grams, 0.446 mol), 2-chloro-4,6dimethoxy-1,3,5- triazine (75.90 grams, 0.425 mol), N-methyl morpholine (88.13 grams, 0.871 mol), and diethyl ether (1000 mL) was stirred at 10 ambient temperature until complete as determined by hplc (3 hours). The 2-amino-5-phenylpentanoic acid, ethyl ester hydrochloride (109.55 grams, 0.425 mol) was added and the reaction mixture stirred for 16 hours at ambient temperature. The 15 reaction mixture was partitioned between 10% citric acid solution (1000 mL) and ethyl acetate (3 x 500 mL). The organic phase was washed with 10% citric acid solution (3 x 500 mL), saturated sodium bicarbonate solution (3 x 500 mL), water (1 20 x 500 mL), dried using sodium sulfate, and concentrated to dryness. The residue was recrystallized from hexane (3000 mL) to give 155.11 grams of compound 2, mp 97-99 °C. ¹H nmr (25 CDCl₃): δ 1.25-1.28 (t, 3H), 1.43 (s, 9H), 1.48 (s, 3H), 1.50 (s, 3H), 1.70-1.73 (m, 3H), 1.87-1.93 (m, 1H), 2.62-2.67 (m, 2H), 4.16-4.21 (m, 2H), 4.57-4.62 (m, 1H), 4.95 (s, 1H), 6.96 (s, broad, 1H), 7.16-7.19 (m, 3H), 7.26-7.33 (m, 2H). ¹³C nmr (CDCl₃): δ 14.53, 26.32, 27.17, 28.67, 30 32.47, 35.73, 52.54, 57.17, 61.62, 126.21, 128.69,

620

128.79, 142.12, 154.99, 172.81, 174.69. MS (FIA) m/z 407.5 ([M+H] $^+$). IR (KBr, cm $^{-1}$) 1652.75, 1685.52 (amides), 1741.73 (C=O). [α] $^{20}_D$ = 7.83 (c = 10.22 mg / 1mL, MeOH). UV (0.1% trifluoroacetic acid in water : acetonitrile) λ_{max} 215.6 nm. Anal. Calcd. for $C_{22}H_{34}N_2O_5$: C, 65.00; H, 8.43; N, 6.89. Found: C, 65.23; H, 8.34; N, 6.94.

10

15

20

25

30

Preparation 36

solution consisting of the product οf Preparation 35 (152.53 grams, 0.3752 mol) and tetrahydrofuran (884 mL) was cooled to 5 °C. solution consisting of lithium hydroxide (26.96 grams, 1.126 mol) and water (1419 mL) was added to the reaction dropwise over 10 minutes maintaining a temperature of 5-10 °C. Ethanol (183 mL) was added and the reaction stirred at 5-10 °C until complete as determined by hplc (2 hours). The pH of the reaction mixture was adjusted to 2.0 using 6 N hydrochloric acid solution while maintaining The product was extracted from solution 5-10 °C. with ethyl acetate $(3 \times 500 \text{ mL})$. The ethvl acetate extracts were combined, dried using sodium sulfate, and concentrated to dryness to give 141.51 grams (100%) of 427623. 1 H nmr (DMSO-d₆): δ 1.32-1.37 (m, 15H), 1.57-1.75 (m, 4H), 2.51-2.58 (m, 2H), 4.23-4.27 (m, 1H), 6.85 (s, broad, 1H),7.15-7.28 (m, 5H), 7.42 (d, 1H), 12.5 (s, broad, ¹³C nmr (DMSO-d₆): δ 26.31, 27.85, 1H).

621

29.00, 31.86, 35.60, 52.53, 56.60, 78.95, 126.52, 129.05, 129.10, 142.69, 155.06, 174.40, 175.17. MS (FIA) m/z 379.5 ([M+H]⁺). IR (KBr, cm⁻¹) 1641.98, 1692.22 (amides), 1719.72 (C=O). $[\alpha]^{20}_D = -5.73$ (c = 10.48 mg / 1mL, MeOH). Anal. Calcd. for $C_{20}H_{30}N_2O_5$: C, 63.47; H, 7.99; N, 7.40. Found: C, 63.25; H, 7.84; N, 7.46.

Preparation 37

10

N-Methyl morpholine (4.79 mL, 2 eg, 47.3 mm) was stirred slurry of N-Boc-aa added to aminoisobutyric acid (4.43 g, 21.7 mm, 1 eq) and $3.89 ext{ g}$ (21.7 mm, 1.0 eq) of 2-chloro-(4,6)-15 dimethoxy-1,3,5-triazine (CDMT) in 100 diethyl ether. After stirring the reaction mixture at ambient temperature for 1.5 hours, Dtryptophan ester hydrochloride was added. 20 stirring overnight, the reaction mixture was quenched by the addition of 150 mL of 10% aqueous citric acid solution. The layers were separated and the ether layer was washed with 50 mL of saturated sodium bicarbonate solution and 50 mL of water. Lithium hydroxide (2.43 g, 5 eq) was 25 dissolved in 100 ml of water and the solution was added to the diethyl ether solution and stirred vigorously for 4 hours at room temperature. The layers were separated and the pH of the 30 layers was adjusted to 5.6 with 1M HCl. was then adjusted to 3.95 with 10% citric acid

622

solution and the aqueous layer was extracted with 100 mL of ethyl acetate. The ethyl acetate layers were washed with brine, dried over magnesium sulfate and filtered. The volatiles were removed under vacuum to give 82 % yield of the desired product as a white foam. 1H-NMR consistent with structure.

Coupling of Dipeptide Acids to Nitroimidazoles Preparation 38

10

20

25

A solution of the product of Preparation 17 (1.66 g, 5.29 mmol) in THF (5 mL) was added to a 15 suspension of 5% palladium on carbon (0.80 g, catalytic, 25 mL THF) under inert atmosphere. resulting mixture was placed under hydrogen (40 psi) on a Parr shaker for 1.5 hours. The resulting mixture was placed under nitrogen and celite added. The mixture was then filtered and rinsed with THF. The filtrate was place under nitrogen and HOBT (0.71 g, 5.29 mmol), the product of Preparation 30 (2.01 g, 5.29 mmol), EDC (1.00 g, 5.81 mmol), and DIEA (1.0 mL, 5.81 mmol) were added. The resulting mixture was stirred 18 hours ambient temperature, then concentrated in The crude material was dissolved in ethyl vacuo. acetate and washed with sodium bicarbonate and

623

brine. The organic layer was dried over sodium sulfate, filtered, and concentrated in vacuo. resulting crude foam was purified by flash 100 chromatography (silica, g, 2% methanol/dichloromethane) to yield the desired product (0.66 g, 19% yield) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{35}H_{46}N_6O_6$; 65.00 C, 7.17 H, 12.99 N; found 63.21 C, 6.92 H, 12.54 N; FDMS (M+) - 646.

Preparation 39

10

Prepared as in Preparation 38 using the product of 15 Preparation 18 (1.39 g, 3.74 mmol) and 5% palladium on carbon (0.70 g, catalytic, 25 mL THF) to give the crude amine. The resulting filtrate was reacted with HOBT (0.50 g, 3.74 mmol), the product of Preparation 30 (1.42 g, 3.74 mmol), 20 diisopropylethylamine (0.65 mL, 3.74 mmol), and EDCI (0.71 g, 4.11 mmol) to yield the desired product (0.86 g, 33% yield) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{37}H_{48}O_8N_6$; 63.05 C, 6.86 25 H, 11.92 N; found 63.01 C, 6.64 H, 11.85 N; FDMS (M+) - 705.

WO 99/08697


624

Preparation 40

Prepared as in Preparation 38 using the product of Preparation 19 (1.21 g, 3.53 mmol) and 5% palladium on carbon (0.80 g, catalytic, 25 mL THF) to give the crude amine. The resulting filtrate was reacted with HOBT (0.48 g, 3.53 mmol), the product of Preparation 30 (1.34 g, 3.53 mmol), diisopropylethylamine (0.6 mL, 3.53 mmol), and EDCI (0.67 g, 3.88 mmol) to yield the desired product (0.97 g, 41% yield) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₃₇H₅₀N₆O₆; 65.85 C, 7.47 H, 12.45 N; found 64.96 C, 7.48 H, 12.04; FDMS (M+) - 675.

625

Preparation 41



Prepared as in Preparation 38 using the product of Preparation 20 (0.77 g, 2.24 mmol) and 5% palladium on carbon (0.80 g, catalytic, 25 mL THF) to give the crude amine. The resulting filtrate was reacted with HOBT (0.30 g, 2.46 mmol), the product of Preparation 36 (0.85 g, 2.24 mmol), and DCC (0.51 g, 2.46 mmol) to yield the desired product (0.70 g, 46% yield) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₃₇H₅₀N₆O₆; 65.85 C, 7.47 H, 12.45 N; found 65.83 C, 7.27 H, 12.38 N; ISMS (M+) - 675.

10

626

Pr paration 42

Prepared as in Preparation 38 using the product of Preparation 22 (0.92 g, 2.47 mmol) and 5% palladium on carbon (1.00 g, catalytic, 30 mL THF) to give the crude amine. The resulting filtrate was reacted with HOBT (0.35 g, 2.47 mmol), the product of Preparation 36 (0.94 g, 2.47 mmol), and DCC (0.56 g, 2.72 mmol) to yield the desired product (0.92 g, 53% yield) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₃₉H₅₄N₆O₆; 66.64 C, 7.74 H, 11.96 N; found 66.65 C, 7.65 H, 12.02 N; ISMS (M+) - 702.

15

Preparation 43

Prepared as in Preparation 38 using the product of Preparation 22 (1.32 g, 3.55 mmol) and 5% palladium on carbon (1.4 g, catalytic, 50 mL THF) to give the crude amine. The resulting filtrate was reacted with HOBT (0.48 g, 3.55 mmol), the product of Preparation 30 (1.35 g, 3.55 mmol), and DCC (0.81 g, 3.91 mmol) to yield the desired product (0.82 g, 33% yield) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₃₈H₅₂N₆O₇; 64.75 C, 7.44 H, 11.92 N; found 66.19 C, 7.17 H, 12.10 N; ISMS (M+) - 705.

Preparation 44

628

Prepared as in Preparation 38 using the product of 21 (0.27 q, 0.85 mmol) Preparation and 5% palladium on carbon (0.30 g, catalytic, 20 mL THF) to give the crude amine. The resulting filtrate was reacted with HOBT (0.11 g, 0.85 mmol), the product of Preparation 36 (0.32 g, 0.85 mmol), Nmethylmorpholine (0.10 mL, 0.85 mmol), and EDCI (0.16 g, 0.93 mmol) to yield the desired product (0.70 g, 46% yield) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{35}H_{48}N_6O_6$; 66.43 C, 7.65 H, 13.28 N; found 63.53 C, 6.83 H, 12.38 N; ISMS (M+) - 649.

Preparation 45

15

20

25

5

10

Prepared as in Preparation 38 using the product of Preparation 21 (0.75 g, 2.36 mmol) and 5% palladium on carbon (0.80 g, catalytic, 25 mL THF) to give the crude amine. The resulting filtrate was reacted with HOBT (0.32 g, 2.36 mmol), the product of Preparation 30 (0.90 g, 2.36 mmol), and DCC (0.54 g, 2.60 mmol) to yield the desired product (0.86 g, 56% yield) as a light yellow foam: 1 H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{37}H_{50}N_6O_6$; 62.75 C, 7.12 H, 12.91 N; found 62.65 C, 6.95 H, 12.76 N; ISMS (M+) - 651.

Preparation 46

Prepared as in Preparation 38 using the product of Preparation 21 (0.80 g, 2.52 mmol) and 5% palladium on carbon (0.80 g, catalytic, 25 mL THF) to give the crude amine. The resulting filtrate was reacted with HOBT (0.34 g, 2.52 mmol), the product of Preparation 37 (0.99 g, 2.52 mmol), and DCC (0.57 g, 2.77 mmol) to yield the desired product (0.77 g, 46% yield) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₃₇H₅₀N₆O₆; 63.72 C, 6.87 H, 14.86 N; found 63.45 C, 6.86 H, 14.76 N; ISMS (M+) - 660.

Preparation 47

10


Prepared as in Preparation 38 using the product of Preparation 23 (0.80 g, 2.05 mmol) and 5% palladium on carbon (0.80 g, catalytic, 25 mL THF) to yield the crude amine. The filtrate was reacted with HOBT (0.28 g, 2.05 mmol), the product of Preparation 30 (0.78 g, 2.05 mmol), and DCC (0.46 g, 2.05 mmol) to yield the desired product (0.76 g, 51% yield) as a light yellow foam: 1 H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{41}H_{50}N_6O_6$; 68.12 C, 6.97 H, 11.63 N; found 66.93 C, 6.74 H, 11.24 N; ISMS (M+) - 723.

Preparation 48

Prepared as in Preparation 38 using the product of 15 Preparation 23 (0.60 q, 1.54 mmol) palladium on carbon (0.60 g, catalytic, 25 mL THF) to give the crude amine. The resulting filtrate was reacted with HOBT (0.21 g, 1.54 mmol), the product of Preparation 36 (0.58 g, 1.54 mmol), and 20 DCC (0.35 g, 1.69 mmol) to yield the desired product (0.56 g, 50% yield) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{42}H_{52}N_6O_5$; 69.98 C, 7.27 25 H, 11.66 N; found 68.71 C, 6.92 H, 11.39 N; ISMS (M+) - 721.

631


Preparation 49



Prepared as in Preparation 38 using the product of Preparation 23 (0.20 g, 0.51 mmol) and 5% 5 palladium on carbon (0.20 g, catalytic, 25 mL THF) to give the crude amine. The resulting filtrate was reacted with HOBT (0.07 g, 0.51 mmol), the product of Preparation 37 (0.20 g, 0.51 mmol), and DCC (0.12 g, 0.51 mmol) to yield the desired 10 product (0.17 g, 45% yield) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₄₂H₄₉N₇O₆; 68.93 C, 6.75 H, 13.40 N; found 67.02 C, 6.54 H, 12.71 N; ISMS (M+) - 732.

15

Preparation 50



Prepared as in Preparation 38 using the product of Preparation 24 (0.30 g, 0.82 mmol) and 5% palladium on carbon (0.30 g, catalytic, 25 mL THF) to give the crude amine. The resulting filtrate was reacted with HOBT (0.11 g, 0.82 mmol), the product of Preparation 36 (0.31 g, 0.82 mmol), and DCC (0.19 g, 0.90 mmol) to yield the desired product (0.32 g, 56% yield) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₄₀H₅₀N₆O₅; 69.14 C, 7.25 H, 12.09 N; found 67.82 C, 7.07 H, 11.62 N; ISMS (M+) - 695.

633

Preparation 51

Prepared as in Preparation 38 using the product of Preparation 25 (0.35 g, 0.84 mmol) and 5% palladium on carbon (0.35 g, catalytic, 25 mL THF) to give the crude amine. The resulting filtrate was reacted with HOBT (0.11 g, 0.84 mmol), the product of Preparation 36 (0.32 g, 0.84 mmol), and 10 DCC (0.17 g, 0.92 mmol) to yield the desired product (0.22 g, 35% yield) as a light yellow foam: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₄₄H₅₆N₆O₅; 70.56 C, 7.54 H, 11.22 N; found 70.22 C, 7.58 H, 11.21 N; ISMS (M+) - 749.

Preparation 52

Prepared as in Preparation 38 using the product of 20 Preparation 26 (0.42 g, 1.15 mmol) and 5% palladium on carbon (0.40 g, catalytic, 25 mL THF)

634

to yield the crude amine. The filtrate was reacted with HOBT (0.16 g, 1.15 mmol), the product of Preparation 30 (0.44 g, 1.15 mmol), and DCC (0.26 g, 1.28 mmol) to yield the desired product (0.41 g, 51% yield) as a light yellow foam: 1 H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{39}H_{48}N_6O_6$; 67.22 C, 6.94 H, 12.06 N; found 67.66 C, 6.95 H, 11.66 N; ISMS (M+) - 697.

10 Exampl

5

15

20

25

A solution of the product of Preparation 38 (0.52 g, 0.80 mmol) in dichloromethane (20 mL) was stirred under nitrogen with anisole (0.4 mL) and triflouroacetic acid (4.0mL) at ambient temperature for 3 hours. The mixture was concentrated in vacuo to approximately 5 mL and excess diethyl ether added. The mixture was filtered and rinsed with diethyl ether to yield the desired product (0.40 g, 65% yield) as an off white solid: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C34H40N6O8F6; 52.71 C, 5.20 H, 10.85 N; found 52.60 C, 5.08 H, 10.69 N; FDMS (M+) - 546.

Example 2

Prepared as in Example 1 using the product of Preparation 39 (0.86 g, 1.22 mmol), triflouroacetic acid (4.0 mL), anisole (0.4 mL), and dichloromethane (20 mL) to yield the desired product (0.86 g, 85%) as a pale yellow solid: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₃₆H₄₂N₆O₁₀F₆; 51.92 C, 5.08 H, 10.09 N; found 51.63 C, 4.96 H, 10.22 N; FDMS (M+) - 604.

Example 3

Prepared as in Example 1 using the product of Preparation 40 (0.95 g, 1.41 mmol), triflouroacetic acid (4.0 mL), anisole (0.4 mL), and dichloromethane (20 mL) to yield the desired product (0.82 g, 92%) as a pale yellow solid: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₃₆H₄₄N₆O₈F₆; 53.86 C, 5.53 H, 10.47

PCT/US98/17201 WO 99/08697

636

N; found 52.73 C, 5.50 H, 10.07 N; FDMS (M+) -574.

Example 4

5 .

10

15

A solution of the product of Preparation 41 (0.69 g, 1.02 mmol) in dichloromethane (10 mL) was stirred under nitrogen with anisole (0.2 mL) and triflouroacetic acid (4.0mL) at temperature for 3 hours. The mixture was quenched with saturated sodium bicarbonate and stirred 10 min. at ambient temperature. Dichloromethane was added and the mixure washed with bicarbonante and brine. The organic layer was dried over sodium sulfate, concentrated in vacuo, and redissolved in 2 mL ethyl acetate. Diethyl ether (saturated HCl (g), 5 mL) was added and the mixture stirred 10 The mixture was filtered to yield the desired product (0.57 g, 86% yield) as a white solid: ¹H NMR (300 MHz, CDCl₃) - consistent with 20 structure; Anal. calcd. for $C_{32}H_{42}N_6O_4Cl_2$; 59.35 C, 6.85 H, 12.98 N; found 58.74 C, 6.77 H, 12.85 N; ISMS (M+) - 575.

Exampl 5

Prepared as in Example 4 using the product of Preparation 42 (0.26 g, 0.37 mmol), triflouroacetic acid (4.0 mL), anisole (0.4 mL), and dichloromethane (20 mL) to yield the desired product (0.19 g, 76%) as a pale yellow solid: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₃₄H₄₈N₆O₄Cl₂; 60.44 C, 7.16 H, 12.44 N; found 60.08 C, 7.03 H, 12.06 N; ISMS (M+) - 603.

Example 6

Prepared as in Example 4 using the product of Preparation 44 (0.19 g, 0.29 mmol), triflouroacetic acid (4.0 mL), anisole (0.4 mL), and dichloromethane (20 mL) to yield the desired product (0.16 g, 84%) as a pale yellow solid: ¹H NMR (300 MHz, CDCl₃) - consistent with structure;

638

Anal. calcd. for $C_{30}H_{42}N_6O_4Cl_2$; 57.97 C, 6.81 H, 13.52 N; found 57.54 C, 6.36 H, 13.04 N; FDMS (M+) - 549.

Example 7

5

10

Prepared as in Example 4 using the product of Preparation 45 (0.84 g, 1.29 mmol), triflouroacetic acid (4.0 mL), anisole (0.4 mL), and dichloromethane (20 mL) to yield the desired product (0.69 g, 86%) as a pale yellow solid: ^{1}H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{29}H_{38}N_{6}O_{5}Cl_{2}$; 55.86 C, 6.47 H, 13.48 N; found 55.31 C, 6.52 H, 13.01 N; ISMS (M+) - 551.

15

Example 8

Prepared as in Example 4 using the product of 20 Preparation 46 (0.75 g, 1.13 mmol), triflouroacetic acid (4.0 mL), anisole (0.4 mL),

and dichloromethane (20 mL) to yield the desired product (0.62 g, 87%) as a pale yellow solid: ^{1}H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{30}H_{37}N_{7}O_{4}Cl_{2}$; 56.96 C, 6.21 H, 15.50 N; found 55.48 C, 6.03 H, 14.63 N; ISMS (M+) - 560.

Example 9

Prepared as in Example 4 using the product of Preparation 43 (0.82 g, 1.16 mmol), triflouroacetic acid (4.0 mL), anisole (0.4 mL), and dichloromethane (20 mL) to yield the desired product (0.71 g, 90%) as a pale yellow solid: ¹H

NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₃₃H₄₆N₆O₅Cl₂; 58.49 C, 6.84 H, 12.40 N; found 55.40 C, 6.48 H, 11.80 N; ISMS (M+) - 605.

Example 10

Prepared as in Example 4 using the product of Preparation 47 (0.76 g, 1.05 mmol), triflouroacetic acid (2.0 mL), anisole (0.2 mL), and dichloromethane (8.0 mL) to yield the desired product (0.76 g, 85% yield) as an off white solid:

¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₄₀H₄₄N₆O₈F₆; 56.47 C, 5.21 H, 9.88 N; found 56.24 C, 5.32 H, 9.86 N; ISMS (M+) - 623.

Example 11

Prepared as in Example 4 using the product of Preparation 48 (0.52 g, 0.72 mmol), triflouroacetic acid (4.0 mL), anisole (0.4 mL), and dichloromethane (20 mL) to yield the desired

641

product (0.47 g, 94%) as a pale yellow solid: ^{1}H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for $C_{37}H_{46}N_{6}O_{3}Cl_{2}$; 64.06 C, 6.68 H, 12.11 N; found 62.18 C, 6.59 H, 11.78 N; ISMS (M+) - 621.

Example 12

10

15

5

Prepared as in Example 4 using the product of Preparation 50 (0.32 g, mmol), triflouroacetic acid (4.0 mL), anisole (0.4 mL), and dichloromethane (20 mL) to yield the desired product (0.26 g, %) as a pale yellow solid: ^{1}H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₃₅H₄₄N₆O₃Cl₂; 62.96 C, 6.64 H, 12.59 N; found 60.05 C, 6.31 H, 11.98 N; FDMS (M+) - 595.

Example 13

Prepared as in Example 4 using the product of Preparation 51 (0.22 g, 0.29 mmol), triflouroacetic acid (4.0 mL), anisole (0.4 mL), and dichloromethane (20 mL) to yield the desired product (0.19 g, %) as a pale yellow solid: ¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₃₉H₅₀N₆O₃Cl₂; 64.90 C, 6.98 H, 11.64 N; found 66.48 C, 7.24 H, 11.96 N; FDMS (M+) - 649.

Example 14

Prepared as in Example 4 using the product of Preparation 49 (0.96 g, 1.31 mmol), triflouroacetic acid (4.0 mL), anisole (0.4 mL), and dichloromethane (20 mL) to yield the desired product (0.54 g, 59%) as a pale yellow solid: ¹H NMR (300 MHz, CDCl₃) - consistent with structure;

643

Anal. calcd. for $C_{37}H_{43}N_7O_3Cl_2$; 63.06 C, 6.15 H, 13.91 N; found 58.22 C, 5.48 H, 12.32 N; ISMS (M+) - 632.

5

Example 15

Prepared as in Example 1 using the product of Preparation 52 (0.41 g, 0.59 mmol), triflouroacetic acid (2.0 mL), anisole (0.2 mL), and dichloromethane (8.0 mL) to yield the desired product (0.48 g, 99% yield) as an off white solid:

¹H NMR (300 MHz, CDCl₃) - consistent with structure; Anal. calcd. for C₃₆H₄₂N₆O₆F₆; 55.34 C, 5.13 H, 10.19 N; found 55.60 C, 4.92 H, 9.89 N; ISMS (M+) - 597.

644

EXAMPLES PART 4

Preparation 1

5

To a solution of Preparation 306 from Examples Part 2 (1.58 g, 10 mmol) and (1R, 2S)-(+)-cis-1amino-2-indanol (1.49 g, 10 mmol) in methanol (10 mL) was added sodium bicarbonate (840 mg, 10 mmol) and water (5 mL). The resulting 10 mixture was stirred at room temperature overnight. The solvent was removed in vacuo and the residue partitioned between ethyl acetate and brine. The organic phase was dried (MgSO₄) 15 and concentrated in vacuo. The residue was purified by flash chromatography on silica eluting with 80% ethyl acetate / hexanes then 100% ethyl acetate, to give 1.65 g, 67% of the desired product as a tan foam. ^{1}H NMR (CDCl₃) δ 7.70 (s, 1H), 7.5-7.3 (m, 4H), 7.2 (d, J=9 Hz,20 1H), 5.65 (d, J=5 Hz, 1H), 4.97 (m, 1H), 3.83(br s, 1H), 3.35 (dd, J= 6.5, 16 Hz, 1H), 3.12(dd, J= 4, 16 Hz, 1H).

645

Preparation 2

To a solution of Preparation 1 (490 mg, 2 mmol) in dry DMF (6 mL) was added tbutyldimethylsilyl chloride (300 mg, 2 mmol) and imidazole (136 mg, 2 mmol). The solution was stirred at room temperature for 24 h and 10 an additional 50 mg of t-butyldimethylsilyl chloride was added. After stirring an additional 16 h the solvent was removed in vacuo. Flash chromatography of the residue, eluting with 50% ethyl acetate / hexanes gave 540 mg, 75% of the desired product as a white 15 solid. ¹H-NMR is consistent with the desired product. MS (ion spray) 360 (M+1). Anal. Calcd for $C_{18}H_{25}N_3O_3Si$, C: 60.14, H: 7.01, N: 11.69. Found: , C: 60.40, H: 6.91, N: 11.58. [a]_D +51.6 20 (c=10).

Pr paration 3

10% Pd on carbon (350 mg) was wet under nitrogen atmosphere with dry THF (5 mL). A solution of Preparation 2 (500 mg, 1.39 mmol) in dry THF (15 mL) was added and the mixture hydrogenated for 2 h under 40 psi hydrogen gas 10 pressure. The catalyst was removed by filtration through celite and the filtrate concentrated in vacuo to approximately a volume of 10 mL. To this solution was added Preparation 4 of Examples Part 1 (532 mg, 1.4 15 mmol), HOBt hydrate (214 mg, 1.4 mmol), DCC (289 mg, 1.4 mmol), and the mixture stirred at room temperature overnight. The precipitated DCU was removed by filtration. The filtrate was diluted with ethyl acetate and the organic solution washed with 1N hydrochloric acid 20 solution, saturated aqueous sodium bicarbonate solution, and brine then dried (MgSO4) and concentrated in vacuo. Flash chromatography of

the residue, eluting with 80% ethyl acetate /

647

hexanes gave 570 mg, 59% of the desired product as a white foam. ^{1}H -NMR is consistent with the desired product. MS (ion spray) 692 (M+1). IR 1709, 1669 cm $^{-1}$.

5

Example 1

10 A compound of Preparation 3 (440 mg, 0.63 mmol) was dissolved in glacial acetic acid that had been previously saturated with dry hydrogen chloride gas (10 mL, approx. 3N in HCl). The mixture was stirred at room temperature for 30 min, then water (0.5 mL) was added and the 15 mixture stirred an additional 1 h. The mixture was concentrated in vacuo, toluene was added and the solvent removed in vacuo to give a tan solid. This material was converted to the free base by partitioning between chloroform and 20 saturated aqueous sodium bicarbonate solution. The organic phase was dried (MgSO₄) and concentrated in vacuo. Flash chromatography of the residue, eluting with 5% methanol /

648

chloroform containing a trace of ammonium hydroxide gave two products. The more polar product, 80 mg, 23% yield was the desired compound. MS 478 (M+). The more polar product was reconverted to the bis hydrochloride salt by dissolving in chloroform and adding ethyl ether that had been previously saturated with dry hydrogen chloride gas. 1 H NMR (DMSO-d₆) δ 11.25 (br s, 1H), 8.6 (d, J=7 Hz, 1H), 8.4-8.1(m, 5H), 7.4-7.15 (m, 10H), 7.02 (s, 1H), 5.75(d, J= 5 Hz, 1H), 4.70 (m, 1H), 4.56 (m, 1H),4.45 (m, 2H), 3.65 (m, 2H), 3.12 (dd, J=6.5, 16 Hz, 1H), 2.82 (dd, J= 4, 16 Hz, 1H), 1.45(s, 6H). The less polar product was identical 15 to acetyl analog (the product from Example 172, Examples Part 2A) (160 mg, 48% yield).

Preparation 4

20

25

10

To a 0°C solution of Preparation 1 (490 mg, 2 mmol) in pyridine (10 mL) was added acetic anhydride (0.2 mL, 2 mmol). The mixture was allowed to warm to room temperature while stirring overnight. The solvent was removed in vacuo and the residue dissolved in ethyl

649

acetate. The organic solution was washed with 1N hydrochloric acid solution, saturated aqueous sodium bicarbonate solution, and brine then dried (MgSO₄) and concentrated in vacuo.

- Flash chromatography of the residue on silica gel, eluting with 50% ethyl acetate / hexanes gave 420 mg, 73% of the desired product as a colorless oil. ¹H-NMR is consistent with the desired product. MS (ion spray) 288 (M+1).
- 10 Anal. Calcd for $C_{14}H_{13}N_3O_4$, C: 58.53, H: 4.56, N: 14.63. Found: , C: 58.30, H: 4.62, N: 14.41.

Racemic trans-1-amino-2-hydroxyindan was prepared from racemic indene oxide according to the method of D. R. Boyd, N. D. Sharma, N. I. Bowers, P. A. Goodrich, M. R. Groocock, A. J. Blacker, D. A. Clarke, T. Howard, H. Dalton. Tetrahedron Asymmetry 7, 1559-1562, 1996.

650

Pr paration 5

To a solution of Preparation 306 from Examples Part 2 (1.85 g, 11.67 mmol) and trans-1-amino-2hydroxyindan (1.74 g, 11.76 mmol) in methanol (15 mL) was added sodium bicarbonate (1.0 g, 11.9 mmol) and water (8 mL). The resulting mixture was stirred at room temperature 10 overnight. The solvent was removed in vacuo and the residue partitioned between ethyl acetate and brine. The organic phase was dried (MgSO₄) and concentrated in vacuo. The residue was purified by flash chromatography on silica gel 15 eluting with 80% ethyl acetate / hexanes to give 1.73 g, 60% of the desired product as a tan foam. ¹H -NMR is consistent with the desired

product. MS (ion spray) 246 (M+1). Anal. Calcd

20 for $C_{12}H_{11}N_3O_3$, C: 58.77, H: 4.52, N: 17.13. Found: , C: 58.87, H: 4.65, N: 17.18.

651

Preparation 6

To a solution of Preparation 5 (290 mg, 1.18 5 mmol) in pyridine (10 mL) was added acetic anhydride (0.2 mL, 2 mmol). The mixture stirred at room temperature overnight. The solvent was removed in vacuo and the residue dissolved in ethyl acetate. The organic solution was washed 10 with 1N hydrochloric acid solution, saturated aqueous sodium bicarbonate solution, and brine then dried (MgSO₄) and concentrated in vacuo to give 280 mg, 82% yield of the desired product as an off-white foam. H-NMR is consistent with 15 the desired product. MS (ion spray) 288 (M+1). IR 1744, 1549 cm⁻¹

652

10% Pd on carbon (260 mg) was wet under a nitrogen atmosphere with dry THF (5 mL). A solution of Preparation 6 (275 mg, 0.95 mmol) in dry THF (7 mL) was added and the mixture hydrogenated for 1.5 h under 40 psi hydrogen gas pressure. The catalyst was removed by filtration through celite and the filtrate 10 concentrated in vacuo to approximately a volume of 8 mL. To this solution was added Preparation 4 of Examples Part 1(380 mg, 1.0 mmol), HOBt hydrate (153 mg, 1.0 mmol), DCC (206 mg, 1.0 mmol), and the mixture stirred at room 15 temperature for 60 h. The precipitated DCU was removed by filtration. The filtrate was diluted with ethyl acetate and the organic solution washed with 1N hydrochloric acid solution, 20 saturated aqueous sodium bicarbonate solution, and brine then dried (MgSO₄) and concentrated in vacuo. Radial chromatography of the residue, eluting with 80% ethyl acetate / hexanes gave 310 mg, 52% of the desired product. 1H-NMR is

653

consistent with the desired product. MS (ion spray) 620 (M+1). Anal. Calcd for $C_{33}H_{41}N_5O_7$, C: 63.96, H: 6.67, N: 11.30. Found: , C: 63.30, H: 6.86, N: 11.13.

5

Example 2

A compound of Preparation 7 (290 mg, 0.46 mmol) 10 was dissolved in glacial acetic acid that had been previously saturated with dry hydrogen chloride gas (10 mL, approx. 3N in HCl). The mixture was stirred at room temperature for 20 min then concentrated in vacuo, toluene was 15 added and the solvent removed in vacuo to give a yellow solid. This material was converted to the free base by partitioning between ethyl acetate and 0.1N aqueous sodium hydroxide solution. The organic phase was dried (MgSO₄) 20 and concentrated in vacuo. Flash chromatography of the residue, eluting with 5% methanol / chloroform gave the desired product. The hydrochloride salt was reformed by dissolving

654

the product in glacial acetic acid that had been previously saturated with dry hydrogen chloride gas (5 mL, approx. 3N in HCl), then toluene was added and the solvent removed in vacuo to give 140 mg, 50% yield of the desired product a white solid. ^{1}H NMR (DMSO-d₆) δ 11.05 (br s, 1H), 8.55 (d, J= 8 Hz, 1H), 8.24 (br s,2H), 8.10 (br s, 1H), 7.22-7.05 (m, 10H), 5.95 (d, J= 7 Hz, 1H), 5.50 (m, 1H), 5.30 (br s,2H), 4.78 (m, 1H), 4.52, (s, 2H), 3.73 (m, 2H), 10 3.52 (dm, J= 16 Hz, 1H), 2.98 (dd, J= 7, 16 Hz,1H), 2.04 (s, 3H), 1.50 9s, 6H). MS (ion spray) 520 (M+1). Anal. Calcd for $C_{28}H_{35}Cl_2N_5O_5+0.4$ $H_2O.$ C: 56.02, H: 5.96, N: 11.67. Found: , C: 56.33, H: 6.33, N: 11.27. 15

Preparation 8

20 Preparation 391 of Examples Part 2A (950 mg, 4.35 mmol) was dissolved in THF (20 mL) and methyl isocyanate (0.3 mL, 5 mmol) was added. After 4 h of stirring there was no apparent change by TLC. An additional 10 mmol of methyl isocyanate was added along with triethyl amine (0.7 mL, 5 mmol) and the mixture was stirred an additional 60 h. The white precipitate that

655

formed was collected by filtration and dried in vacuo to give 1.04 g, 87% of the desired urea. $^{1}\text{H-NMR}$ is consistent with the desired product. MS (ion spray) 276 (M+1). Anal. Calcd for $C_{12}H_{13}N_{5}O_{3}$, C: 52.36, H: 4.76, N: 25.44. Found: , C: 52.61, H: 4.74, N: 25.25.

Preparation 9

10

15

20

10% Pd on carbon (550 mg) was wet under a nitrogen atmosphere with ethyl acetate (5 mL). A slurry of Preparation 8 (550 mg, 2.0 mmol) in ethyl acetate (40 mL), methanol (40 mL) and acetic acid (1 mL) was added and the mixture hydrogenated for 2 h under 40 psi hydrogen gas pressure. The catalyst was removed by filtration through celite and the filtrate concentrated in vacuo. Toluene was added and the mixture reconcentrated in vacuo to give a purple solid. This solid was dissolved in THF (20 mL) and to this solution was added

656


Preparation 4 of Examples Part 1 (760 mg, 2.0 mmol), HOBt hydrate (306 mg, 2.0 mmol), DCC (412 mg, 2.0 mmol), and triethyl amine (0.28 mL, 2.0 mmol). The mixture stirred at room temperature overnight. The precipitated DCU was removed by filtration. The filtrate was diluted with ethyl acetate and the organic solution washed with 1N hydrochloric acid solution, saturated aqueous sodium bicarbonate solution, and brine then dried (MgSO₄) and concentrated in 10 vacuo. Flash chromatography of the residue, eluting with a step gradient of 2% methanol / ethyl acetate, to 5% methanol / ethyl acetate, to 5% methanol / 1% acetic acid / ethyl acetate, to 10% methanol / 1% acetic acid / 15 ethyl acetate gave 240 mg, 19% of the desired product as an off-white solid. 1H-NMR is consistent with the desired product. MS (ion spray) 608 (M+1).

657

A compound of Preparation 9 (210 mg, 0.34 mmol) 5 was dissolved in glacial acetic acid that had been previously saturated with dry hydrogen chloride gas (5 mL, approx. 3N in HCl). The mixture was stirred at room temperature for 1 h then concentrated in vacuo, toluene was added and the solvent removed in vacuo to give a gum. 10 The residue was dissolved in methanol (2 mL), toluene was added and the solvents removed in vacuo to give 190 mg of the desired product as white powder. ¹H NMR (DMSO-d₆) d 11.20 (br s, 1H), 8.71 (br s, 1H), 8.62 (d, J=7.5 Hz, 1H), 15 8.25 (br's, 2H), 7.72 (d, J=8.5 Hz, 1H), 7.34-7.12 (m, 9H), 7.04 (t, J=7.5 Hz, 1H), 6.86 (br)s, 1H), 5.42 (ABq, J = 14 Hz, 2H), 4.73 (q, J = 7Hz, 1H), 4.53 (s, 2H), 3.73 (m, 2H) 2.65 (s, 20 3H), 1.50 (s, 6H). MS (ion spray) 508 (M+1).

658

Preparation 10



5 3- Aminobenzyl alcohol(615 mg, 5 mmol) was dissolved in methylene chloride (10 mL) and methyl isocyanate (0.3 mL, 5 mmol) was added. The mixture was stirred at room temperature overnight. The white precipitate that formed was collected by filtration and dried in vacuo to give 775 mg, 86% of the desired urea. ¹H-NMR is consistent with the desired product. MS (ion spray) 181 (M+1). Anal. Calcd for C₉H₁₂N₂O₂, C: 59.99, H: 6.71, N: 15.55. Found: C: 60.23, H: 6.68, N: 15.37.

Preparation 11

15

To a 0°C solution of Preparation 10 (540 mg, 3 mmol) in THF (30 mL) was added triethyl amine (0.46 mL, 3.3 mmol) and methanesulfonyl chloride (0.255 mL, 3.3 mmol). The mixture was stirred at 0°C for 1 h at which time TLC (EtOAc)

659

indicated the alcohol had been consumed and a less polar product had formed. The solvent was removed in vacuo and the residue dissolved in dry DMF (10 mL). Potassium carbonate (966 mg, 7 mmol) and 4-nitro imidazole (339 mg, 3 mmol) were added, and the mixture was stirred at room temperature overnight. The DMF was removed in vacuo, the residue dissolved in chloroform/isopropanol (3/1), and washed with water, brine, then dried (MgSO₄) and concentrated to a solid. Flash chromatography on silica gel eluting with 3% methanol in ethyl acetate gave the desired product 410 mg 50%, contaminated with approximately 20% of the undesired 4-alkylated isomer (by NMR analysis). Anal. Calcd for $C_{12}H_{13}N_5O_3$, C: 52.36, H: 4.76, N: 25.44. Found: , C: 52.66, H: 4.64, N: 25.31.

15

660

Preparation 12

10% Pd on carbon (380 mg) was wet under a nitrogen 5 atmosphere with ethyl acetate (20 mL). A solution of Preparation 11 (380 mg, 1.4 mmol) in methanol (20 mL) was added and the mixture hydrogenated for 2 h under 40 psi hydrogen gas pressure. The catalyst was removed by filtration through celite and the filtrate concentrated in 10 vacuo. THF was added and the mixture reconcentrated in vacuo to give a solid. This solid was dissolved in THF (10 mL) and to this solution was added Preparation 4 from Examples Part 1(532 mg, 1.4 mmol), HOBt hydrate (214 mg, 1.4 mmol), DCC 15 (289 mg, 1.4 mmol). The mixture stirred at room temperature overnight. The precipitated DCU was removed by filtration. The filtrate was diluted with ethyl acetate and the organic solution washed with saturated aqueous sodium bicarbonate solution, and brine then dried (MgSO₄) and concentrated in vacuo. Flash chromatography of the residue, eluting with a 20 step gradient of 2% methanol / ethyl acetate, to 10% methanol / ethyl acetate gave 400 mg, of impure product as an off-white solid. Radial chromatography, 5% methanol / chloroform gave the desired product, 370mg, 44% as a

661

yellowish tinted foam. H-NMR is consistent with the desired product. MS (ion spray) 608 (M+1). Anal. Calcd for C31H41N7O6 +0.2CHCl₃ C: 59.33, H: 6.58, N: 15.52. Found: , C: 59.21, H: 6.65, N: 15.18.

5

15

20

Example 5

To a compound of Preparation 12 (0.3 g, 0.5 mmol) stirring in dichloromethane (6 mL) at room temperature was added 10 trifluoroacetic acid (2 mL). After 2 h, the reaction mixture was concentrated to dryness. The residue was partitioned between ethyl acetate and saturated sodium bicarbonate and was then extracted with ethyl acetate. combined organics were washed with brine, dried over sodium sulfate and concentrated to dryness. The residue was chromatographed over silica gel using ethanol/chloroform as eluant to yield 0.12 g (49%) of the desired product as a white foam. 1H-NMR is consistent with structure; MS (ion spray) 508.3 (M+1); Anal. Calc'd for $C_{26}H_{33}N_7O_4^{\circ}0.41CHCl_3$: C, 57.00; H, 6.05; N, 17.62. Found: C, 57.17; H, 6.24; N, 17.22.

662

EXAMPLES PART 5

Preparation 1

Methyl 2R-{2-[(tert-butoxy)carbonylamino]-2-methylpropanoylamino)-2-phenylacetate

5

To a solution of N-[(1,2-dimethylethoxy)carbonyl]-2methylalanine (5.04 g, 24.79 mmol) and 2-chloro-4,6dimethoxy-1,3,5-triazine (4.43 g, 24.79 mmol) in 100 mL 10 of THF was added N-methylmorpholine (5.45 mL, 49.58 mmol). After 1h, (R)-phenylglycine methyl ester hydrochloride (5.0 g, 24.79 mmol) was added in one portion, and the reaction mixture was stirred at room temperature for 16 h. The reaction was concentrated in vacuo, and the resulting residue was dissolved in 100 mL 15 each of EtOAc and 10 % (w/w) aqueous citric acid. phases were separated and the organic layer was washed with saturated aqueous NaHCO3 and saturated aqueous NaCl, dried over Na₂SO₄, filtered and concentrated in vacuo to give 8.51 g (98 %) of a white solid. ¹H NMR (DMSO, 300 20 MHz) d 8.05 (m, 1H), 7.40 (m, 5H), 6.90 (br s, 1H), 5.40 (d, 1H), 3.65 (s, 3H), 1.30 (m, 15H); ¹³C NMR (DMSO, 75 MHz) 174.2, 170.9, 154.3, 136.8, 128.4, 128.1, 128.0, 127.4, 78.2, 56.0, 52.3, 28.0, 25.2, 24.7 ppm; Anal. 25 Calcd. for $C_{18}H_{26}N_2O_5$: C, 61.70; H, 7.48; N, 7.99. Found: C, 61.50; H, 7.36; N, 8.19.

663

Preparation 2

(R)-2-[(Tert-butoxy)carbonylamino]-2-methyl-N-({N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]carbamoyl)phenylmethyl)propanamide

5

10

To a 0 °C solution of the methyl ester in Example 1, (1.0 g, 2.85 mmol) in dioxane (19 mL) and H_2O (9.5 mL) was added LiOH (0.682 g, 28.5 mmol). The reaction stirred for 40 min and was diluted with 50 mL of EtOAc. The pH was adjusted to 3 with conc. HCl and the phases were separated. The aqueous layer was extracted with EtOAc (2X25 mL). The combined organics were dried over Na_2SO_4 , filtered and concentrated in vacuo to give 958 mg of the acid which was used as is in subsequent reactions.

15 N-methylmorpholine (0.25 mL, 2.27 mmol) was added to a suspension of 2-chloro-4,6-dimethoxy-1,3,5-triazine (405 mg, 2.27 mmol) and $2-\{2-[(\text{tert-butoxy})\}$ carbonylamino]-2-methylpropanoylamino}-2-phenylacetic acid (570 mg, 1.69 mmol) in CH_2Cl_2 (7 mL) and THF (2 mL) at 23 °C. The mixture stirred for 2h and 2-(4-20 aminoimidazolyl) -2-phenyl-1-pyrrolidinylethan-1-one dihydrochloride (779 mg, 2.27 mmol) was added in one portion. After 16 h, the reaction was diluted with EtOAc (50 mL) and 10% (w/w) aqueous citric acid. The phases 25 were separated, and the organic layer was washed with 20 mL each of saturated aqueous NaHCO3 and saturated aqueous NaCl, dried over Na2SO4, filtered and concentrated in vacuo. The residue was purified by flash chromatography (gradient 2/98 to 6/94 MeOH/CH₂Cl₂) to give 405 mg (40 %) of the title compound. 1H NMR (CDCl₃, 300 MHz) d 10.61 (m, 30 1H), 7.75 (s, 1H), 7.45-7.15 (m, 12H), 5.95 (s, 1H), 5.80(m, 1H), 3.60-3.38 (m, 3H), 3.10 (m, 1H), 2.15 (m, 1H), 1.98-1.65 (m, 3H), 1.50-1.05 (m, 15H); IR (CHCl₃) 3400,

664

3000,1706, 1661, 1486, 1455, 1157 cm⁻¹; MS (electrospray) m/z 588 (M+H), 589 (M-H), 587.

Example 1

(R)-2-Amino-2-methyl-N-({N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]carbamoyl}
phenylmethyl)propanamide, bis-trifluoroacetate

Trifluoroacetic acid (1.4 mL, 18.3 mmol) was added to a solution of a compound of Example 2 (360 mg, 0.611 10 mmol), anisole (0.20 mL, 1.83 mmol) and CH_2Cl_2 (14 mL). After 4 h the solution was concentrated to an oil which was dissolved in CH_2Cl_2 (3.0 mL). This solution was added dropwise to Et₂O (25 mL) with vigorous stirring, and a white precipitate formed. After 12 h, the solid was 15 filtered and washed with Et2O (20 mL). The wet cake was dried at 40 °C in a vacuum oven for 12 h to yield 391 mg (91 %). ¹H NMR (DMSO-d₆, 300 MHz) d 10.70 (d, J = 2.5 Hz, 1H), 8.66 (d, J = 6.7 Hz, 1H), 8.02 (m, 3H), 7.54-7.10(m, 12H), 6.39 (s, 1H), 5.58 (m, 1H), 3.60 (m, 1H), 3.30 20 (m, 2H), 2.90 (m, 1H), 1.80-1.55 (m, 4H), 1.07-0.70 (m, 2H)6H); IR (KBr) 3064, 3035, 2980, 1672, 1575, 1539, 1497, 1201, 1133, 721, 700 cm⁻¹; MS (electrospray) m/z(freebase) 488; (freebase M+H), 489 (freebase M-H), 487.

665

Preparation 3

(R)-2-[(Tert-butoxy)carbonylamino]-2-methyl-N-[(N-{1-[2-(4-methylpiperidyl)-2-oxo-1-phenylethyl] imidazol-4-yl)carbamoyl)phenylmethyl] propanamide

5

N-methylmorpholine (0.16 mL, 1.49 mmol) was added to a suspension of 2-chloro-4,6-dimethoxy-1,3,5-triazine (268 mg, 1.53 mmol), 2-{2-[(tert-butoxy)carbonylamino]-2methylpropanoylamino}-2-phenylacetic acid (502 mg, 1.49 10 mmol), CH_2Cl_2 (7 mL) and THF (2 mL) at 23 °C. heterogeneous reaction stirred for two hours and 2-(4aminoimidazolyl) -1-(4-methylpiperidyl)-2-phenylethan-1one dihydrochloride (613 mg, 1.65 mmol) was added. The mixture was stirred for 15.5 h, and EtOAc (40 mL) and 10 % (w/w) aqueous citric acid (18 mL) were added. 15 layers were separated, and the organic layer was extracted with 15 mL each of saturated NaHCO3 and saturated NaCl. The organic layer was dried (MgSO4), concentrated in vacuo to 590 mg of solid, and purified by flash chromatography (10 % i-PA/CHCl3) to yield 440 mg (48 20 %) of a foamy solid: ^{1}H NMR (CDCl₃, 300 MHz) d 10.34 (m, 1H), 7.67 (s, 1H), 7.40-7.19 (m, 12H), 6.10 (m, 1H), 5.73 (m, 1H), 4.57 (m, 1H), 3.84 (m, 1H), 3.60 (m, 1H), 2.95 (m, 1H), 2.74-2.58 (m, 1H), 1.65-0.79 (m, 23H); ¹³C NMR 25 (CDCl₃, 75 MHz) 174.0, 167.0, 165.3, 137.7, 136.7, 134.2, 133.5, 133.4, 129.4, 129.3, 129.2, 129.2, 128.7, 128.5, 128.5, 128.3, 128.2, 128.0, 127.2, 127.2, 108.1, 62.1, 62.0, 62.0, 61.9, 57.1, 56.6, 46.0, 46.0, 45.8, 429, 34.2, 33.6, 33.5, 30.8, 30.7, 28.2, 21.5, 21.4 ppm; IR $(CHCl_3)$ 1456, 1487, 1560, 1661, 1707, 3010 cm⁻¹; MS 30 (electrospray) m/z 617 (M+H), 438 & 615 (M-H), 436.

666

Example 2

(R)-2-Amino-2-methyl-N-[(N-{1-[2-(4-methylpiperidyl)-2-oxo-1-phenylethyl]imidazol-4-yl}carbamoyl)

phenylmethyl]propanamide bis-trifluoroacetate

5

Trifluoroacetic acid (1.5 mL, 19.9 mmol) was added to a solution of a compound of Example 4 (410 mg, 0.66 mmol), anisole (0.22 mL, 2.0 mmol) and CH_2Cl_2 (16 mL). After 5 h the solution was concentrated to an oil which 10 was dissolved in CH₂Cl₂ (3.5 mL). This solution was added dropwise to Et₂O (30 mL) with vigorous stirring, and a white precipitate formed. After 30 min the solid was filtered and washed with copious Et₂O. The wet cake was dried in a room temperature vacuum desiccator overnight to yield 283 mg (58 %) of a white, hygroscopic powder: 15 mp 182 °C (dec); ¹H NMR (DMSO-d₆, 300 MHz) d 10.79 (d, J =2.7 Hz, 1H), 8.72 (d, J = 6.5 Hz, 1H), 8.15 (m, 2H), 7.54-7.20 (m, 11H), 7.19 (m, 1H), 6.73 (d, J = 14.2 Hz, 1H), 5.61 (d, J = 6.7 Hz, 1H), 4.36 (m, 1H), 3.72 (m, 1H), 2.69-2.48 (m, 2H), 1.59-1.24 (m, 9H), 1.07-0.70 (m, 20 5H); ¹³C NMR (DMSO-d₆, 75 MHz) 171.5, 171.4, 166.4, 165.6, 165.6, 165.4, 165.4, 158.3, 157.9, 137.2, 137.1, 137.0, 136.2, 136.1, 135.6, 135.5, 135.4, 133.2, 133.1, 129.2, 128.9, 128.7, 128.4 128.3, 128.3, 128.1, 128.0, 127.9, 127.6, 107.2, 107.2, 97.8, 60.3, 60.3, 60.0, 57.0, 56.9, 25 56.6, 45.2, 42.2, 42.0, 42.0, 34.0, 33.9, 33.3, 3.2, 33.0, 30.0, 23.2, 23.2, 21.5, 21.4 ppm; IR (KBr) 1137, 1201, 1540, 1576, 1670, 2955 cm⁻¹; MS (electrospray) 517 (freebase M+H), 438 & 515 (freebase M-H).

667

EXAMPLES PART 6

Example 1

Chromatographic Separation

5

10

Diastereomeric secretagogue compounds are separated on a 8 x 15 cm Prochrom® column packed with a Kromasil® CHI-DMB chiral phase. The chromatographic eluent consisted of 12% to 15% 3A alcohol, 0.2% dimethylethylamine in heptane. One to two gram loadings of mixtures of diastereomers have been separated by this technique. The active isomer has been demonstrated to be the second eluting component. Isolation of the desired isomer is completed by evaporation of the solvent by using a roto evaporator.

Since this is a chiral phase, enantiomers of the individual diastereomers are also resolved to yield compounds which are enantiomerically pure.

The procedure for packing the Prochrom® column, separation of the diastereomeric secretagogue SAR candidates, and analysis of the purified compounds is provided as follows:

Materials

25 Chemicals:

Kromasil® CHI-I, 10m CSP Bulk 3A alcohol, QA 041N Heptane, QD440N Dimethylethylamine, Aldrich

30

668

Equipment:

8 cm Prochrom® column

Prochrom® LC-80 pumping system, fraction collector

5 Chromatography:

10

Column Preparation: Slurry 500 grams of Kromasil® CHI bulk packing in 400 ml of propanol. Pack in 8 x 16 cm Prochrom® column. Equilibrate with eluent consisting 13% 3A alcohol, 0.2% dimethylethylamine in heptane at a flow rate of 250 ml/min. Detector setting is 260 nm to 270 nm for the 200 mg loading per run. Change wavelength to 280 nm for 1.0 gram loadings.

15 <u>Sample preparation:</u> Dissolve approximately 1.0 grams of secretagogue mixture in small amount of 3A alcohol and dilute with approximately 50 ml of eluent.

Chromatographic separation: The first isomer from each series elutes in range of 6 to 8 minutes. The second isomer elutes in range of 8 to 10 minutes. Shave the front and back of isomer 1 to eliminate other isomeric impurities.

Fraction work up: Strip fractions to dryness using 25 roto vap.

Sample Analysis Conditions:

30 a. Operating Conditions

Column: 0.46 x 25 cm Kromasil® CHI

Eluent: 15% ethanol, 0.2% dimethylethylamine in

heptane

669

Flow Rate: 1.0 ml/min
Temperature: ambient

uv: 250 nm

Injection: as needed

5 b. System Suitability

I) isomer 1 elutes in 5 to 6 minutes, isomer 2 elutes in 8 to 10 minutes

c. Analysis Procedure

I) calculate the diastereomeric excess of each

10 isolated isomer.

Both diastereomerically and optically pure secretagogue compounds are produced by this process.

670

Exampl 2

Chiral Synthesis of Nitroimidazole (L-isomer)

Preparation 2A

5

Boc-L- α -Phenylglycine (2.0 g, 9.12 mmol) was dissolved in 15mL anhydrous THF and cooled to 0°C. (1.97 g, 9.58 mmol) was added, followed by the addition of 4-methylpiperidine (1.08 mL, 9.12 mmol) and HOBt (1.29 10 g, 9.58 mmol). After stirring 30 min the ice bath was removed and stirring continued overnight at ambient temperature. The reaction was filtered, concentrated, then diluted with EtOAc and washed with 10% aqueous sodium bicarbonate (1 X 50 mL), 0.1N HCl (1 X 50 mL), and 15 dried over sodium sulfate. Filtered, and concentrated to 3.2 g of a crude yellow oil. The oil was purified was radial chromatography eluting with 30:70 EtOAc: Hexanes. The appropriate fractions were concentrated to 2.5g (83% yield) of a colorless foam. H-NMR (d, CDCl₃, 250 MHz, 20 rotamers present) 0.60 (d, J = 7.5 Hz, 1.5 H), 0.77 (d, J)= 7.5 Hz, 1.5 H), 1.00-1.50 (m, 5H), 1.25 (s, 9H), 2.45(m, 1H), 2.80 (m, 1H), 3.62 (m, 1H), 4.43 (m, 1H), 5.43 (m, 1H), 6.02 (d, J = 8 Hz, 1H), 7.10-7.30 (m, 5H); MS 2.5 (ion spray) 333.5 (M+1); $[\alpha]_D = +95.9$ (MeOH); Anal. Calc'd for $C_{19}H_{28}N_2O_3$: C, 68.65; H, 8.49; N, 8.43. Found: C, 69.04; H, 8.51; N, 8.91.

671

Preparation 2B

5 The Boc group was removed from the above compound (1.09 g, 3.28 mmol) by dissolving in 3 mL of methylene chloride and added 3 mL of trifluoroacetic acid. stirring 1 h at room temperature the reaction was concentrated in vacuo (3 X CH2Cl2). The compound was 10 vacuum dried to 1.5 g (100%) of the ditrifluoroacetic acid salt. H-NMR (d, CDCl₃, 400 MHz, equal intensity rotamers present at room temperature) 0.01 (m, 0.5H), 0.70 (d, J = 7.5 Hz, 1.5 H), 0.83 (m, 0.5H), 0.88 (d, J =7.5 Hz, 1.5 H), 1.00-1.60 (m, 4H), 2.55 (m, 1H), 2.90 (m, 1H), 3.52 (m, 1H), 4.35 (m, 1H), 5.42 (m, 1H), 7.32-7.43 15 (m, 5H), 8.18 (br s, 2H), 9.30 (br s, 2H); MS (ion spray) 233.1 (M+1); $[\alpha]_D = +19.7$ (1N HCl).

Preparation 2C

20

25

The above amine (1.5 g, 3.26 mmol) was dissolved in 10mL methanol and 10 mL water and cooled to 0°C. Sodium bicarbonate (0.58 g, 6.85 mmol) was added and stirred 10 min followed by the addition of 1,4-dinitroimidazole. The solution quickly turned bright yellow and after stirring 30 min the ice bath was removed and stirring continued overnight at ambient temperature. The product was extracted into CH₂Cl₂, dried over sodium sulfate,

672


filtered, and concentrated to 0.93 g of a crude white foam. A pure sample is obtained by crystallization from EtOAc (0.64 g, 60%). ¹H-NMR $(d, CDCl_3, 400 \text{ MHz}, rotamers)$ present) 0.05 (m, 0.5H), 0.59 (d, J = 7.5 Hz, 1.5 H), 0.72 (d, J = 7.5 Hz, 1.5 H), 0.79 (m, 0.5H), 0.93 (m,0.5H), 1.14 (m, 0.5H), 1.23-1.56 (m, 3H), 2.44 (m, 1H), 2.82 (m, 1H), 3.36 (m, 1H), 4.37 (m, 1H), 5.88 (s, 0.5H), 5.90 (s, 0.5H), 7.02 (s, 1H), 7.10-7.30 (m, 5H), 7.52 (s, 1.50)1H); MS (ion spray) 329.3 (M+1); $[\alpha]_D = +177.6$ (MeOH); 98% ee by chiral HPLC: Anal. Calc'd for $C_{17}H_{20}N_4O_3$: C, 62.18; H, 10 6.14; N, 17.06. Found: C, 62.25; H, 6.08; N, 17.09.

Example 3 Chiral Synthesis of Nitroimidazole D-isomer

15

20

25



Boc-D- α -Phenylglycine (2.0 g, 9.12 mmol) was dissolved in 15mL anhydrous THF and cooled to 0°C. DCC (1.97 g, 9.58 mmol) was added, followed by the addition of 4methylpiperidine (1.08 mL, 9.12 mmol) and HOBt (1.29 g, 9.58 mmol). After stirring 30 min the ice bath was removed and stirring continued overnight at ambient temperature. The reaction was filtered, concentrated, then diluted with EtOAc and washed with 10% aqueous sodium bicarbonate (1 X 50 mL), 0.1N HCl (1 X 50 mL), and dried over sodium sulfate. Filtered, and concentrated to 3.1 g of a crude yellow oil. The oil was purified was radial chromatography eluting with 30:70 EtOAc: Hexanes. 30 The appropriate fractions were concentrated to 2.2g (73%

673

yield) of a colorless foam. $^{1}\text{H-NMR}$ (d, CDCl₃, 250 MHz, rotamers present) 0.60 (d, J = 7.5 Hz, 1.5 H), 0.77 (d, J = 7.5 Hz, 1.5 H), 1.00-1.50 (m, 5H), 1.25 (s, 9H), 2.45 (m, 1H), 2.80 (m, 1H), 3.62 (m, 1H), 4.43 (m, 1H), 5.43 (m, 1H), 6.02 (d, J = 8 Hz, 1H), 7.10-7.30 (m, 5H); MS (ion spray) 333.5 (M+1); [α]_D = -112.9 (MeOH); Anal. Calc'd for C₁₉H₂₈N₂O₃: C, 68.65;H, 8.49; N, 8.43. Found: C, 68.41; H, 8.11; N, 8.53.

10

5

Preparation 3B

The Boc group was removed from the compound of Preparation 3A (1.0 g, 3.01 mmol) by dissolving in 2 mL of methylene chloride and added 2 mL of trifluoroacetic acid. After stirring 1 h at room temperature the reaction was concentrated in vacuo (3 X CH₂Cl₂). The compound was vacuum dried to 1.4 g (100%) of the ditrifluoroacetic acid salt. ¹H-NMR (d, CDCl₃, 400 MHz, equal intensity rotamers present at room temperature) 0.01 (m, 0.5H), 0.70 (d, J = 7.5 Hz, 1.5 H), 0.83 (m, 0.5H), 0.88 (d, J = 7.5 Hz, 1.5 H), 1.00-1.60 (m, 4H), 2.55 (m, 1H), 2.90 (m, 1H), 3.52 (m, 1H), 4.35 (m, 1H), 5.42 (m, 1H), 7.32-7.43 (m, 5H), 8.18 (br s, 2H), 9.30 (br s, 2H); MS (ion spray) 233.1 (M+1); [\alpha]_D = -38.1 (MeOH).

Preparation 3C

674

The amine from Preparation 3B (1.4 g, 3.04 mmol) was dissolved in 10mL methanol and 10 mL water and cooled to 0°C. Sodium bicarbonate (0.51 g, 6.08 mmol) was added and stirred 10 min followed by the addition of 1,4dinitroimidazole. The solution quickly turned bright vellow and after stirring 30 min the ice bath was removed and stirring continued overnight at ambient temperature. The product was extracted into CH2Cl2, dried over sodium sulfate, filtered, and concentrated to 0.93 g of a crude 10 yellow foam. A pure sample is obtained by crystallization from EtOAc (0.60 g, 60%). H-NMR (d, CDCl₃, 400 MHz, rotamers present) 0.05 (m, 0.5H), 0.59 (d, J = 7.5 Hz, 1.5 H), 0.72 (d, J = 7.5 Hz, 1.5 H), 0.79 (m,15 0.5H), 0.93 (m, 0.5H), 1.14 (m, 0.5H), 1.23-1.56 (m, 3H), 2.44 (m, 1H), 2.82 (m, 1H), 3.36 (m, 1H), 4.37 (m, 1H), 5.88 (s, 0.5H), 5.90 (s, 0.5H), 7.02 (s, 1H), 7.10-7.30 $(m, 5H), 7.52 (s, 1H); MS (ion spray) 329.2 (M+1); <math>[\alpha]_D =$ -175.7 (MeOH); 96% ee by chiral HPLC; Anal. Calc'd for $C_{17}H_{20}N_4O_3$: C, 62.18; H, 6.14; N, 17.06. Found: C, 62.93; 20 H, 6.14; N, 17.21.

675

Example 4 Chiral Synthesis of Nitroimidazole (D-isomer)

Preparation 4A

5

D-(p-hydroxyphenyl)glycine was Boc protected following the procedure of Salituro, JACS, 760 (1990).

10

15

20

25

Example 4B

To a 0°C solution of the phenol compound from Preparation 4A (3.0 g, 11.2 mmol) in 100 mL of DMF was added 1.03 g (60% dispersion, 25.8 mmol) NaH. The mixture was then sonicated at room temperature for 30 min and methyl iodide added (0.77 mL, 12.3 mmol). The reaction becomes near homogenous after an additional 30 min of sonication. An additional 0.2 mL of methyl iodide was added and the reaction stirred overnight at room temperature. reaction was diluted with EtOAc and a few drops of water. The reaction was transferred to a separatory funnel and acidified with a 10% sodium bisulfate solution and brine. The organic layer was separated and dried with sodium sulfate to 3.4 g of a crude yellow oil. This oil was purified by radial chromatography eluting with 80/20/5 CH₂Cl₂/MeOH/NH₄OH. The appropriate fractions were taken up in EtOAc and washed with 10% sodium bisulfate/brine

676

solution, dried over sodium sulfate, filtered and concentrated to 1.5 g (48%) of the desired ether. $^{1}\text{H-NMR}$ (d, CDCl₃, 400 MHz, equal intensity rotamers present) 1.22 (s, 4.5H), 1.40 (s, 4.5 H), 3.77 (s, 3H), 5.02 (d, J = 5 Hz, 0.5 H), 5.22 (d, J = 5 Hz, 0.5H), 5.39 (d, J = 5 Hz, 0.5H), 6.82 (m, 2H), 7.29 (m, 2H), 7.62 (m, 0.5H); MS (ion spray) 280.2 (M-1); [α]_D = -134.1 (MeOH); Anal. Calc'd for C₁₄H₁₉NO₅: C, 59.78;H, 6.81; N, 4.98. Found: C, 60.07; H, 7.01; N, 4.99.

10

5

Preparation 4C

To a 0°C solution of the acid from Preparation 4B (540 mg, 1.92 mmol) and dimethylamine hydrochloride (157 mg, 1.92 15 mmol) in 8 mL of anhydrous DMF was added diethylcyanophosphonate (DECP) (0.29 mL, 1.92 mmol) and triethylamine (0.27 mL, 1.92 mmol). After 30 min at 0°C a second equivalent of dimethylamine hydrochloride, DECP, and triethylamine were added. After an additional 30 min 20 the reaction was complete by TLC. The reaction was diluted with EtOAc , washed with 1:1 10% sodium bisulfate:brine, 1:1 10% sodium bicarbonate:brine, and brine. The organic layer was dried over sodium sulfate, filtered and concentrated to 0.90 g of a light yellow 25 oil. This material was purified by radial chromatography eluting with 1:1 EtOAc: Hexanes. The appropriate fractions were vacuum dried to 0.50 g (85%) of a white solid. ¹H-NMR (d, CDCl₃, 400 MHz) 1.40 (s, 9H), 2.85 (s, 3H), 2.95 (s, 3H), 3.79 (s, 3H), 5.50 (d, J = 10 Hz, 1H), 30 5.96 (d, J = 10 Hz, 1H), 6.83 (d, J = 10 Hz, 2H), 7.29

677

(d, J = 10 Hz, 2H); MS (ion spray) 309.3 (M+1); $[\alpha]_D = -165.5$ (MeOH).

Preparation 4D

5

The Boc group was removed from the compound of Preparation 4C (450 mg, 1.46 mmol) by dissolving in 2 mL of methylene chloride and added 2 mL of trifluoroacetic acid. After stirring 90 min at room temperature the reaction was complete by TLC and HPLC and concentrated in vacuo (3 X CH₂Cl₂). The compound was vacuum dried to 620 mg (97%) of the ditrifluoroacetic acid salt. This material was used without further characterization.

Preparation 4F

The amine from Preparation 4E (637 mg, 1.46 mmol) was dissolved in 10mL methanol and 10 mL water and cooled to 0°C. Sodium bicarbonate (270 mg, 3.21 mmol) was added and stirred 10 min followed by the addition of 1,4-dinitroimidazole. The solution quickly turned bright yellow and after stirring 30 min the ice bath was removed and stirring continued 6h. The product was extracted into CH₂Cl₂, washed with brine, dried over sodium sulfate, filtered, and concentrated to 450 mg of a crude yellow

678

oil. The compound was purified by radial chromatography eluting with EtOAc. The appropriate fractions were dried to 410 mg (92%) of a light yellow solid. $^{1}\text{H-NMR}$ (d, CDCl₃, 400 MHz) 2.88 (s, 3H), 3.01 (s, 3H), 3.80 (s, 3H), 6.01 (s, 1H), 6.94 (d, J = 10 Hz, 2H), 7.29 (d, J = 10 Hz, 2H), 7.37 (s, 1H), 7.68 (s, 1H); MS (ion spray) 305.2 (M+1); $[\alpha]_{D} = -258.2$ (MeOH); Anal. Calc'd for $C_{14}H_{16}N_{4}O_{4}$: C, 55.26;H, 5.30; N, 18.41. Found: C, 55.23; H, 5.27; N, 18.23.

679

Example 5

Alternate Chiral Synthesis of Nitroimidazole-D-isomer)

Preparation 5A

5

D-Phenylglycine (400 mg, 2.53 mmol) was suspended in 5 mL methanol and 5 mL water and cooled to 0°C. Sodium bicarbonate (400 mg, 5.06 mmol) was added and stirred 10 10 min followed by the addition of 1,4-dinitroimidazole. The solution quickly turned bright yellow and after stirring 30 min the ice bath was removed and stirring continued overnight at ambient temperature. The reaction was diluted with EtOAc and acidified with 0.2N HCl. The organic extract was washed with brine and dried over 15 sodium sulfate. Filtered, and concentrated to 620 mg (99%) of a light yellow solid. ¹H-NMR (d, DMSO, 400 MHz) 6.50 (s, 1H), 7.42 (m, 3H), 7.52 (m, 2H), 7.94 (s, 1H), 8.40 (s ,1H); MS (ion spray) 248.4 (M+1); $[\alpha]_D = -146.2$ 20 (MeOH); Anal. Calc'd for $C_{11}H_9N_3O_4 \bullet 0.3 H_2O$: C, 52.30; H, 3.83; N, 16.63. Found: C, 52.17; H, 3.95; N, 16.79.

680

Preparation 5B

The acid from Preparation 5A (215 mg, 0.87 mmol) was dissolved in 10 mL anhydrous THF and cooled to 0°C. Added 4-methylpiperidine (103 μ L, 0.87 mmol), and diethylcyanophosphonate (DECP) (132 μ L, 0.87 mmol). After 45 min a second equivalent of 4-methylpiperidine and DECP were added and stirring continued at 0°C for an additional 45 min. Diluted the reaction with ethyl 10 acetate (300 mL) and washed with aqueous 10% NaHSO4 / brine, 5% NaHCO3 / brine, brine (150 mL each) and dried over NaSO₄. Purified by radial chromatography eluting with EtOAc which gave 236 mg (83%) of the desired product 15 as a yellow oil. An analytical sample was obtained by crystallization from EtOAc. 1H-NMR (d, DMSO, 400 MHz, rotamers present) -0.05 (m, 0.5H), 0.68 (d, J = 7.5 Hz, 1.5 H), 0.80 (m, 0.5H), 0.83 (d, J = 7.5 Hz, 1.5 H), 0.97(m, 0.5H), 1.18 (m, 0.5H), 1.40-1.60 (m, 3H), 2.60 (m, 3H)20 1H), 2.98 (m, 1H), 3.59 (m, 1H), 4.34 (m, 1H), 6.85 (s, 0.5H), 6.92 (s, 0.5H), 7.37-7.50 (m, 5H), 7.78 (s, 1H), 8.12 (s, 0.5H), 8.15 (s, 0.5H); MS (ion spray) 329.2 (M+1); $[\alpha]_D = -204.7$ (MeOH); Anal. Calc'd for $C_{17}H_{20}N_4O_3$: C, 62.18; H, 6.14; N, 17.06. Found: C, 62.47; H, 6.21; N,

25 17.07.

681

EXAMPLES PART 7

Melting points were determined with a Thomas-Hoover capillary melting point apparatus and are uncorrected.

5 Nuclear magnetic resonance studies were performed on a Bruker ARX 500 spectrometer. Merck silica gel 60 F254 plates (0.25mm) were used for thin layer chromatography. Merck silica gel 60 230-400 mesh was used for flash chromatography. Biotage KP-SIL, 60A cartridges were used for Biotage Flash 40 purifications. HPLC conditions: Eluent: 0.1% trifluoroacetic acid in water: acetonitrile at 2 mL/min. Column: Zorbax RX-C8. Detection: 230 nm.

Preparation 1

Ethyl 2-(2-Naphthyl)acetate

15

A steady stream of anhydrous hydrochloric acid was 20 bubbled subsurface into a solution of 2-naphthylacetic acid (251.38 grams, 1.35 mol) dissolved in ethanol (1760 mL) over a period of 10 minutes. The resulting solution was stirred at ambient temperature until complete as determined by hplc (2 hours). The reaction mixture was concentrated to 25 dryness. The resulting oil was dissolved in ethyl acetate (200 mL) and filtered through silica gel (300 grams) eluting the product with ethyl acetate (1400 mL). The filtrate was concentrated to give 286.33 grams (99%) of ethyl 2-(2naphthyl)acetate as a colorless oil. MS (FIA) m/z 215.3 $[(M+H)^{+}]$. ¹H nmr (DMSO-d₆): δ 1.15-1.24 (t, 3H), 3.81-3.86 30 (d, 2H), 4.07-4.15 (q, 2H), 7.41-7.55 (m, 3H), 7.80-7.92 (m, 3H)4H).

682

Preparation 2

Ethyl 2-Bromo-2-(2-naphthyl)acetate

5

A solution consisting of ethyl 2-(2-naphthyl)acetate (1.07 grams, 5.0 mmol), N-bromosuccinimide (0.89 grams, 5.0 mmol), benzoyl peroxide (0.05 grams), and carbon tetrachloride (50 mL) was heated at reflux until complete as 10 determined by hplc (3 hours). The reaction was cooled to ambient temperature, washed with water (2 x 25 mL), dried using sodium sulfate, and filtered. The filtrate was concentrated to dryness. The residue was purified using a 15 Biotage Flash 40M system eluting with hexane : ethyl acetate (49:1) to give 1.20 grams (82%) of ethyl 2-bromo-2-(2naphthyl) acetate, mp 80-82° C. MS (FIA) m/z 293.0 [(M+H) $^{+}$]. Anal. calcd. for $C_{14}H_{13}O_{2}Br$: C: 57.36; H: 4.47. Found: C: 57.62; H: 4.54. ¹H nmr (CDCl₃): δ 1.27-1.33 (t, 3H), 4.18-4.36 (m, 2H), 5.56 (s, 1H), 7.52-7.55 (m, 2H), 20 7.71-7.76 (m, 1H), 7.82-7.92 (m, 3H), 7.97 (s, 1H).

Preparation 3

Ethyl 2-(2-Naphthyl)-2-(4-nitroimidazolyl)acetate

683

A yellow slurry consisting of ethyl 2-bromo-2-(2naphthyl) acetate (384.04 grams, 1.31 mol), 4-nitroimidazole (148.13 grams, 1.31 mol), potassium carbonate (362.11 grams, 2.62 mol), and dimethyl formamide (2500 mL) was stirred atambient temperature until complete as determined by hplc (16 hours). The reaction mixture was diluted with water (2000 mL) and extracted with ethyl acetate (4 x 500 mL). organic extracts were combined and washed with saturated sodium bicarbonate solution (2 x 500 mL), 10% citric acid 10 solution (2 x 500 mL), saturated sodium chloride solution (2 x 500 mL), dried using sodium sulfate, and evaporated. A portion (50 grams) of the crude product was purified by column chromatography on silica gel eluting with dichloromethane: heptane (16:3) gradient to dichloromethane 15 : heptane : methanol (16:3:0.2) giving 30.99 grams of ethyl 2-(2-naphthy1)-2-(4-nitroimidazoly1)acetate which was 90% pure by hplc. A 1 gram sample of the product was purified a second time using a Biotage Flash 40S system eluting with dichloromethane : heptane : methanol (16.9:3:0.1) to give 0.90 grams (46%) of ethyl 2-(2-naphthyl)-2-(4-120 nitroimidazolyl)acetate as a tan oil. MS (FIA) m/z 326.4 $[(M+H)^{+}]$. H nmr (CDCl₃): δ 1.25-1.31 (t, 3H), 4.28-4.39 (m, 2H), 6.16 (s, 1H), 7.36-7.44 (dd, 1H), 7.54-7.62 (m, 2H)3H), 7.84-7.90 (m, 3H), 7.90-7.95 (m, 2H).

25

684

Preparation 3

Ethyl 2-[4-((2R)-2-{2-[(Tert-butoxy)carbonylamino]-2-methyl propanoylamino}-3-indol-3-ylpropanoylamino)imidazolyl]-2-(2-naphthyl)acetate

5

10

15

20

A mixture of ethyl 2-(2-naphthyl)-2-(4-nitroimidazolyl) acetate (2.04 grams, 6.27 mmol), tetrahydrofuran (20 mL), and 10% palladium on carbon (2.04 gram) was hydrogenated at ambient temperature and pressure until complete as determined by hplc (20 hours). The catalyst was removed by filtration and rinsed with tetrahydrofuran (10 mL). The filtrate was added to a slurry consisting of 1-[3-(dimethyl amino)propyl-3-ethylcarbodiimide hydrochloride (1.20 grams, 6.27 mmol), tetrahydrofuran (10 mL), and (2R)-2-{2-[(tertbutoxy)carbonylamino]-2-methylpropanoylamino}-3-indol-3ylpropanoic acid (2.44 grams, 6.27 mmol) and stirred 16 hours at ambient temperature. The reaction mixture was partitioned between water (150 mL) and ethyl acetate (3 \times 50 mL). The organic extracts were combined, washed with saturated sodium chloride solution, dried using sodium sulfate, and evaporated. The resulting crude oil was purified by column chromatography on silica gel with hexane : ethyl acetate : methanol (10:10:1) as an eluent giving 1.72 grams (41%) of ethyl $2-[4-((2R)-2-\{2-[(tert-butoxy)]$ carbonylamino]-2-methylpropanoylamino}-3-indol-3-ylpropanoyl

685

amino)imidazolyl]-2-(2-naphthyl)acetate. A 0.2 gram sample was further purified using preparative reverse phase hplc to give 0.16 grams of ethyl 2-[4-((2R)-2-{2-[(tert-butoxy) carbonylamino]-2-methylpropanoylamino}-3-indol-3-ylpropanoyl amino)imidazolyl]-2-(2-naphthyl)acetate for analytical study. MS (FIA) m/z 667.4 [(M+H) $^+$]. Anal. calcd. exact mass for C₃₇H₄₃N₆O₆ [(M+H) $^+$] = 667.3244. Exact mass found by mass spectrometry: C₃₇H₄₃N₆O₆ [(M+H) $^+$] = 667.3254. ¹H nmr (CDCl₃): 1.25-1.42 (m, 19H), 3.24-3.33 (m, 2H), 4.28-4.33 (m, 2H), 4.98-5.01 (m, 1H), 5.94 (s, 1H), 6.85-7.01 (m, 3H), 7.18-7.21 (m, 2H), 7.35-7.39 (m, 2H), 7.49-7.58 (m, 4H), 7.78-7.84 (m, 4H), 8.69 (s, 1H), 10.65 (s, broad, 1H).

15

10

Preparation 4

Ethyl 2-[4-((2R)-2-{2-[(Tert-butoxy)carbonylamino]-2-methyl propanoylamino}-3-indol-3-ylpropanoylamino)imidazolyl]-2-phenylacetate

20

25

This compound was obtained from the reduction of ethyl 2-(4-nitroimidazolyl)-2-phenylacetate and subsequent reaction with (2R)-2-{2-[(tert-butoxy) carbonylamino]-2-methyl propanoylamino}-3-indole-3-ylpropanoic acid as a yellow foam in 73% yield after purification by flash chromatography using dichloromethane: methanol (19:1) as

686

the eluent. MS (FIA) m/z 617.5 [(M+H) $^{+}$]. 1 H nmr (CDCl $_{3}$): δ 1.19-1.32 (m, 18H), 3.10-3.12 (m, 1H), 3.16-3.17 (m, 1H), 3.32 (s, 1H), 4.22-4.27 (m, 2H), 4.69 (s, broad, 1H), 6.44 (s, 1H), 6.85-6.91 (m, 2H), 7.00 (t, 1H), 7.07-7.08 (m, 1H), 7.38-7.40 (m, 1H), 7.42-7.45 (m, 6H), 7.55-7.56 (m, 2H), 10.16 (s, broad, 1H), 10.75 (s, 1H).

Preparation 5

Ethyl 2-[4-((2R)-2-{2-[(Tert-butoxy)carbonylamino]-2methylpropanoylamino)-3-cyclohexylpropanoylamino) imidazolyl]-2-phenylacetate

5

This compound was obtained from the reduction of ethyl 2-(4-nitroimidazolyl)-2-phenylacetate and subsequent reaction with (2R)-2-{2-[(tert-butoxy) carbonylamino]-2-methyl propanoylamino}-3-cyclohexylpropanoic acid as a yellow foam in 45% yield after after purification by flash chromatography using dichloromethane: methanol (29:1) as the eluent. ¹H nmr (CDCl₃): δ 0.77-0.99 (m, 4H), 1.00-1.21 (m, 5H), 1.22-1.27 (m, 7H), 1.33-1.45 (m, 8H), 1.48-1.74 (m, 6H), 4.19-4.30 (m, 2H), 4.70-4.75 (m, 1H), 5.10 (s, broad, 1H), 5.84-5.86 (t, 1H), 6.92 (s, broad, 1H), 7.25-7.41 (m, 7H), 7.51 (m, 1H), 10.5 (s, broad, 1H).

687

Preparation 6

Ethyl 2-[4-((2R)-2-{2-[(Tert-butoxy)carbonylamino]-2-methylpropanoylamino)pentanoylamino)imidazolyl]-2-phenylacetate

5

This compound was obtained from the reduction of ethyl 2-(4-nitroimidazolyl)-2-phenylacetate and subsequent reaction with (2R)-2-{2-[(tert-butoxy)carbonylamino]-2-methyl propanoylamino}pentanoic acid as a yellow foam in 61% yield after purification using preparative reverse phase hplc. MS (FD+) m/z 529 (M*). ¹H nmr (CDCl₃): δ 0.82-0.84 (m, 3H), 1.25-1.46 (m, 20H), 1.60-1.75 (m, 1H), 1.77-1.83 (m, 1H), 4.24-4.32 (m, 2H), 4.76-4.78 (m, 1H), 5.25 (s, broad, 1H), 5.93-5.96 (m, 1H), 7.15-7.30 (m, 1H), 7.36-7.39 (m, 5H), 7.48-7.49 (m, 1H), 7.61-7.64 (d, 1H), 11.10-11.20 (m, 1H).

688

Preparation 7

Ethyl 2-[4-((2R)-2-{2-[(Tert-butoxy)carbonylamino]-2-methylpropanoylamino)hexanoylamino)imidazolyl]-2-phenylacetate

5

This compound was obtained from the reduction of ethyl 2-(4-nitroimidazolyl)-2-phenylacetate and subsequent

10 reaction with (2R)-2-{2-[(tert-butoxy) carbonylamino]-2-methyl propanoylamino}hexanoic acid as a yellow foam in 74% yield after purification by flash chromatography using dichloromethane: methanol (19:1) as the eluent. H nmr (CDCl₃): δ 0.87-0.88 (d, 3H), 1.28-1.32 (m, 7H), 1.44-1.46 (m, 8H), 1.51 (s, 6H), 1.66-1.68 (m, 3H), 1.95 (s, broad, 1H), 4.28-4.34 (m, 2H), 4.60-4.62 (m, 1H), 5.05 (s, broad, 1H), 5.85 (s, 1H), 6.90-7.00 (m, 1H), 7.36-7.46 (m, 6H), 9.65 (s, broad, 1H).

689

Pr paration 8

5

This compound was obtained from the reduction of ethyl 2-(4-nitroimidazolyl)-2-phenylacetate and subsequent reaction with (2R)-2-{2-[(tert-butoxy)carbonylamino]-2-10 methyl propanovlamino}-3-(3-thienyl)propanoic acid as a yellow foam in 53% yield after purification using preparative reverse phase hplc. MS (FD+) m/z 583 (M⁺). Anal. calcd. for $C_{29}H_{37}N_5O_6\cdot 1/2H_2O$: C: 58.77; H: 11.82. Found: C: 58.83; H: 6.03; N: 11.83. ¹H nmr 15 $(CDCl_3): \delta 1.20-1.23 (m, 3H), 1.33 (s, 9H), 1.40 (s, 6H),$ 3.29-3.30 (d, 2H), 4.18-4.27 (m, 2H), 5.04 (s, broad, 1H), 5.40 (s, broad, 1H), 5.94 (s, broad, 1H), 6.72-6.76 (m, 2H), 6.96-6.98 (t, 1H), 7.34 (s, 5H), 7.48 (s, 2H), 7.54-7.62 (m, 20 1H), 11.38 (m, 1H).

690

Preparation 9

Ethyl 2-[4-((2R)-3-Benzo[b]thiophen-3-yl-2-{2-[(tert-butoxy) carbonylamino]-2-methylpropanoylamino}propanoylamino) imidazolyl]-2-phenylacetate

5

This compound was obtained from the reduction of ethyl 2-(4-nitroimidazolyl)-2-phenylacetate and subsequent 10 reaction with (2R)-3-benzo[b]thiophen-3-y1-2-{2-[(tertbutoxy) carbonylamino]-2-methylpropanoylamino)propanoic acid as a yellow foam in 82% yield. H nmr (CDCl₃): δ 1.26-1.46 (m, 18H), 3.34-3.35 (m, 1H), 3.55-3.56 (m, 1H), 4.26-4.33 (m, 2H), 5.15 (s, broad, 2H), 5.82-5.84 (t, 1H), 7.20-7.47 (m, 10H), 7.60-7.65 (m 0.5H), 7.72-7.73 (m, 1H), 8.11-15 8.12 (m, 0.5H), 11.10 (m, broad, 1H).: 13 C nmr (CDCl₃): δ 14.46, 14.60, 19.34, 21.42, 25.69, 28.62, 36.64, 53.29, 54.39, 57.01, 60.77, 62.67, 64.05, 108.00, 122.26, 124.23, 124.34, 124.51, 125.58, 125.60, 125.87, 126.37, 127.92, 20 128.18, 128.24, 128.28, 128.35, 128.94, 129.58, 129.64, 129.68, 134.11, 134.53, 137.61, 140.55, 168.81, 168.86, 171.51, 174.81.

691

Preparation 10

2-[4-((2R)-2-{2-[(Tert-butoxy)carbonylamino]-2-methylpropanoylamino}-3-indol-3-ylpropanoylamino)
imidazolyl]-2-(2-naphthyl)acetic Acid

5

A solution consisting of ethyl 2-[4-((2R)-2-{2-[(tertbutoxy)carbonylamino]-2-methylpropanoylamino}-3-indol-3-10 ylpropanoylamino)imidazolyl]-2-(2-naphthyl)acetate (1.52 grams, 2.28 mmol), lithium hydroxide (0.11 grams, 4.56 mmol), dioxane (10 mL), and water (10 mL) was stirred at ambient temperature until complete as determined by hplc (30 The reaction mixture was concentrated to dryness and the residue was dissolved in water (20 mL). The aqueous 15 solution was adjusted to a pH of 3 using a 10% sodium bisulfate solution and extracted with ethyl acetate (3 \times 25 The organic layers were combined, dried using sodium sulfate, filtered, and concentrated to give 1.34 grams (92%) 20 of $2-[4-((2R)-2-\{2-[(tert-butoxy) carbonylamino]-2-methyl]$ propanoylamino}-3-indol-3-ylpropanoylamino)imidazolyl]-2-(2naphthyl)acetic acid.

692

Preparation 11

2-[4-((2R)-2-{2-[(Tert-butoxy)carbonylamino]-2-methyl propanoylamino}-3-cyclohexylpropanoylamino)imidazolyl]-2-phenylacetic Acid

5

This compound was obtained from the hydrolysis of ethyl $2-[4-((2R)-2-\{2-[(tert-butoxy)\ carbonylamino]-2-methyl\ propanoylamino\}-3-cyclohexylpropanoylamino)imidazolyl]-2- 10 phenylacetate as a white powder in 69% yield. ¹H nmr (DMSO-d₆): <math>\delta$ 0.64-0.92 (m, 3H), 0.93-1.15 (m, 3H), 1.16-1.41 (m, 7H), 1.42-1.73 (m, 4H), 4.36 (s, broad, 1H), 6.08 (s, 1H), 6.98 (s, 1H), 7.15 (s, 1H), 7.31 (s, 5H), 7.49 (s, 2H), 10.0 (d, broad, 1H).

15

693

Preparation 12

2-[4-((2R)-2-{2-[(Tert-butoxy)carbonylamino]-2-methyl propanoylamino}-3-(2-naphthyl)propanoylamino)imidazolyl]-2phenylacetic Acid

5

This compound was obtained from the reduction of ethyl 2-(4-nitroimidazolyl)-2-phenylacetate and subsequent

10 reaction with (2R)-2-{2-[(tert-butoxy)carbonylamino]-2-methyl propanoylamino}-3-(2-naphthyl)propanoic acid followed by hydrolysis to give a white solid in 76% yield. ¹H nmr (DMSO-d₆): δ 1.02-1.13 (t, 7H), 1.27 (s, 8H), 3.24 (m, 3H), 3.50 (m, 1H), 4.58-4.78 (m, 1H), 6.24 (s, 1H), 7.00 (s, broad, 1H), 7.13-7.56 (m, 9H), 7.73 (d, 1.5H), 7.84 (d, 1.5H), 8.16 (m, 1H), 9.98 (s, broad, 0.5H), 10.11 (s, broad, 0.5H), 13.51 (s, broad, 1H).

Preparation 13

20 N-[(1R)-2-Indol-3-yl-1-(N-{1-[2-(4-methylpiperidinyl)-1-(2-naphthyl)-2-oxoethyl]imidazol-4-yl}carbamoyl)ethyl]-2[(tert-butoxy)carbonylamino]-2-methylpropanamide

694

A solution consisting of $2-[4-((2R)-2-\{2-[(tert-butoxy)\}$ carbonylamino] -2-methylpropanoylamino} -3-indol-3-yl propanoylamino)imidazolyl]-2-(2-naphthyl)acetic acid (0.55 5 grams, 0.861 mmol), 4-methylpiperidine (0.085 grams, 0.861 mmol), 1,3-dicyclohexylcarbodiimide (0.195 grams, 0.947 mmol), 1-hydroxybenzotriazole hydrate (0.116 grams, 0.861 mmol) and dimethyl formamide (5 mL) was stirred at ambient 10 temperature until complete as determined by hplc (7 hours). The reaction mixture was diluted with water (40 mL) and extracted with ethyl acetate (4 x 25 mL). The organic extracts were combined, washed with saturated sodium chloride solution (2 x 35 mL), dried using sodium sulfate, and concentrated to an oil. The crude product was purified 15 using preparative reverse phase hplc to give 0.32 grams (52%) of N- $[(1R)-2-indol-3-yl-1-(N-{1-[2-(4-methyl)]}$ piperidyl) -1-(2-naphthyl) -2-oxoethyl]imidazol-4-yl} carbamoy1) ethy1]-2-[(tert-butoxy) carbonylamino]-2methylpropanamide. ¹H nmr (CDCl₃): 0.76-0.77 (d, 2H), 20 0.91-0.95 (m, 2H), 1.23-1.36 (m, 18H), 1.54 (m, 1H), 1.67 (m, 1H), 2.70-2.72 (m, 2H), 3.25-3.29 (m, 2H), 3.68 (m, 1H),4.55-4.70 (m, 1H), 4.98 (m, 1H), 6.24 (m, 1H), 6.81-6.83 (d, 1H), 6.92 (m, 1H), 7.00-7.01 (m, 1H), 7.18-7.28 (m, 3H), 7.37-7.55 (m, 5H), 7.76-7.83 (m, 4H), 8.80 (s, broad, 1H), 10.38 (s, broad, 1H). ¹³C nmr (CDCl₃): δ 14.60, 19.32, 19.47, 21.41, 21.83, 21.90, 25.39, 25.55, 26.04, 28.56,

695

28.63, 28.84, 31.05, 31.16, 31.21, 33.98, 34.08, 34.29, 34.69, 43.42, 46.28, 46.52, 49.38, 54.55, 56.99, 60.77, 62.31, 69.97, 71.02, 108.80, 110.24, 111.79, 119.02, 119.36, 121.86, 124.10, 125.99, 127.12, 127.36, 127.97, 128.08, 128.10, 128.16, 128.33, 128.63, 128.71, 129.77, 132.26, 133.63, 133.75, 134.02, 136.58, 137.29, 155.16, 157.65, 166.07, 166.18, 166.22, 166.34, 169.40, 171.52, 175.12.

Preparation 14

10 N-[(1R)-2-Indol-3-yl-1-(N- $\{1-[2-(4-methylpiperidy1)-2-oxo-1-phenylethyl]imidazol-4-yl\}carbamoyl)ethyl]-2-[(tert-butoxy) carbonylamino]-2-methylpropanamide$

15

20

25

This compound was obtained from the hydrolysis of ethyl $2-[4-((2R)-2-\{2-[(tert-butoxy)\ carbonylamino]-2-methyl\ propanoylamino\}-3-indol-3-ylpropanoylamino)imidazolyl]-2-phenylacetate and subsequent reaction with 4-methyl piperidine in 84% yield after Biotage Flash 40M purification using dichloromethane: methanol (24:1) as the eluent. MS (FIA) m/z 670.5 [(M+H)<math>^+$]. ¹H nmr (CDCl $_3$): δ 0.74-0.75 (d, 2H), 0.89-0.90 (d, 2H), 1.17-1.32(m, 18H), 1.53-1.63 (m, 3H), 2.66-2.70 (m, 1H), 3.05 (t, 1H), 3.15-3.20 (m, 1H), 3.69-3.83 (m, 1H), 4.36-4.49 (m, 1H), 4.67 (s, broad, 1H), 6.90-6.93 (m, 2H), 7.01-7.04 (m, 2H), 7.11 (s, 1H), 7.26-

696

7.32 (m, 2H), 7.40-7.54 (m, 5H), 7.67 (s, broad, 1H), 8.16 (m, broad, 1H), 10.49 (s, broad, 1H), 10.84 (s, 1H).

Preparation 15

pyrrolidinylethyl)imidazol-4-yl]carbamoyl)ethyl)-2-[(tertbutoxy)carbonylamino]-2-methylpropanamide

10

20

5

This compound was obtained from the hydrolysis of ethyl $2-[4-((2R)-2-\{2-[(tert-butoxy) carbonylamino]-2-methyl]$ propanoylamino}-3-indol-3-ylpropanoylamino)imidazolyl]-2phenylacetate and subsequent reaction with pyrrolidine in 80% yield after purification by flash chromatography using 15 dichloromethane: methanol (19:1) as the eluent. 1H nmr $(CDCl_3): \delta 1.10-1.40 \text{ (m, 15H)}, 1.67-1.92 \text{ (m, 3H)}, 2.92-$ 3.60 (m, 5H), 4.90 (s, broad, 1H), 5.33 (s, broad, 1H), 5.85 (d, 1H), 6.80-7.05 (m, 3H), 7.13-7.39 (m, 10H), 7.44-7.80 (m, 2H), 8.96 (s, broad, 1H), 10.20 (s, broad, 1H). ¹³C nmr $(CDC1_3): \delta 14.25, 21.11, 24.02, 25.63, 26.08, 28.24,$ 33.87, 46.39, 46.64, 54.28, 56.67, 60.46, 63.07, 63.09, 108.33, 109.73, 110.69, 111.47, 118.36, 118.56, 119.05, 121.57, 123.77, 125.01, 126.42, 127.60, 128.51, 129.38, 25 133.14, 133.85, 136.23, 136.45, 136.49, 165.79, 165.85, 169.17, 174.87.

697

Preparation 16

 $N-((1R)-2-Cyclohexyl-1-\{N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]carbamoyl\}ethyl)-2-[(tert-butoxy)carbonylamino]-2-methylpropanamide$

5

This compound was obtained from the reaction of 2-[4-((2R)-2-{2-[(tert-butoxy) carbonylamino]-2-

10 methylpropanoylamino}-3-cyclohexylpropanoylamino)
 imidazolyl]-2-phenylacetic acid and pyrrolidine in 99% yield
 after purification by flash chromatography using
 dichloromethane : methanol (19:1) as the eluent. ¹H nmr
 (CDCl₃): δ 0.71-1.00 (m, 2H), 1.00-1.16 (m, 2H), 1.17-2.18
15 (m, 28H), 3.07-3.20 (s, broad, 1H), 3.38-3.64 (m, 3H), 4.624.77 (m, 1H), 5.91 (t, 1H), 7.38 (m, 7H), 9.93 (s, broad, 1H), 10.60 (s, broad, 1H).

698

Preparation 17

Methyl 1-{2-[4-((2R)-2-{2-[(Tert-butoxy)carbonylamino]-2-methylpropanoylamino}-3-cyclohexylpropanoylamino) imidazolyl]-2-phenylacetyl}(2S)pyrrolidine-2-carboxylate

5

This compound was obtained from the reaction of 2-[4-((2R)-2-{2-[(tert-butoxy) carbonylamino]-2-methylpropanoylamino}-3-cyclohexylpropanoylamino)

10 imidazolyl]-2-phenylacetic acid and L-proline methyl ester in 88% yield after purification by flash chromatography using dichloromethane: methanol (19:1) as the eluent. ¹H nmr (CDCl₃): δ 0.75-0.98 (m, 4H), 0.98-1.21 (m, 4H), 1.21-1.50 (m, 12H), 1.50-1.80 (m, 5H), 1.80-2.07 (m, 3H), 2.07-1.228 (m, 1H), 3.19-3.36 (m, 1H), 3.37-3.72 (m, 2H), 3.72 (s, 3H), 4.03-4.61 (m, 1H), 4.61-4.75 (m, 1H), 5.11 (s, broad, 1H), 5.92-6.00 (m, 1H), 7.00 (s, broad, 1H), 7.21-7.49 (m, 7H), 8.00 (s, 1H), 9.60 (s, broad, 1H).

699

Preparation 18

 $(2R)-2-\{2-[(Tert-butoxy) carbonylamino]-2-methylpropanoyl$ amino}-N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]pentanamide

5

This compound was obtained from the hydrolysis of ethyl $2-[4-((2R)-2-\{2-[(tert-butoxy) carbonylamino]-2-methyl]$ propanoylamino)pentanoylamino)imidazoly1]-2-phenylacetate 10 and subsequent reaction with pyrrolidine in 87% yield after purification by preparative reverse phase hplc. 1H nmr $(CDCl_3): \delta 0.72-0.75 \text{ (m, 3H)}, 1.18-1.41 \text{ (m, 17H)}, 1.75-$ 1.79 (m, 4H), 1.80-1.92 (m, 1H), 3.00 (s, broad, 1H), 3.09- $3.11 \, (m, 1H), 3.44-3.52 \, (m, 3H), 4.74 \, (m, 1H), 5.45 \, (s, 1H)$ 15 broad, 1H), 6.09-6.10 (d, 1H), 7.28-7.52 (m, 8H), 10.98-11.07 (m, 1H). ¹³C nmr (CDCl₃): δ 14.14, 14.54, 18.95, 21.36, 24.35, 24.88, 26.43, 26.49, 26.54, 28.65, 35.62, 35.68, 46.72, 46.86, 53.55, 56.85, 60.70, 63.31, 108.46, 20 128.86, 129.46, 129.59, 133.70, 134.69, 134.74, 137.31, 166.20, 166.26, 170.06, 171.46, 175.02.

700

Preparation 19

 $(2R)-2-\{2-[(Tert-butoxy) carbonylamino]-2-methylpropanoylamino\}-N-\{1-[2-(4-methylpiperidyl)-2-oxo-1-phenylethyl]$ imidazol-4-yl}pentanamide

5

This compound was obtained from the hydrolysis of ethyl $2-[4-((2R)-2-\{2-[(tert-butoxy) carbonylamino]-2-methyl]$ propanoylamino)pentanoylamino)imidazolyl]-2-phenylacetate 10 and subsequent reaction with 4-methylpiperidine in 72% yield after purification by preparative reverse phase hplc. 1H nmr (CDCl₃): δ 0.30-0.40 (m, 0.5H), 0.77-0.80 (m, 4H), 0.91-0.92 (d, 1H), 1.05-1.20 (m, 0.5H), 1.23-1.56 (m, 23H), 1.62-1.67 (m, 1H), 1.76 (s, broad, 1H), 2.59-2.72 (m, 1.5H), 15 3.00 (t, 0.5H), 3.71 (m, 1H), 4.53-4.61 (dd, 1H), 4.73 (s, 1.73)broad, 1H), 5.40 (s, broad, 1H), 6.26-6.32 (m, 1H), 7.29-7.45 (m, 7H), 10.89-10.93 (m, 1H). 13 C nmr (CDCl₃): δ 14.15, 14.16, 14.56, 18.97, 21.82, 21.93, 24.90, 25.39, 26.01, 26.50, 26.55, 28.69, 31.12, 31.20, 33.93, 34.25, 20 34.70, 35.53, 35.64, 43.35, 46.31, 46.42, 53.63, 53.85, 56.90, 60.74, 62.23, 62.32, 108.61, 128.65, 128.93, 129.47, 129.60, 129.70, 134.01, 134.97, 137.16, 137.24, 165.99, 166.06, 166.12, 166.21, 170.04, 175.05.

25

701

Preparation 20

(2R)-2-{2-[(Tert-butoxy)carbonylamino]-2-methylpropanoyl amino}-N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]hexanamide

5

This compound was obtained from the hydrolysis of ethyl $2-[4-((2R)-2-\{2-[(tert-butoxy) carbonylamino]-2-methyl]$ propanoylamino)hexanoylamino)imidazolyl]-2-phenylacetate and 10 subsequent reaction with pyrrolidine in 63% yield after purification by preparative reverse phase hplc. 1H nmr (CDCl₃): δ 0.68-0.71 (m, 3H), 1.20-1.22 (m, 5H), 1.33 (s, 11H), 1.41 (s, 3H), 1.68 (m, 1H), 1.74-1.78 (m, 4H), 1.86-15 1.88 (m, 1H), 3.08-3.09 (m, 1H), 3.44-3.51 (m, 3H), 4.74 (s, broad, 1H), 5.50 (s, broad, 1H), 6.09-6.10 (d, 1H), 7.27-7.33 (m, 3H), 7.37-7.39 (m, 2H), 7.45-7.57 (m, 3H), 11.05-11.06 (m, 1H). ¹³C nmr (CDCl₃): δ 14.11, 19.35, 22.73, 24.35, 26.42, 26.46, 26.51, 27.65, 28.68, 31.15, 33.22, 33.30, 46.72, 46.84, 53.72, 56.84, 63.27, 69.71, 108.47, 20 128.84, 129.43, 129.57, 133.68, 133.76, 134.74, 134.80, 137.30, 166.19, 166.26, 170.11, 175.01.

702

Preparation 21

 $(2R)-2-\{2-[(Tert-butoxy) carbonylamino]-2-methylpropanoylamino}-N-\{1-[2-(4-methylpiperidyl)-2-oxo-1-phenylethyl]$ $imidazol-4-yl\}hexanamide$

5

This compound was obtained from the hydrolysis of ethyl $2-[4-((2R)-2-\{2-[(tert-butoxy) carbonylamino]-2-methyl]$ propanoylamino)hexanoylamino)imidazolyl]-2-phenylacetate and 10 subsequent reaction with 4-methylpiperidine in 66% yield after purification by preparative reverse phase hplc. ¹H nmr (CDCl₃): δ 0.73-0.74 (m, 3H), 0.89-0.91 (m, 1H), 1.21-1.66 (m, 28H), 1.75-1.87 (m, 1H), 2.57-2.63 (q, 1H), 2.72 (t, 0.5H), 2.99 (t, 0.5H), 3.70-3.72 (m, 1H), 4.53-4.61 (dd, 15 1H), 4.73 (s, broad, 1H), 5.43 (s, broad, 1H), 6.28-6.31 (d, 1H), 7.33-7.50 (m, 7H), 11.01-11.03 (m, 1H). 13 C nmr $(CDCl_3)$: δ 14.15, 14.55, 19.34, 21.39, 21.83, 21.93, 22.76, 25.40, 26.49, 26.56, 27.63, 28.70, 28.73, 31.12, 31.21, 33.24, 33.33, 33.94, 34.24, 34.70, 43.32, 46.30, 20 46.41, 53.72, 56.88, 60.72, 62.10, 62.17, 62.26, 108.57, 128.64, 128.92, 129.42, 129.56, 129.67, 133.84, 135.00, 135.04, 135.07, 137.20, 137.25, 137.28, 137.32, 165.97, 166.02, 166.12, 166.18, 170.00, 174.95.

25

703

Preparation 22

N-[(1R)-1-(N-{1-[2-(4-Methylpiperidy1)-2-oxo-1-phenylethy1] imidazol-4-yl}carbamoy1)-2-(3-thienyl)ethyl]-2-[(tertbutoxy)carbonylamino]-2-methylpropanamide

5

This compound was obtained from the hydrolysis of ethyl $2-[4-((2R)-2-\{2-[(tert-butoxy) carbonylamino]-2-methyl]$ 10 propanoylamino}-3-(3-thienyl)propanoylamino)imidazolyl]-2phenylacetate and subsequent reaction with 4-methyl piperidine in 70% yield after purification by preparative reverse phase hplc. ¹H nmr (CDCl₃): δ 0.79-0.80 (d, 2H), 0.91-0.92 (m, 2H), 0.93-1.15 (m, 3H), 1.32-1.40 (m, 15H), 1.45-1.55 (m, 1H), 1.64 (m, 1H), 2.60-2.62 (m, 1H), 2.75 (t, 15 0.5H), 2.98 (t, 0.5H), 3.32-3.35 (d, 2H), 3.70 (m, 1H), 4.50-4.67 (dd, 1H), 5.00 (m, 1H), 5.50 (s, broad, 1H), 6.23-6.24 (m, 1H), 6.73-6.76 (m, 2H), 6.97-6.98 (m, 1H), 7.32-7.45 (m, 6H), 7.56 (s, broad, 1H), 10.94 (s, broad, 1H). ¹³C nmr (CDCl₃): δ 14.58, 19.45, 21.40, 21.95, 25.58, 20 25.62, 25.69, 28.75, 31.12, 33.15, 33.94, 33.98, 34.26, 43.36, 46.32, 46.45, 49.28, 54.74, 56.97, 60.74, 62.17, 62.28, 69.84, 108.79, 124.54, 126.95, 127.23, 128.60, 128.93, 129.60, 129.70, 134.03, 135.01, 135.14, 137.05, 137.19, 138.80, 157.69, 165.97, 168.51, 171.49, 175.14. 25

704

Preparation 23

N-((1R)-1-{N-[1-(2-Oxo-1-phenyl-2-pyrrolidinylethyl) imidazol-4-yl]carbamoyl}-2-(3-thienyl)ethyl)-2-[(tertbutoxy)carbonylamino]-2-methylpropanamide

5

This compound was obtained from the hydrolysis of ethyl $2-[4-((2R)-2-\{2-[(tert-butoxy) carbonylamino]-2-methyl]$ 10 propanoylamino}-3-(3-thienyl)propanoylamino)imidazolyl]-2phenylacetate and subsequent reaction with pyrrolidine in 71% yield after purification by preparative reverse phase hplc. ¹H nmr (CDCl₃): δ 1.32-1.39 (m, 15H), 1.77-1.81 (m, 3H), 1.90 (m, 1H), 3.13 (m, broad, 1H), 3.35 (s, 2H), 3.48-3.54 (m, 3H), 4.97-5.08 (m, 1H), 5.50 (s, broad, 1H), 6.03-15 6.05 (m, 1H), 6.73-6.76 (m, 2H), 6.98-6.99 (m, 1H), 7.34-7.40 (m, 5H), 7.45-7.48 (m, 2H), 10.91 (s, broad, 1H). 13 C nmr (CDCl₃): δ 14.58, 19.37, 21.41, 24.39, 25.55, 25.60, 25.66, 26.46, 28.73, 31.19, 33.15, 46.76, 46.91, 54.73, 56.98, 60.74, 63.36, 69.84, 108.56, 108.63, 124.56, 126.94, 20 127.24, 128.79, 128.82, 128.87, 129.53, 129.65, 133.82, 134.70, 134.83, 134.87, 137.14, 137.19, 137.24, 138.79, 166.11, 168.52, 171.50, 175.08.

705

Preparation 24

N-((1R)-2-Benzo[b]thiophen-3-yl-1-{N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]carbamoyl}ethyl)-2-[(tert-butoxy)carbonylamino]-2-methylpropanamide

5

This compound was obtained from the hydrolysis of ethyl $2-[4-((2R)-3-benzo[b]-thiophen-3-yl-2-{2-[(tert-butoxy)]}$ 10 carbonylamino] -2-methylpropanoylamino) propanoylamino) imidazolyl]-2-phenylacetate and subsequent reaction with pyrrolidine in 68% yield after purification by preparative reverse phase hplc. ¹H nmr (CDCl₃): δ 1.21-1.34 (m, 15H), 1.76-1.87 (m, 3H), 3.00-3.10 (m, 1H), 3.34-3.38 (m, 2H), 3.48-3.57 (m, 3H), 5.17 (s, 1H), 5.40 (s, broad, 1H), 5.92 15 (s, 1H), 7.16-7.30 (m, 1H), 7.34 (s, 10H), 7.39-7.46 (m, 1H)1H), 7.57-7.59 (m, 0.5H), 7.68-7.69 (m, 1H), 8.06-8.07 (d, 0.5H), 10.85 (s, broad, 1H). ¹³C nmr (CDCl₃): δ 14.58, 19.38, 21.41, 24.38, 25.54, 25.64, 25.70, 26.03, 26.43, 28.58, 31.20, 34.26, 36.36, 46.70, 46.90, 49.27, 53.40, 20 54.44, 56.93, 60.75, 63.23, 69.80, 108.47, 108.61, 122.87, 124.29, 124.35, 124.44, 125.64, 125.84, 126.45, 127.78, 128.17, 128.75, 128.79, 128.83, 129.51, 129.61, 132.48, 133.42, 134.04, 134.78, 137.14, 139.34, 140.45, 157.71, 25 166.11, 168.93, 171.51, 175.01, 175.13.

706

Preparation 25

N-[(1R)-2-Benzo[b]thiophen-3-yl-1-(N-{1-[2-(4-methyl piperidyl)-2-oxo-1-phenylethyl]imidazol-4-yl}carbamoyl) ethyl]-2-[(tert-butoxy)carbonylamino]-2-methylpropanamide

5

This compound was obtained from the hydrolysis of ethyl $2-[4-((2R)-3-benzo[b]-thiophen-3-yl-2-{2-[(tert-butoxy)]}$ 10 carbonylamino] -2-methylpropanoylamino) propanoylamino) imidazolyl]-2-phenylacetate and subsequent reaction with 4methylpiperidine in 71% yield after purification by preparative reverse phase hplc. ¹H nmr (CDCl₃): δ 0.78-0.79 (d, 1H), 0.90-1.00 (d, 1H), 1.12-1.33; (m, 20H), 1.51 15 (m, 1H), 1.64 (m, 1H), 2.61-2.67 (m, 1.5H), 2.90 (m, 1H), 3.32-3.33 (m, 1H), 3.51-3.62 (m, 2.5H), 4.57-4.68 (m, 1H), 5.45 (s, 1H), 6.11-6.14 (s, broad, 1H), 7.19-7.45 (m, 10H), 7.57-7.59 (m, 0.5H), 7.67-7.69 (m, 1H), 8.06 (s, broad, 0.5H), 11.00 (s, broad, 1H). ¹³C nmr (CDCl₃): δ 14.59, 19.41, 21.41, 21.86, 21.88, 21.98, 25.45, 25.69, 25.79, 20 26.04, 28.71, 31.15, 32.10, 33.97, 34.26, 34.68, 36.40, 43.27, 46.27, 46.40, 49.24, 53.42, 54.44, 56.90, 60.75, 62.13, 108.68, 108.81, 122.87, 124.28, 124.35, 124.41, 125.64, 125.84, 126.45, 127.78, 128.20, 128.61, 128.85, 25 129.47, 129.57, 129.66, 132.50, 133.40, 134.03, 135.05,

707

135.15, 139.38, 140.44, 157.75, 165.86, 166.05, 169.02, 171.50, 174.98, 175.14.

Example 1

5 N-[(1R)-2-Indol-3-yl-1-(N-{1-[2-(4-methylpiperidyl)-1-(2-naphthyl)-2-oxoethyl]imidazol-4-yl}carbamoyl)ethyl]-2-amino-2-methylpropanamide Dihydrochloride

10

15

20

25

A solution consisting of $N-((1R)-2-indol-3-yl-1-(N-\{1-indol-3-yl-1-i$ [2-(4-methylpiperidyl)-1-(2-naphthyl)-2-oxoethyl]imidazol-4y1}carbamoy1)ethy1]-2-[(tert-butoxy)carbonylamino]-2-methyl propanamide (0.32 grams, 0.445 mmol) and anisole (0.25 mL) dissolved in methylene chloride (20 mL) was added trifluoroacetic acid (2.5 mL). The resulting reaction mixture was stirred at ambient temperature until complete as determined by hplc (2.5 hours). The reaction mixture was concentrated to dryness. The residue was dissolved in methanol (5 mL) and applied to a Varian Mega Bond Elut SCX ion exchange column (5 gram). The column was washed with methanol (50 mL). The product was eluted from the column with 2N ammonia in methanol (30 mL). The eluent was concentrated to dryness to give the free base (0.28 grams). A 1.95 M solution of anhydrous HCl in ethyl acetate (0.456 mL, 0.89 mmol) was added to the free base which was dissolved in ethyl acetate (10 mL). The resulting

708

precipitate was collected by filtration and dried in vacuum (4-methylpiperidyl)-1-(2-naphthyl)-2-oxoethyl]imidazol-4y1}carbamoy1)ethy1]-2-amino]-2-methy1propanamide dihydrochloride. MS (FIA) m/z 620.7 [(M+H)⁺]. Anal. calcd. for $C_{36}H_{41}N_7O_3\cdot 2HC1\cdot 1/2H_2O$: C: 61.62; H: 6.32; N: 13.97. Found: C: 61.42; H: 6.18; N: 13.62. Anal. calcd. exact mass for $C_{36}H_{42}N_7O_3$ [(M+H)⁺] = 620.3349. Exact mass found by mass spectrometry: $C_{36}H_{42}N_7O_3$ [(M+H)⁺] = 620.3355. 10 1 H nmr (DMSO-d₆): 0.65-0.67 (d, 2H), 0.89-0.90 (d, 2H), 1.16-1.24 (m, 2H), 1.35-1.36 (d, 4H), 1.51-1.53 (d, 4H), 1.63-1.65 (m, 1H), 2.68-2.74 (m, 1.5H), 3.08 (t, 0.5H), 3.17-3.19 (m, 1H), 3.26-3.27 (m, 1H), 3.71-3.82 (m, 1H), 4.40-4.55 (m, 1H), 4.71-4.72 (t, 1H), 6.90-7.00 (m, 1H), 15 7.02-7.04 (m, 1H), 7.26-7.33 (m, 3H), 7.52 (m, 1H), 7.59-

20 **Example 2**

11.35 (s, 1H).

 $N-[(1R)-2-Indol-3-yl-1-(N-\{1-[2-(4-methylpiperidyl)-2-oxo-1-phenylethyl]imidazol-4-yl)carbamoyl)ethyl]-2-amino-2-methyl propanamide Dihydrochloride$

7.62 (m, 3H), 7.74 (m, 1H), 7.98-8.09 (m, 4H), 8.31-8.32 (d,

3H), 8.49-8.61 (m, 1H), 8.66-8.68 (d, 1H), 10.94 (s, 1H),

709

This compound was obtained from N- $\{(1R)-2-indol-3-yl-1-(N-\{1-[2-(4-methylpiperidyl)-2-oxo-1-phenylethyl]imidazol-4-yl\}carbamoyl)ethyl]-2-<math>\{(tert-butoxy) carbonylamino\}-2-methyl$ propanamide as a red foam in 100% yield. MS (FIA) m/z 570.5 $[(M+H)^*]. \quad ^1H \text{ nmr } (d-MeOH): \quad \delta \quad 0.81-0.82 \quad (d, 2H), \quad 0.98-0.99 \quad (d, 2H), \quad 1.18-1.21 \quad (m, 2H), \quad 1.34-1.37 \quad (m, 1H), \quad 1.43 \quad (s, 3H), \quad 1.61 \quad (s, 6H), \quad 1.71 \quad (t, 1H), \quad 2.73-2.76 \quad (m, 1.5H), \quad 3.14 \quad (t, 0.5H), \quad 3.27-3.33 \quad (m, 1H), \quad 3.40-3.44 \quad (m, 1H), \quad 3.61-3.65 \quad (m, 1H), \quad 3.75-3.77 \quad (d, 1H), \quad 4.45-4.60 \quad (m, 1H), \quad 4.81 \quad (s, broad, 4H), \quad 6.94-6.99 \quad (m, 1.5H), \quad 7.06-7.07 \quad (m, 1.5H), \quad 7.19 \quad (s, 1H), \quad 7.31-7.35 \quad (m, 2H), \quad 7.52-7.61 \quad (m, 6H), \quad 8.62-8.65 \quad (d, 1H).$

5

10

Example 3

N-((1R)-2-Indol-3-yl-1-{N-[1-(2-oxo-1-phenyl-2-pyrrolidinyl ethyl)imidazol-4-yl]carbamoyl}ethyl)-2-amino-2-methyl propanamide Bistrifluoroacetic Acid

This compound was obtained from N-((1R)-2-indol-3-yl-1-{N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]carbamoyl}ethyl)-2-[(tert-butoxy)carbonylamino]-2-methylpropanamide as a white solid in 50% yield. MS (FD+)m/z 541 (M*). Anal. calcd. for C₃₀H₃₅N₇O₃·2C₂HF₃O₂: C: 53.06;

H: 4.85; N: 12.74. Found: C: 52.93; H: 4.88; N: 12.55. ¹H nmr (DMSO-d₆): δ 1.29 (s, 3H), 1.46-1.48 (d,

710

3H), 1.72-1.88 (m, 4H), 2.94 (m, 1H), 3.06-3.07 (m, 1H), 3.19-3.20 (m, 1H), 3.40-3.41 (d, 2H), 3.67-3.69 (m, 1H), 4.78 (s, broad, 1H), 6.53 (s, 1H), 6.93-6.97 (m, 1H), 7.06 (m, 1H), 7.20 (d, 1H), 7.31-7.36 (m, 2H), 7.42-7.42 (m, 4H), 7.73-7.80 (m, 2H), 8.01 (s, broad, 2H), 8.36-8.38 (d, 1H), 10.82-10.85 (d, 2H).

Example 4

N-((1R)-2-Cyclohexyl-1-{N-[1-(2-oxo-1-phenyl-2-pyrrolidinyl ethyl)imidazol-4-yl]carbamoyl}ethyl)-2-amino-2-methyl propanamide Bistrifluoroacetic Acid

This compound was obtained from N-((1R)-2-cyclohexyl-1-{N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl] carbamoyl}ethyl)-2-[(tert-butoxy)carbonylamino]-2-methyl propanamide as a white solid in 81% yield. MS (FD+) m/z 508.2 (M*). Anal. calcd. for C₂₈H₄₀N₆O₃·2C₂HF₃O₂: C: 52.17; H: 5.75; N: 11.41. Found: C: 52.11; H: 5.81; N: 11.40. ¹H nmr (DMSO-d₆): δ 0.91 (m, 2H), 1.11-1.13 (m, 3H), 1.31 (s, broad, 1H), 1.48-1.79 (m, 18H), 2.91-2.93 (m, 1H), 3.38 (m, 2H), 3.65-3.68 (m, 1H), 4.53-4.54 (m, 1H), 6.52 (s, 1H), 7.30 (s, 1H), 7.41-7.48 (m, 5H), 7.77-7.81 (d, 25 1H), 8.13-8.14 (d, 3H), 8.30-8.31 (d, 1H), 10.68 (s, 1H).

711

Example 5

Methyl 1-(2-{4-[(2R)-2-(2-Amino-2-methylpropanoylamino)-3-cyclohexylpropanoylamino]imidazolyl}-2-phenylacetyl)
(2S)pyrrolidine-2-carboxylate Bistrifluoroacetic Acid

5

25

This compound was obtained from methyl $1-\{2-[4-((2R)-2-$ {2-[(tert-butoxy) carbonylamino]-2-methylpropanoylamino}-3cyclohexylpropanoylamino)imidazolyl]-2-phenylacetyl}(2S) 10 pyrrolidine-2-carboxylate as an off white solid in 53% yield. MS (FD+) m/z 566.2 (M⁺). Anal. calcd. for 5.58; N: $C_{30}H_{42}N_6O_5 \cdot 2C_2HF_3O_2$: C: 51.38; H: 10.57. Found: 10.66. ¹H nmr (DMSO-d₆): δ 0.91 51.44; H: 5.59; N: (m, 2H), 1.12-1.16 (m, 3H), 1.30 (s, broad, 1H), 1.47-1.67 15 (m, 13H), 1.81-1.87 (m, 3H), 2.10-2.30 (m, 1H), 2.95-3.15 (m, 1H), 3.64 (s, 1H), 3.68 (s, 2H), 3.72-3.75 (m, 1H),4.41-4.43 (t, 1H), 4.53-4.54 (m, 1H), 6.59-6.62 (m, 1H), 7.17-7.19 (d, 0.5H), 7.37-7.47 (m, 6.5H), 7.57-7.62 (d, 1H), 8.11 (s, broad, 3H), 8.25-8.28 (m, 1H), 10.55 (s, 1H). 20

Example 6

(2R)-2-(2-Amino-2-methylpropanoylamino)-N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]pentanamide

Bistrifluoroacetic Acid

712

This compound was obtained from (2R)-2-{2-[(tert-butoxy) carbonylamino]-2-methylpropanoylamino}-N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]pentanamide as an off white solid in 82% yield. MS (FD+) m/z 454 (M*). Anal. calcd. for C₂₄H₃₄N₆O₃·2C₂HF₃O₂: C: 49.27; H: 5.32; N: 12.31. Found: C: 49.19; H: 5.29; N: 12.25. ¹H nmr (DMSO-d₆): δ 0.88-0.90 (m, 3H), 1.20-1.30 (m, 1H), 1.30-1.39 (m, 1H), 1.50-1.51 (m, 6H), 1.68-1.85 (m, 7H), 2.89-2.90 (m, 1H), 3.40 (s, broad, 2H), 3.67 (s, broad, 1H), 4.41-4.42 (m, 1H), 6.59 (s, 1H), 7.35 (s, 1H), 7.44-7.46 (m, 5H), 7.96-8.00 (d, 1H), 8.19-8.20 (m, 3H), 8.33-8.35 (d, 1H), 10.85 (s, 1H).

713

Example 7

(2R)-2-(2-Amino-2-methylpropanoylamino)-N-{1-[2-(4-methylpiperidyl)-2-oxo-1-phenylethyl]imidazol-4-yl} pentanamide Bistrifluoroacetic Acid

5

This compound was obtained from (2R)-2-{2-[(tert-butoxy) carbonylamino}-2-methylpropanoylamino}-N-{1-[2-(4-10 methyl piperidyl)-2-oxo-1-phenylethyl]imidazol-4-yl}pentanamide as an off white solid in 85% yield. MS (FD+) m/z 482 (M[†]). Anal. calcd. for C₂₆H₃₈N₆O₃·2C₂HF₃O₂·1/2H₂O: C: 50.07; H: 5.74; N: 11.68. Found: C: 49.94; H: 5.53; N: 11.51. ¹H nmr (DMSO-d₆): δ 0.73-0.74 (d, 2H), 0.86-0.89 (m, 7H), 0.95-1.69 (m, 12H), 2.61-2.68 (m, 2H), 3.02 (t, 1H), 3.69-3.76 (t, 1H), 4.36-4.41 (m, 2H), 6.82-6.87 (d, 1H), 7.29 (s, 1H), 7.37-7.49 (m, 5H), 7.86-7.91 (m, 1H), 8.17-8.19 (d, 3H), 8.32-8.33 (d, 1H), 10.79 (s, 1H).

714

Example 8

(2R)-2-(2-Amino-2-methylpropanoylamino)-N-{1-[2-(4-methylpiperidyl)-2-oxo-1-phenylethyl]imidazol-4-yl} hexanamide Bistrifluoroacetic Acid

5

20

This compound was obtained from (2R)-2-{2-[(tert-butoxy) carbonylamino]-2-methylpropanoylamino}-N-{1-[2-(4-10 methyl piperidyl)-2-oxo-1-phenylethyl]imidazol-4-yl}hexanamide as an off white solid in 90% yield. MS (FD+) m/z 496.3 (M*). Anal. calcd. for C₂₇H₄₀N₆O₃·2C₂HF₃O₂: C: 51.38; H: 5.84; N: 11.60. Found: C: 51.38; H: 5.83; N: 11.59. ¹H nmr (DMSO-d₆): δ 0.73-0.74 (d, 2H), 0.84-15 0.89 (m, 5H), 1.00-1.28 (m, 1H), 1.28-1.30 (m, 5H), 1.49-1.69 (m, 11H), 2.64-2.66 (m, 1.5H), 3.02 (t, 0.5H), 3.69-3.72 (t, 1H), 4.39-4.43 (m, 2H), 6.82-6.84 (m, 1H), 7.28-7.30 (m, 1H), 7.38-7.49 (m, 5H), 7.87-7.90 (m, 1H), 8.16-8.17 (d, 3H), 8.32-8.33 (d, 1H), 10.80 (s, 1H).

715

Example 9

(2R)-2-(2-Amino-2-methylpropanoylamino)-N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]hexanamide

Bistrifluoroacetic Acid

5

25

This compound was obtained from (2R)-2-{2-[(tert-10 butoxy) carbonylamino]-2-methylpropanoylamino}-N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]hexanamide as an off white solid in 92% yield. MS (FD+) m/z 468.2 (M⁺).

Anal. calcd. for C₂₅H₃₆N₆O₃·2C₂HF₃O₂: C: 50.00; H: 5.50; N: 12.06. Found: C: 49.79; H: 5.55; N: 12.07. ¹H nmr

15 (DMSO-d₆): δ 0.84-0.88 (m, 3H), 1.28-1.30 (m, 4H), 1.50-1.52 (d, 6H), 1.71-1.78 (m, 6H), 2.87-2.89 (m, 1H), 3.39-3.40 (m, 2H), 3.67-3.69 (m, 1H), 4.37-4.40 (m, 1H), 6.62 (s, 1H), 7.38 (s, 1H), 7.42-7.49 (m, 5H), 8.09-8.13 (d, 1H), 8.20-8.21 (m, 3H), 8.37-8.38 (d, 1H), 10.98 (s, 1H), 11.69

20 (s, broad, 1H).

Example 10

N-((1R)-1-{N-[1-(2-0xo-1-phenyl-2-pyrrolidinylethyl) imidazol-4-yl]carbamoyl}-2-(3-thienyl)ethyl)-2-amino-2methylpropanamide Bistrifluoroacetic Acid

716

This compound was obtained from $N-((1R)-1-\{N-[1-(2-oxo-$ 1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]carbamoyl}-2-(3thienyl)ethyl)-2-[(tert-butoxy)carbonylamino]-2-methyl 5 propanamide as an off white solid in 90% yield. MS (FD+) m/z 508.2 (M^{+}). Anal. calcd. for $C_{26}H_{32}N_{6}O_{3}S\cdot 2C_{2}HF_{3}O_{2}$: C: 48.91; H: 4.65; N: 10.9911.41. Found: C: 48.68; H: 11.26. ¹H nmr (DMSO-d₆): δ 1.40 (s, 3H), 1.50-4.53; N: 1.52 (d, 3H), 1.72-1.85 (m, 4H), 2.92-2.93 (m, 1H), 3.22 (m, 10 1H), 3.32 (m, 1H), 3.39-3.42 (m, 2H), 3.67-3.69 (m, 1H), 4.70-4.74 (m, 1H), 6.59 (s, 1H), 6.93-6.98 (d, 2H), 7.32-7.33 (d, 1H), 7.39-7.49 (m, 6H), 7.92-7.96 (d, 1H), 8.15-8.16 (d, 3H), 8.57-8.58 (d, 1H), 8.70 (s, broad, 1H), 10.99 15 (s, 1H).

717

Example 11

N-[(1R)-1-(N- $\{1-[2-(4-Methylpiperidy1)-2-oxo-1-phenylethy1]$ imidazol-4-yl $\}$ carbamoyl)-2-(3-thienyl)ethyl]2-amino-2-methyl propanamide Bistrifluoroacetic Acid

5

This compound was obtained from $N-[(1R)-1-(N-\{1-[2-(4-1)])]$ 10 methylpiperidyl)-2-oxo-1-phenylethyl]imidazol-4-yl} carbamov1)-2-(3-thieny1)ethy1]-2-[(tert-butoxy) carbonyl amino]-2-methylpropanamide as an off white solid in 84% yield. MS (FD+) m/z 536.2 (M^{+}). Anal. calcd. for $C_{28}H_{36}N_6O_3S\cdot 2C_2HF_3O_2$: C: 50.26; H: 5.01; N: 10.99. Found: 50.06; H: 4.89; N: 11.00. ¹H nmr (DMSO-d₆): δ 15 0.73-0.74 (d, 2H), 0.89-0.90 (d, 2H), 0.95-1.20 (m, 1H), 1.28-1.52 (m, 8H), 1.60-1.63 (m, 1H), 2.64-2.69 (m, 1.5H), 3.03 (t, 0.5H), 3.19-3.22 (t, 1H), 3.30-3.34 (t, 1H), 3.71-3.75 (m, 1H), 4.40-4.44 (m, 1H), 4.71 (m, 1H), 6.85-6.98 (m, 1H)3H), 7.32-7.48 (m, 7H), 7.92-7.97 (m, 1H), 8.16-8.17 (d, 20 3H), 8.57-8.58 (d, 1H), 11.02 (s, 1H), 11.3 (s, broad, 1H).

718

Example 12

N-((1R)-2-Benzo[b]thiophen-3-yl-1-{N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]carbamoyl}ethyl)-2-amino-2-methylpropanamide Bistrifluoroacetic Acid

5

This compound was obtained from N-((1R)-2-benzo[b]thiophen-3-yl-1-{N-[1-(2-oxo-1-phenyl-2-10 pyrrolidinylethyl)imidazol-4-yl]carbamoyl}ethyl)-2-[(tert-butoxy)carbonylamino]-2-methyl propanamide as an off white solid in 89% yield. MS (FD+) m/z 557.9 (M⁺). ¹H nmr (DMSO-d₆): δ 1.21-1.22 (d, 2H), 1.28-1.29 (d, 1H), 1.46-1.49 (m, 3H), 1.75-1.87 (m, 4H), 2.93-2.95 (m, 1H), 3.36-3.42 (m, 3.5H), 3.68-3.69 (m, 1.5H), 4.95-4.98 (m, 1H), 6.57 (s, 1H), 7.39-7.54 (m, 9H), 7.79-8.05 (m, 6H), 8.15 (m, 0.5H), 8.30 (m, 0.5H), 8.55-8.56 (m, 1H), 10.95-11.05 (d, 1H).

719

Example 13

N-[(1R)-2-Benzo[b]thiophen-3-yl-1-(N-{1-[2-(4-methyl piperidyl)-2-oxo-1-phenylethyl]imidazol-4-yl}carbamoyl) ethyl]-2-amino-2-methylpropanamide Bistrifluoroacetic Acid

5

This compound was obtained from N-[(1R)-2-benzo[b]thiophen-3-yl-1-(N-{1-[2-(4-methylpiperidyl)-2-oxo-1-phenylethyl] imidazol-4-yl}carbamoyl)ethyl]-2-[(tert-butoxy)carbonyl amino]-2-methylpropanamide as an off white solid in 72% yield. MS (FIA) m/z 586.7 [(M+H)⁺]. ¹H nmr (DMSO-d₆): δ 0.74-0.76 (d, 2H), 0.89-0.90 (d, 2H), 1.0-1.64 (m, 10H), 2.65-2.70 (m, 1.5H), 3.04 (t, 0.5H), 3.15-3.35 (m, 1.5H), 3.62-3.75 (m, 1H), 4.42-4.45 (m, 1H), 4.97 (m, 1H), 6.84-6.89 (d, 1H), 7.39-7.57 (m, 10H), 7.79-8.20 (m, 6H), 8.36 (t, 0.5H), 8.53-8.57 (m, 1H), 10.97-11.06 (d, 1H).

720

Example 14

N-[(1R)-1-(N-{1-[(N,N-Dimethylcarbamoyl)-2-naphthylmethyl})
imidazol-4-yl}carbamoyl)-2-indol-3-ylethyl]-2-amino-2-methyl
propanamide Dihydrochloride

5

This compound was obtained from the reaction of 2-[4-((2R)-2-{2-[(tert-butoxy)carbonylamino]-2-

721

Example 15

 $N-((1R)-2-(2-Naphthyl)-1-\{N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]carbamoyl]ethyl)-2-amino-2-methylpropanamide$

5

This compound was obtained from 2-[4-((2R)-2-{2-[(tert-butoxy) carbonylamino]-2-methylpropanoylamino}-3-(2-naphthyl)

10 propanoylamino)imidazolyl]-2-phenylacetic acid and pyrrolidine followed by deprotection according to the general procedure to give a tan foam in 53% yield. MS (FD⁺) m/z 552 (M⁺). ¹H nmr (CDCl₃): δ 1.06-1.25 (m, 9H), 1.81-1.92 (m, 3H), 3.12-3.13 (m, 1H), 3.44-3.61 (m, 4H), 5.03-5.05 (m, 1H), 5.88 (s, 1H), 7.26-7.44 (m, 13H), 7.6 (m, 0.5H), 7.75-7.76 (m, 1H), 8.15 (m, 0.5H), 8.24 (m, 1H), 10.5 (m, 1H).

722

Exampl 16

Methyl 1-(2-{4-[(2R)-2-(2-Amino-2-methylpropanoylamino)-3-(2-naphthyl)propanoylamino]imidazolyl}-2-phenylacetyl)(2S)
pyrrolidine-2-carboxylate Bistrifluoroacetic Acid

5

10

This compound was obtained from 2-[4-((2R)-2-{2-[(tert-butoxy)carbonylamino]-2-methylpropanoylamino}-3-(2-naphthyl) propanoylamino)imidazolyl]-2-phenylacetic acid and L-proline methyl ester followed by deprotection according to the general procedure to give a white solid in 76% yield. MS (FD⁺) m/z 610.0 (M⁺). IR (KBr): 1744.15, 1669.60, 1535.15, 1437.31, 1201.83, 1136.77, 721.83.

723

EXAMPLES PART 8

Preparation 1

2-(4-Nitroimidazolyl)-2-phenylacetic acid

5

Lithium hydroxide (18.1 g, 750 mm, 2 eq) was added to a stirred slurry of ethyl 2-(4-nitroimidazoyl)-2phenylacetate (Preparation 11 from Examples Part 1) 10 (104 g, 379 mm) in 250 mL of ethanol. Deionized water was added to the resulting mixture and the stirring was continued for 4 hours. The ethanol was removed under vacuum and the resulting aqueous solution was washed with 100 mL of diethyl ether. 15 The aqueous layer was diluted with 100 mL of deionized water and the pH was adjusted to 1.8 with concentrated HCl after cooling to 12 ℃. resulting slurry was stirred for 30 minutes at less 20 than 5 degrees and filtered. The wet cake was washed with 100 mL of deionized water and dried under a stream of air on the filter overnight to vield 90.34 g (96%) of a brown solid. The product may be recrystallized from isopropyl alcohol to give 72.31 g (80% recovery, 77% overall yield) of a tan 25 solid. Elemental analysis: Calculated: %C 53.45, %H 3.67, %N 16.97; Found: : %C 53.67, %H 3.79, %N 16.65. MS: 247 (M^{+}) : IR (cm^{-1}) 1719; H¹ nmr (d^{6}) DMSO): d 6.51 (s, 1H), 7.43-7.55 (m, 5H), 7.95 (s, 1H), 8.40 (s, 1H) 30

724

Preparation 2

1-(4-Methylpiperidiny1)-2-(4-nitroimidazoly1)-2phenylethan-1-one

5

N-Methyl morpholine (2.22 ml, 1.0 eg) was added to a 10 stirred solution of 2-(4-nitroimidazoly1)-2phenylacetic acid (1) (5.0 g, 20.2 mm) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (3.61 g, 20.2 mm, 1.0 eq) in 50 ml of anhydrous tetrahydrofuran at 25°C. After stirring the reaction mixture at ambient temperature for 1 h, 4-methylpiperidine (2.39 mL, 15 24.14 mm, 1.1 eq) was added and stirring continued for 50 min. The reaction mixture was quenched by the addition of 1M HCl. The layers were separated and the organic layer was washed with saturated 20 NaHCO3. The resulting emulsion was broken and all the solids dissolved by the addition of deionized water. The layers were separated and the organic layer washed with brine, dried over MgSO4 and filtered. The volatiles were removed under vacuum 25 to give a tan foam (6.08 g, 91% yield). The foam was slurried in hexane overnight to achieve partial crystallization of the product. This was filtered to give a cream colored solid (1.29g, 19%) which assayed 97% by HPLC. Analytical data are under 30 PX026948. The gummy residue was treated with

725

refluxing methyl t-butyl ether to facilitate crystallization. After refluxing, the slurry was cooled and filtered to give a cream colored product (3.02 g, 45%). Elemental analysis: Calculated: %C 62.18, %H 6.14, %N 17.06; Found: %C 62.05, %H 3.79, %N 16.65 MS: 328 (M⁺) IR (cm⁻¹)1659; H¹ nmr: d (d⁶ DMSO):0.5 (m,0.4H), 0.70 (d, J+6Hz, 1.5 H), 0.87 (d, J=6Hz, 1.5 H), 0.75 - 1.65 (m, 4.6H), 2.50-2.70 (m, 2.5H), 2.9-3.1 (m, 0.5H), 3.5-3.7 (m, 1H), 4.28-4.45 (m, 1H), 6.89 (s, 0.5H), 6.94 (s, 0.5H), 7.4-7.5 (m, 5H), 7.79 (s, 1H), 8.18 (s, 1H)

Preparation 3

2-(4-Nitroimidazolyl)-2-phenyl-1-pyrrolidinylethane-

15 **1-one**

N-Methyl morpholine (22.25 ml, 2 eq) was added to a stirred solution of 2-(4-nitroimidazolyl)-2phenylacetic acid (1) (25.03 g, 101.2 mm) and 2chloro-4,6-dimethoxy-1,3,5-triazine (18.1 g, 101.2 mm, 1.0 eq) in 50 ml of anhydrous tetrahydrofuran at 25°C. After stirring the reaction mixture at ambient temperature for 1 h, 7.2 mL (101.2 mm, 1.0 eq) of pyrrolidine was added dropwise. The reaction was stirred at room temperature for 2 hours. The reaction mixture was quenched by the addition of 200 mL of ethyl acetate and 200 mL of 1M HCl. The

726

washed with 100 mL of saturated sodium bicarbonate solution. The mixture resulting from the bicarbonate wash was diluted 1:1 with deionized water to dissolve the resulting solids and the layers were separated. The organic layer was washed with brine, dried over magnesium sulfate, filtered and the volatiles were removed under vacuum to give a brown foam. This foam was dissolved in methanol, diethyl ether and methylene chloride. Evaporation of the solvents overnight yielded a brown solid 10 which was slurried in 200 mL of diethyl ether for 4 hours. The resulting slurry was filtered and the cake was washed with diethyl ether. The solids were dried under vacuum overnight to give a cream colored 15 product (21.68 g, 71%) $d (d^6 DMSO): 1.69-1.84 (m, 3H), 2.80-2.85 (m, 0.7H),$ 3.32 - 3.41 (m, 3.6H), 3.64-3.67 (m, 0.7H), 6.65 (s, 1H), 7.42-7.50 (m, 5H), 7.83 (s, 1H), 8.22 (s, 1H)

727

Preparation 4

5

N-Methyl morpholine (17.79 ml, 2 eq) was added to a stirred solution of 2-(4-nitroimidazolyl)-2phenylacetic acid (1) (20 g, 80.9 mm) and 2-chloro-4,6-dimethoxy-1,3,5-triazine (14.45 g,80.1 mm, 1.0 10 eq) in 175 ml of anhydrous tetrahydrofuran at 25°C. After stirring the reaction mixture at ambient temperature for 1.5 h, 14.45g (80.9 mm, 1.0 eq) of proline methyl ester hydrochloride was added. 15 reaction was stirred overnight at room temperature and the solvent was evaporated under a stream of nitrogen. The residue was partitioned between 200 Ml of 1M HCl and 200 mL of deionized water. layers were separated and the organic layer was washed with saturated sodium bicarbonate solution. 20 A quantity of water sufficient to dissolve the resulting solids was added. The organic layer was washed with brine, dried over magnesium sulfate, filtered and the volatiles were removed under vacuum to give a brown foam. The foam was dissolved in 25 methyl t-butyl ether, methylene chloride and methanol to effect crystallization. The resulting slurry was filtered and dried under vacuum to give 17.45 g (60%) of a cream colored product MS: 358

728

 (M^{+}) IR (cm^{-1}) : 1744, 1668; H^{1} nmr d $(d^{6}$ DMSO): 1.42-1.46 (m, 3H), 1.6-1.9 (m, 1H), 2.42-2.63 (m,1H), 3.21 (s, 1.5H), 3.25 (s, 1.5H), 3.30-3.36 (m, 1H), 4.01-4.06 (m, 1H), 6.36 (s, 0.5H), 6.43 (s, 0.5h), 7.0-7.2 (m, 5H), 7.38 (s, 0.5H), 7.41 (s, 0.5H), 7.75 (s, 0.5H), 7.83 (s, 0.5H)

Preparation 5

Methyl (2S)-1-[2-(4-aminoimidazolyl)-2-phenylacetyl]pyrrolidine-2-carboxylate, dihydrochloride

15

20

25

10

Ethanol (12 mL) was added to a mixture of methyl (2S)-1-[2-(4-nitroimidazolyl)-2-phenylacetyl]pyrrolidine-2-carboxylate (4) (148 mg,0.4 mm) and 10% Pd on carbon (15 mg) in a Bradley hydrogenation apparatus. The stirred reaction mixture was subjected to a 60 psi H₂ atmosphere and warmed to 60 °C. After 2 hours, the reaction mixture was cooled to room temperature and the catalyst was removed by filtration. Anhydrous HCl gas was added to the filtered solution until saturation. The volatiles were then removed under vacuum to give 0.15 g (100%) of a yellow solid mixed with a brown gum.

729

Preparation 6

2-(4-aminoimidazolyl)-2-phenyl-1-pyrrolidinylethan-1-one, dihydrochloride

5

Ethanol (200 mL) was added to a mixture of 2-(4nitroimidazolyl)-2-phenyl-1-pyrrolidinylethan-1-one (3) (0.752 g, 2.8 mm) and 10% Pd on carbon (75 mg) in a Bradley hydrogenation apparatus. The stirred 10 reaction mixture was subjected to a 60 psi H2 atmosphere and warmed to 60 °C. After 2 hours, the reaction mixture was cooled to room temperature and the catalyst was removed by filtration. Anhydrous HCl gas was added to the filtered solution until 15 saturation. The volatiles were then removed under vacuum to give a light yellow foam. Diethyl ether and methylene chloride (25:1) were added to the foam and the resulting mixture was stirred overnight to achieve crystallization. The resulting slurry was 20 filtered and the cake was washed with diethyl ether. The cake was dried under vacuum to give 0.659 g (93%) of a yellow solid. LGD 208.

730

Preparation 7

2-(4-aminoimidazoly1)-1-(4-methylpiperidiny1)-2phenylethan-1-one, dihydrochloride

5

Methanol (140 mL) was added to a mixture of 1-(4methylpiperidyl)-2-(4-nitroimidazolyl)-2phenylethan-1-one (2) (2.16 g, 6.5 mm) and 0.214 g of 10% palladium on carbon in a Bradley 10 hydrogenation apparatus. The stirred reaction mixture was subjected to a 60 psi H2 atmosphere and warmed to 60 °C. After 2 hours, the reaction mixture was cooled to room temperature and the catalyst was removed by filtration. Anhydrous HCl 15 gas was added to the filtered solution until saturation. The volatiles were then removed under vacuum and the residue was crystallized from methylene chloride: methyl t-butyl ether 1:3. 20 resulting slurry was filtered and the wet cake was washed with methyl t-butyl ether and dried under vacuum to give 2.23 (93%)g of a yellow solid. MS: 298 $(M+1^+)$; IR $(cm^{-1}):1648$; H¹ nmr d $(d^6 DMSO):0.11-$ 0.21 (m, 0.5H), 0.725 (d, J=4.9 Hz, 1.5H), 1.11-1.2125 (m, 0.5H), 1.25-1.40 (m, 1H), 1.59-1.80 (m, 2H),2.72-2.81 (m, 1H) 3.08-3.15 (m, 0.5H) 3.32 (s, 0.5H), 3.32 (s, 0.5H). 3.32 (s. 0.5H) 3.75 (d. J=8 Hz, 1H), 4.50-4.57 (dd. J=7.3 Hz, 1H). 4.95 (s, 1H), 6.8 (s, 0.5H), 6.9 (s, 0.5H), 7.39-7.60 (m, 5H), 8.29 (s, 30 0.5H), 8.45 (s, 0.5H)

731

Example 1

N-((1R)-2-indol-3-yl-1-{N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]carbamoyl}ethyl)-2[(tert-butoxy)carbonylamino]-2-methylpropanamide

5

N-Methyl morpholine (0.28 mL, 8.32 mm, 1 eq) was added to a stirred slurry of 2-chloro-4,6-dimethoxy-1,3,5-triazine (0.46 g, 2.57 mm, 1 eq) and (2R)-2-{2-[(tert-butoxy)carbonylamino]-2-methylpropanoylamino}-3-indol-3-ylpropanoic acid (1g, 2.57mm) in 10 mL of anhydrous tetrahydrofuran cooled to less than 0 °C. After 1.5 hours, 2-(4-aminoimidazolyl)-2-phenyl-1-pyrrolidinylethan-1-one, hydrochloride (0.97g, 2.82 mm, 1.1 eq) was added and stirring was continued at ice bath temperatures.

The reaction was stirred for 4 hours and quenched by the addition of 15 mL of deionized water and ethyl acetate. The ethyl acetate layer was washed with a saturated sodium bicarbonate solution, dried over magnesium sulfate and the volatiles were removed under vacuum to give the crude product as a light

purple foam (1.4 g, 84%) The crude product was

732

purified by preparative chromatography to provide 0.52 g (31.5%) of the product as a foam. 1 H nmr (CDCl₃): δ 1.10-1.40 (m, 15H), 1.67-1.92 (m, 3H), 2.92-3.60 (m, %H), 4.90 (s, broad, 1H), 5.33 (s, broad, 1H), 5.85 (d, 1H), 6.80-7.05 (m, 3H), 7.13-7.39 (m, 10H), 7.44-7.80 (m, 2H), 8.96 (s, broad, 1H), 10.20 (s, broad, 1H).

Example 2

N-((1R)-2-indol-3-yl-1-{N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]carbamoyl}ethyl)-2-amino-2-methylpropanamide, 2,2,2-trifluoroacetic acid, 2,2,2-trifluoroacetic acid salt

15

Trifluoroacetic acid (0.57 mL, 7.4 mm, 33 eq) was added to a stirred solution of N-((1R)-2-indol-3-yl-1-{N-[1-(2-oxo-1-phenyl-2-

pyrrolidinylethyl)imidazol-4-yl]carbamoyl)ethyl)-2[(tert-butoxy)carbonylamino]-2-methylpropanamide (8)
(0.152 g, 0.22 mm) in 5 mL of methylene chloride.
After stirring at room temperature for 3 hours, the reaction mixture was diluted with 50 mL of diethyl
ether. The resulting solids were isolated by centrifugation and washed with diethyl ether. The solids were dried under vacuum to give the product

733

as a cream colored solid (0.084 g, 48%) MS (FD+) m/z 541 (M*)Anal. calcd. for $C_{30}H_{35}N_7O_3 \cdot 2C_2HF_3O_2$: C: 53.06; H: 4.85; N: 12.74. Found: C: 52.93; H: 4.88; N: 12.55. ¹H nmr (DMSO-d₆): δ 1.29 (s, 3H), 1.46-1.48 (d, 3H), 1.72-1.88 (m, 4H), 2.94 (m, 1H), 3.06-3.07 (m, 1H), 3.19-3.20 (m, 1H), 3.40-3.41 (d, 2H), 3.67-3.69 (m, 1H), 4.78 (s, broad, 1H), 6.53 (s, 1H), 6.93-6.97 (m, 1H), 7.06 (m, 1H), 7.20 (d, 1H), 7.31-7.36 (m, 2H), 7.42-7.42 (m, 4H), 7.73-7.80 (m, 2H), 8.01 (s, broad, 2H), 8.36-8.38 (d, 1H), 10.82-10.85 (d, 2H).

Example 3

Methyl $1-\{2-[4-((2R)-2-\{2-[(tert-$

butoxy)carbonylamino]-2-methylpropanoylamino}-3indol-3-ylpropanoylamino)imidazolyl]-2phenylacetyl}pyrrolidine-2-carboxylate

20

NMR data support the assigned structure. MS: 699.9

734

Example 4

Methyl 1-(2-{4-[(2R)-2-(2-amino-2-methylpropanoylamino)-3-indol-3-ylpropanoylamino]imidazolyl}-2-phenylacetyl)pyrrolidine-2-carboxylate, dihydrochloride

10

15

5

NMR data support the assigned structure. MS: 599

Example 5

N-((1R)-2-(2-naphthyl)-1-{N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]

carbamoyl}ethyl)-2-[(tert-butoxy)carbonylamino]-2-methylpropanamide

735

NMR data support the assigned structure. MS: 652

Example 6

5 N-((1R)-2-(2-naphthyl)-1-{N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]carbamoyl}ethyl)-2-amino-2-methylpropanamide, dihydrochloride

10

NMR data support the assigned structure. MS:552

Example 7

N-[(1R)-1-(N-{1-[2-(4-methylpiperidy1)-2-oxo-1-phenylethyl]imidazol-4-yl}carbamoy1)-2-(2-naphthyl)ethyl]-2-[(tert-butoxy)carbonylamino]-2-methylpropanamide

20

NMR data support the assigned structure. MS: 680

736

Exampl 8

N-[(1R)-1-(N-{1-[2-(4-methylpiperidyl)-2-oxo-1-phenylethyl]imidazol-4-yl)carbamoyl)-2-(2-naphthyl)ethyl]-2-amino-2-methylpropanamide dihydrochloride

H—CI HN N—N—15

NMR data support the assigned structure. MS: 581 $(M+1^+)$

Example 9

N-((1R)-1-{N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]carbamoyl}-2-phenylethyl)-2-[(tert-butoxy)carbonylamino]-2-methylpropanamide

NMR data support the assigned structure. MS: 602

737

Example 10

N-((1R)-1-{N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]carbamoyl}-2-phenylethyl)-2-amino-2-methylpropanamide,
dihydrochloride

5

NMR data support the assigned structure. MS: 502

10

Example 11

N-[(1R)-1-(N-{1-[2-(4-methylpiperidy1)-2-oxo-1-phenylethyl]imidazol-4-yl}carbamoyl)-2-phenylethyl]2-[(tert-butoxy)carbonylamino]-2-methylpropanamide

15

NMR data support the assigned structure. MS: 630

738

Example 12

N-[(1R)-1-(N-{1-[2-(4-methylpiperidyl)-2-oxo-1-phenylethyl]imidazol-4-yl}carbamoyl)-2-phenylethyl]2-amino-2-methylpropanamide, dihydrochloride

5

NMR data support the assigned structure. MS: 530

10

Example 13

(2R)-2-{2-[(tert-butoxy)carbonylamino]-2methylpropanoylamino}-N-[1-(2-oxo-1-phenyl-2pyrrolidinylethyl)imidazol-4-yl]-4-phenylbutanamide

15

NMR data support the assigned structure. MS: 616

739

Example 14

(2R)-2-(2-amino-2-methylpropanoylamino)-N-[1-(2-oxo-1-phenyl-2-pyrrolidinylethyl)imidazol-4-yl]-4-phenylbutanamide, dihydrochloride

5

NMR data support the assigned structure. MS: $517 \, (m+1^+)$


10

Example 15

(2R)-2-(2-[(tert-butoxy)carbonylamino]-2-

15 methylpropanoylamino)-N-{1-[2-(4-methylpiperidinyl)-2-oxo-1-phenylethyl]imidazol-4-yl}-4-

phenylbutanamide



740

NMR data support the assigned structure.

Example 16

(2R)-2-(2-amino-2-methylpropanoylamino)-N-{1-[2-(4-5 methylpiperidinyl)-2-oxo-1-phenylethyl]imidazol-4-yl}-4-phenylbutanamide, dihydrochloride

10 NMR data support the assigned structure. MS: 544

741

Example 17

Methyl $1-\{2-\{4-((2R)-2-\{2-\{(tert-$

butoxy)carbonylamino]-2-methylpropanoylamino}-4-

phenylbutanoylamino)imidazolyl]-

2-phenylacetyl)pyrrolidine-2-carboxylate

10 NMR data support the assigned structure.

742

Exampl 18

Methyl 1-(2-{4-[(2R)-2-(2-amino-

2-methylpropanoylamino)-4-

phenylbutanoylamino]imidazolyl)

-2-phenylacetyl)pyrrolidine-2-carboxylate,
dihydrochloride

5

10 NMR data support the assigned structure. MS:575 $(m+1^+)$

PCT/US98/17201 WO 99/08697

743

EXAMPLES PART 9

Example 1

Pituitary Cell Culture Assay for Growth Hormone Secretion

5

10

15

Thirty-two 250 g male Sprague-Dawley rats were used for each assay. The animals were killed by decapitation and anterior pituitaries were removed and placed into ice cold culture medium. The pituitaries were sectioned into eighths and enzymatically digested using trypsin (Sigma Chemical) to weaken connective tissue. Pituitary cells were dispersed by mechanical agitation, collected, pooled and then seeded into 24-well plates (300,000 cells/well). After 4 days of culture, the cells formed an even monolayer. Cells were then washed with medium and challenged to secrete GH by the addition of GH secretagogues to the medium. After 15 min at 37 °C, the medium was removed and stored frozen until radioimmunoassays for rat GH were performed. Doses of secretagogue were added in quadruplicate. Representative 20 Data is provided in Table 1 below. Compounds disclosed herein are active in the assay as described. Both EC50 and efficacy values were calculated by the 4-parameter logistic equation. Such values were pooled and represented as mean +/- standard error, when appropriate.

Table 1		
EXAMPLES	GH	
PART 1	secretion	
Example #	EC ₅₀ (mM)	
6	5.53	
8 2.39		

744

Example 1

Initial Efficacy

5 In vivo growth hormone secretion and growth hormone selectivity (corticosterone levels) screening were assessed in immature rats, initially with parenteral dosing, followed later by oral dosing with selected compounds. In vivo growth hormone secretagogue activity is examined in immature 10 rats (approx. 75 g body weight) by parenteral (s.c.)administration initially. Compounds displaying potent in vitro activity (≤ approx. 0.5 mM) are tested in vivo. Compounds displaying growth hormone secretory activity at < 30 mg/kg s.c. are further tested for growth hormone 15 selectivity by examining effects on serum corticosterone in immature rats. Growth stimulating efficacy has been demonstrated for GHRP-2 (parenteral administration) in such an in vivo assay.

745

Example 2

Effects of Growth Hormone Secretagogues in the Rat Model of Congestive Heart Failure

5

15

20

25

The effects of growth hormone and a GHRP (GRP-2) on the progression of heart failure were examined in a rat genetic model of heart failure, the SHHF/Mcc-fa^{CD} rat (McCune et al., Mechanisms of Heart Failure, Singal, et al., eds., 10 Kluwer Academic Publishers, 1995). This model represents a well-established small animal model of heart failure that mimics the important aspects of dilated cardiomyopathy, the setting in which dramatic effects of growth hormone have been observed clinically (Fazio et al., New England Journal of Medicine, 334: 809-14, 1996).

Rats were dosed with 0.05 to 0.01 mg/kg of GRP-2 per day, or with 0.1 to 1.0 mg/kg of growth hormone per day. The effects of growth hormone and GHRP-2 were compared to those of a positive control drug in this model, Captopril®. during a 2 month course of therapy, starting during the early stages of heart failure (14-16 months of age). Hemodynamic as well as biochemical indices of heart failure together with growth hormone, levels of IGF-1, insulin, and atrial natriuretic peptide (ANP), and overall survival were monitored.

The data presented in Table 2, below, indicates an increased survival rate in rats treated with GRP-2 over those treated with growth hormone. Importantly, none of the animals treated with GRP-2 died of heart failure.

746

Table 2
Survival Analysis

5	Group	Survival	Cause of Death/ Sacrifice
	Control	3/5	CHF (2)
	GH low	2/5	CHF(1), CHF(1), leuk(1)
	GH high	2/5	CHF(1), CHF+PT(1), CHF(1)
	GRP-2 low	4/5	leuk(1)
10	GRP-2 high	4/5	PT(1)

PT= pituitary tumor leuk= leukemia

30

Data presented in Figures 1-8 provide further evidence of the effectiveness of growth hormone secretagogue therapy in the rat model of CHF. Specifically, Figure 1 shows an increase in heart and lung weight in animals treated with growth hormone, as opposed to a trend toward decreased weights in those animals treated with GRP-2.

Figures 2-4 show an increase in the levels of corticosterone levels in those test animals treated with growth hormone, as opposed to those animals treated with GRP-2. Increased levels of corticosterone were observed in rats treated with growth hormone, which was seen as a function of heart failure progression.

Figures 5-7 show an increase in atrial natriuretic peptide (ANP) in animals treated with growth hormone, as opposed to a dose-dependent trend toward decreased ANP in those treated with GRP-2. Levels of atrial natriuretic peptide (ANP) are known to increase in heart failure.

Figure 8 shows the end systolic volume in rats treated with growth hormone or GRP-2, which indicates a decrease in

747

ventricular systolic function in rats treated with growth hormone, as opposed to rats treated with GRP-2.

Overall, these results indicating a decrease in the incidence of heart failure in rats treated with GRP-2 versus those treated with growth hormone are surprising and unexpected.

10 Example 3

20

25

30

Effect of Growth Hormone Secretagogues in the Canine Model of Congestive Heart Failure

Dog models appear to closely mimic human heart failure from the standpoint of response to drug therapy (Sabbah et al., Circulation 89: 2852-9, 1994; Tsutsui et al., J. Clin. Invest. 93: 2639-2648, 1994; and Wang et al., Amer. J. Physiol. 266: H670-80, 1994).

The effect of the administration of a growth hormone secretagogue may be examined in a canine model of congestive heart failure as follows. After table training for several weeks, the subject dogs are anesthetized and by means of a left thorocotomy, pacing wires are placed on the left atrial appendage and right ventricle, tygon catheters are inserted into the descending aorta and the right and left atria, a solid state micromanometer is inserted into the left ventricle, a flow probe is placed around the ascending aorta and the coronary artery, and piezoelectric crystals are placed in the left ventricle. If desired, renal and iliac arterial flow tubes may also be placed. The animals are permitted to recover for 14 days during which time basal measurements are obtained. Pacing of the right ventricle at approximately 240 beats/minute is initiated on day 14 after surgery. As demonstrated by hemodynamic data at baseline

748

and at three weeks after pacing in a conscious dog, this model not only achieves significant left ventricular dysfunction (for example, a decrease in left ventricular change in pressure over change in time and ejection fraction, increase in left ventricular end diastolic pressure and heart rate), but also demonstrates many of the signs of congestive heart failure seen in the clinic, including exercise intolerance, pulmonary edema, and ascites.

On day 14 after surgery, the subject animals are randomized (targeting five animals in each group). A compound of formula I, or GRP-2, or vehicle is administered beginning at the time of the initiation of pacing throughout duration of the study. Cardiac function, peripheral hemodynamics, exercise performance, and body weights are measured in the baseline study and weekly after initiation of pacing (three weeks total). Twenty-one days after initiation of pacing, the animals are sacrificed and the relative effect of administration of the growth hormone secretagogue on ventricular modeling, left ventricular mass, and body weight is determined.

749

PCT/US98/17201

We Claim:

WO 99/08697

- 1. A method of modulating cardiac function which comprises administering to a patient in need thereof an effective amount of a growth hormone secretagogue.
 - 2. A method according to Claim 1 wherein said growth hormone secretagogue comprises GRP-2.
- 3. A method according to Claim 1 wherein said growth hormone secretagogue comprises a compound of formula I

15

I

20 wherein:


25

A is C_1-C_6 alkyl, aryl, C_1-C_6 alkylaryl, C_1-C_6 alkyl(O) C_1-C_6 alkylaryl, C_1-C_6 alkyl(S) C_1-C_6 alkylaryl, indolyl, indolinyl, thienyl, $(C_1-C_6$ alkyl)thienyl, benzothienyl, benzofuranyl, naphthanyl, cyclohexyl, $(C_1-C_6$ alkyl)indolyl, $(C_1-C_6$ alkyl)benzothienyl, $(C_1-C_6$ alkyl)naphthanyl, $(C_1-C_6$ alkyl)benzofuranyl, and $(C_1-C_6$ alkyl)cyclohexyl;

 $\label{eq:BisnH2} B \ is \ NH_2, \ NHR_1, \ C_1-C_6 alkylNH_2, \ C_1-C_6 alkylNHR_1, \\ C_1-C_6 alkylarylNH_2, \ C_1-C_6 alkylarylNHR_1, \\$

```
C_1-C_6alkylcyclohexylNH<sub>2</sub>, C_1-C_6alkylcyclohexylNHR<sub>1</sub>,
     R_1-piperidin-3-yl(C_1-C_6alkyl),
     R_1-piperidin-2-yl(C_1-C_6alkyl), R_1-piperidin-4-yl(C_1-C_6alkyl),
     R_1-quinolin-2-yl(C_1-C_6alkyl),
 5 R_1-(2,4-dihydroquinolin-2-yl(C_1-C_6alkyl),
     R_1-isoquinolin-2-yl(C_1-C_6alkyl), and
     R_1-(2,4-dihydroisoquinolin-2-yl(C_1-C_6alkyl);
                 R_1 is hydrogen, C_1-C_6alkyl, C_1-C_6alkyl(OH), or
     C<sub>1</sub>-C<sub>6</sub>alkylidenyl(OH)R<sub>2</sub>;
                 R_2 is C_1-C_6alkyl, C_1-C_6alkenyl,
10
     C_1-C_6alkyl(0)C_1-C_6 alkyl, C(0)0-C_1-C_6 alkyl, aryl, or
     C<sub>1</sub>-C<sub>6</sub>alkylaryl;
                  X is C_1-C_6alkylidenyl, O, S, NH, or N(C_1-C_6alkyl);
                  V is selected from the group consisting of
15
```





752

and

W is S, O, NH, or CH2; 5 Y is N or CH; Z is N or CH; Y' is N or CH; Z' is N or CH: R_4 and R_5 are independently hydrogen, C_1 - C_6 alkyl, 10 aryl, C_1 - C_6 alkylaryl, $C(0)O(C_1$ - C_6 alkyl), $C(0)N(C_1-C_6alkyl)_2$, or $C_1-C_6alkylCOR_7$; R₇ is hydrogen, C₁-C₆alkyl, pyrrolidinyl, piperidinyl, homoproline, or proline; D is hydrogen, C₁-C₆alkyl, 15 C_1-C_6 alkyl(0)(CO) C_1-C_6 alkyl, C_1-C_6 alkyl(0)(CO) $N(C_1-C_6$ alkyl), C_1 - C_6 alkylaryl, $C(O)R_6$, C_1 - C_6 alkyl $(O)R_6$, C_1 - C_6 alkyl(OH), C_1 - C_6 alkylC(0)R₆, C_1 - C_6 alkylR₆, aryl, $(C_1$ - C_6 alkyl)NHSO₂(C_1 - C_6 alkyl), (C₁-C₆alkyl)NHSO₂(aryl); R₆ is H, C₁-C₆alkyl, aryl, naphthyl, 20 C₁-C₆alkylaryl, acetyl, NH₂, NH(C₁-C₆alkyl), $NH(C_1-C_6alkyl)O(C_1-C_6alkyl)$, $NH(C_1-C_6alkyl)S(C_1-C_6alkyl)$, $NH(C_1-C_6alkylidenyl)OCH_3$, $NH(C_1-C_6alkyl)aryl$, $NH(C_3-C_6 \text{ cycloalkyl})$, $NH(C_1-C_6\text{alkyl})C(O)(C_1-C_6\text{alkyl})$, $NH(C_1-C_6alkyl)NH(C_1-C_6alkyl)$, $NH(C_1-C_6alkyl)NH(C_1-C_6alkylaryl)$, 25 NHSO₂(C_1 - C_6 alkylaryl), NH(C_1 - C_6 alkyl) $C(0)O(C_1$ - C_6 alkyl), $NH(naphthyl), N(C_1-C_6alkyl)_2, N(C_1-C_6alkyl)(aryl)_1$ $N(C_1-C_6alkyl)(C_1-C_6alkylaryl), O(C_1-C_6alkyl), O(aryl),$ O(C₁-C₆alkylaryl), piperidinyl, piperidinyl-C(0)NH(C_1 - C_6 alkyl), piperidinyl-C(0)NH(C_1 -30 C6alkylaryl), piperidinyl-C(0)N(C1-C6alkyl)2. piperidinyl-C(O)N(C₁-C₆alkyl)(aryl),

pyrrolidinyl, pyrrolidinyl C(0)NH(aryl),


753

pyrrolidinyl C(O)NH(C1-C6alkyl),
pyrrolidinyl C(O)NH(C1-C6alkyl),
pyrrolidinyl C(O)NH(C1-C6alkyl),
pyrrolidinyl C(O)NH(C1-C6alkyl) (aryl),

pyrrolinyl, morpholino, hexamethyleneimino,
heptamethyleneimino, quinolinyl, 2,4-dihydroquinolinyl,
1,2,3,4-tetrahydroquinolinyl,
2,4-dihydroisoquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl,
indolinyl, an amino acid selected from the group consisting
of proline, homoproline, glycine, alanine, valine, leucine,
isoleucine, tyrosine, tryptophan, phenylalanine, serine,
threonine, asparagine, glutamic acid, aspartic acid,
lysine, arginine, glutamine, histidine, cysteine, and
methionine, or a nitrogen-containing heterocycle selected

from the group consisting of

754



 $\label{eq:continuous} E \mbox{ is hydrogen, C_1-C_6alkyl, $C(0)$C_1$-$C_6$alkyl, aryl, $(aryl)$C(0)$NR_6, $(aryl)$(C_1-C_6alkyl)$C(0)$R_6, $$$

5 C₁-C₆alkylaryl, C(0)aryl, C₁-C₆ alkylC(0)aryl, naphthyl, C₁-C₆alkylnaphthyl, C(0)naphthyl, C₁-C₆alkylC(0)naphthyl, heteroaryl, C₁-C₆alkylheteroaryl, C(0)heteroaryl, C₁-C₆ alkylC(0)heteroaryl, indanyl,

755

 C_1 - C_6 alkylindanyl, C(0)indanyl, C_1 - C_6 alkylC(0)indanyl, cycloalkyl;

or D and E combine to form indanyl, fluorenyl, or cycloalkyl;

G is hydrogen, C_1 - C_6 alkyl, aryl, C_1 - C_6 alkylaryl, and C_1 - C_6 alkenyl;

J is hydrogen, $C_1\text{--}C_6alkyl,$ aryl, and $C_1\text{--}C_6alkylaryl;$

L is hydrogen, C_1 - C_6 alkyl, $C(0)OC_1$ - C_6 alkyl, aryl,


10 C_1-C_6 alkylaryl, $C(O)OC_1-C_6$ alkylaryl, C_1-C_6 alkenyl, -F, and -CN, C_1-C_6 alkyl-OH, C_1-C_6 alkyl-O- C_1-C_6 alkyl, C_1-C_6 alkyl-C(O)R₆;

or a pharmaceutically acceptable salt or solvate thereof.

15

756

4. A method according to Claim 3 wherein A is selected from the group consisting of




5

5. A method according to Claim 3 wherein B is

10

- 6. A compound according to Claim 4 wherein ${\tt X}$ is NH.
- 7. A method according to Claim 3 wherein V is selected from the group consisting of



20

8. A method according to Claim 3 wherein D is $-C(0)R_6$, and R_6 is

1-pyrrolidinyl, 1-piperidinyl, 4-methoxy-1-piperidinyl,

9. A method according to Claim 3 wherein E is

OMe

CH₃

OCH₃

CF₃

- 10. A method according to Claim 1 wherein said patient 10 is a human.
 - 11. A method for the treatment or prevention of congestive heart failure which comprises administering to a patient in need of said treatment an effective amount of a growth hormone secretagogue, wherein said growth hormone secretagogue increases the levels of endogenous growth hormone.
- 12. A method according to Claim 11 wherein said growth 20 hormone secretagogue is GRP-2.

758

13. A method according to Claim 11 wherein said growth hormone secretagogue is a compound of formula I

5

I

wherein:

A is C₁-C₆alkyl, aryl, C₁-C₆alkylaryl,

C₁-C₆alkyl(O)C₁-C₆alkylaryl, C₁-C₆alkyl(S)C₁-C₆alkylaryl,

indolyl, indolinyl, thienyl, (C₁-C₆alkyl)thienyl,

benzothienyl, benzofuranyl, naphthanyl, cyclohexyl, (C₁-C₆alkyl)indolyl, (C₁-C₆alkyl)benzothienyl,

15 (C_1-C_6alkyl) naphthanyl, (C_1-C_6alkyl) benzofuranyl, and (C_1-C_6alkyl) cyclohexyl;

 $\label{eq:BisnH2} B is NH_2, NHR_1, C_1-C_6alkylNH_2, C_1-C_6alkylNHR_1, \\ C_1-C_6alkylarylNH_2, C_1-C_6alkylarylNHR_1, \\ C_1-C_6alkylcyclohexylNH_2, C_1-C_6alkylcyclohexylNHR_1, \\$

20 R₁-piperidin-3-yl(C₁-C₆alkyl),

 R_1 -piperidin-2-yl(C_1 - C_6 alkyl), R_1 -piperidin-4-yl(C_1 - C_6 alkyl), R_1 -quinolin-2-yl(C_1 - C_6 alkyl),

 $R_1-(2,4-dihydroquinolin-2-yl(C_1-C_6alkyl),$

 R_1 -isoquinolin-2-yl(C_1 - C_6 alkyl), and

25 R_1 -(2,4-dihydroisoquinolin-2-yl(C_1 - C_6 alkyl);

 R_1 is hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkyl(OH), or C_1 - C_6 alkylidenyl(OH) R_2 ;

 $R_2 \text{ is } C_1-C_6 \text{alkyl}, \ C_1-C_6 \text{alkenyl}, \\ C_1-C_6 \text{alkyl}, \ C_1-C_6 \text{ alkyl}, \ C_1-C_6 \text{ alkyl}, \ \text{aryl}, \ \text{or}$

759

C₁-C₆alkylaryl;

X is C_1-C_6 alkylidenyl, O, S, NH, or $N(C_1-C_6$ alkyl); V is selected from the group consisting of

5

760

and

W is S, O, NH, or CH2;

5 Y is N or CH;

Z is N or CH;

Y' is N or CH;

Z' is N or CH;

 R_4 and R_5 are independently hydrogen, C_1 - C_6 alkyl,

10 aryl, C_1 - C_6 alkylaryl, $C(0)O(C_1$ - C_6 alkyl),

 $C(0)N(C_1-C_6alkyl)_2$, or $C_1-C_6alkylCOR_7$;

 R_7 is hydrogen, C_1 - C_6 alkyl, pyrrolidinyl,

piperidinyl, homoproline, or proline;

D is hydrogen, C1-C6alkyl,

 $\begin{array}{lll} 15 & C_1-C_6alkyl\,(O)\,(CO)\,C_1-C_6alkyl\,, & C_1-C_6alkyl\,(O)\,(CO)\,N\,(C_1-C_6alkyl)_2\,, \\ & C_1-C_6alkylaryl\,, & C\,(O)\,R_6\,, & C_1-C_6alkyl\,(O)\,R_6\,, & C_1-C_6alkyl\,(OH)\,, & C_1-C_6alkyl\,(OH)_2\,, \\ & alkyl\,C\,(O)\,R_6\,, & C_1-C_6alkyl\,R_6\,, & aryl\,, & (C_1-C_6alkyl)\,NHSO_2\,(C_1-C_6alkyl)\,, \\ & & (C_1-C_6alkyl)\,NHSO_2\,(aryl)\,; \end{array}$


R₆ is H, C₁-C₆alkyl, aryl, naphthyl,

- C₁-C₆alkylaryl, acetyl, NH₂, NH(C₁-C₆alkyl), NH(C₁-C₆alkyl)O(C₁-C₆alkyl), NH(C₁-C₆alkyl)S(C₁-C₆alkyl), NH(C₁-C₆alkylidenyl)OCH₃, NH(C₁-C₆alkyl)aryl, NH(C₃-C₆ cycloalkyl), NH(C₁-C₆alkyl)C(O)(C₁-C₆alkyl), NH(C₁-C₆alkyl)NH(C₁-C₆alkyl)NH(C₁-C₆alkyl)NH(C₁-C₆alkyl), NH(C₁-C₆alkyl)NH(C₁-C₆alkyl), NH(C₁-C₆alkyl), NH(C₁-C₆alkyl)NH(C₁-C₆alkyl), NH(C₁-C₆alkyl)NH(C₁-C₆alkyl), NH(C₁-C₆alkyl), NH(C₁
- NHSO₂(C₁-C₆alkylaryl), NH(C₁-C₆alkyl)C(O)O(C₁-C₆alkyl),
 NH(naphthyl),N(C₁-C₆alkyl)₂, N(C₁-C₆alkyl)(aryl),
 N(C₁-C₆alkyl)(C₁-C₆alkylaryl), O(C₁-C₆alkyl), O(aryl),
 O(C₁-C₆alkylaryl), piperidinyl,
 piperidinyl-C(O)NH(C₁-C₆alkyl), piperidinyl-C(O)NH(C₁-
- C6alkylaryl), piperidinyl-C(O)N(C1-C6alkyl)2, piperidinyl-C(O)N(C1-C6alkyl)(aryl), pyrrolidinyl, pyrrolidinyl C(O)NH(aryl),

761

pyrrolidinyl $C(0)NH(C_1-C_6alkyl)$, pyrrolidinyl C(O)N(C1-C6alkyl)2, pyrrolidinyl C(0)NH(C1-C6alkylaryl), pyrrolidinyl C(O)NH(C1-C6alkyl)(aryl), pyrrolinyl, morpholino, hexamethyleneimino, heptamethyleneimino, quinolinyl, 2,4-dihydroquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,4-dihydroisoquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl, indolinyl, an amino acid selected from the group consisting 10 of proline, homoproline, glycine, alanine, valine, leucine, isoleucine, tyrosine, tryptophan, phenylalanine, serine, threonine, asparagine, glutamic acid, aspartic acid, lysine, arginine, glutamine, histidine, cysteine, and methionine, or a nitrogen-containing heterocycle selected 15 from the group consisting of





 $\label{eq:condition} E \mbox{ is hydrogen, } C_1-C_6alkyl, \mbox{ } C(O)C_1-C_6alkyl, \mbox{ } aryl, \\ (aryl)C(O)NR_6, \mbox{ } (aryl)(C_1-C_6alkyl)C(O)R_6, \\ \mbox{ }$

5 C₁-C₆alkylaryl, C(0)aryl, C₁-C₆ alkylC(0)aryl, naphthyl, C₁-C₆alkylnaphthyl, C(0)naphthyl, C₁-C₆alkylC(0)naphthyl, heteroaryl, C₁-C₆alkylheteroaryl, C(0)heteroaryl, C₁-C₆ alkylC(0)heteroaryl, indanyl,

763

 C_1 - C_6 alkylindanyl, C(0)indanyl, C_1 - C_6 alkylC(0)indanyl, cycloalkyl;

or D and E combine to form indanyl, fluorenyl, or cycloalkyl;

G is hydrogen, C_1 - C_6 alkyl, aryl, C_1 - C_6 alkylaryl, and C_1 - C_6 alkenyl;

J is hydrogen, $C_1\text{-}C_6alkyl,$ aryl, and $C_1\text{-}C_6alkylaryl;$

L is hydrogen, C₁-C₆alkyl, C(0)OC₁-C₆alkyl, aryl,

10 C_1-C_6 alkylaryl, $C(0)OC_1-C_6$ alkylaryl, C_1-C_6 alkenyl, -F, and -CN, C_1-C_6 alkyl-OH, C_1-C_6 alkyl-O- C_1-C_6 alkyl-C(0)R₆;

or a pharmaceutically acceptable salt or solvate thereof.

- 15 14. A method according to Claim 11 wherein said patient is a human.
- 15. A method according to Claim 1 which further comprises the administration of an effective amount of an angiotensin converting enzyme (ACE) inhibitor.
 - 16. A method according to Claim 1 which further comprises the administration of an effective amount of an antihypertensive agent.

25

- 17. A method according to Claim 1 which further comprises the administration of an effective amount of a diuretic.
- 30 18. A method according to Claim 1 which further comprises the administration of an effective amount of a β -blocker.

764

- 19. A method according to Claim 11 which further comprises the administration of an effective amount of an angiotensin converting enzyme (ACE) inhibitor.
- 5 20. A method according to Claim 11 which further comprises the administration of an effective amount of an antihypertensive agent.
- 21. A method according to Claim 11 which further
 10 comprises the administration of an effective amount of a diuretic.
- 22. A method according to Claim 11 which further comprises the administration of an effective amount of a $\beta-$ blocker.
- 23. A method of diminishing loss of body weight in a mammal following congestive heart failure which comprises the administration of a therapeutically effective amount of GRP-2.
 - 24. A method of enhancing recovery of a mammal following congestive heart failure which comprises the administration of a therapeutically effective amount of GRP-2.

25

30

25. A method for screening compounds having growth hormone secretagogue activity for the ability to modulate cardiac function, which comprises the administration of said compounds to a SHHF/Mcc- fa^{CP} rat, wherein said rat exhibits congestive heart failure.




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6






FIG. 7



.8/8

FIG. 8



INTERNATIONAL SEARCH REPORT

International application No. PCT/US98/17201

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) :A61K 38/00, 31/54, 31/535, 31/495, 31/445 US CL :514/16, 222.2, 231.2, 249, 255, 315 According to International Patent Classification (IPC) or to both national classification and IPC B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/16, 222.2, 231.2, 249, 255, 315 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category* Citation of document, with indication, where app	propriate, of the relevant passages	Relevant to claim No.
X,P US 5,721,250 A (MORRIELLO et a column 43, line 41.	US 5,721,250 A (MORRIELLO et al.) 24 February 1998, see column 43, line 41.	
A,P US 5,663,146 A (BOWERS et al.) 02	September 1997.	1-25
Further documents are listed in the continuation of Box C. See patent family annex.		
T bater document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention to be of particular relevance; the claimed invention cannot be considered novel or cannot be considered to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published after the international filing date when the document is taken alone document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family Date of the actual completion of the international search 27 NOVEMBER 1998 Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT **C** *** **T** **		
Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	RAYMOND J. HENLEY III Telephone No. (703) 308-1235	for for